

Effect of Sewage on Physical and Chemical Properties of Soil in Unnao District (U.P.)

Abstract

This study was carried out during 2021-22 considering various vegetable growing farmers in Unnao District (U.P.) to find out the effect of using sewage as the only source of irrigation to their crops. 250 surface soil samples collected from farmer's field were analyzed for Physical as well as Chemical parameters. The physical properties of soils improved due to sewage application; Bulk density ranged from 1.04 to 1.42 Mg m⁻³ and particle density ranged between 2.61 to 2.88 Mg m⁻³ and Porosity ranged between 53.4 to 61.85 with mean of 56.87 percent. Chemical Parameters such as pH, E.C., O.C., Available N, P & K were also analyzed and the results indicated that soils had a mean normal pH of 7.2, E.C. of 0.76 dSm⁻¹, O.C. of 56.80. Available nitrogen ranged from 565 – 1217 kg ha⁻¹ with mean of 879.56 kg ha⁻¹. The Phosphorus availability was found ranging from 14.40 – 33.5 kg ha⁻¹ with a mean of 23.98 kg ha⁻¹, while Potassium was found with a mean of 205.30 kg ha⁻¹ in range of 174 – 235 kg ha⁻¹.

Key words: Bulk density, Particle density, pH, Nitrogen, Phosphorous and Potassium

INTRODUCTION It is necessary to do a more updated review of the introductory part, there are few recent articles, the most current is from 2004. In addition, the introduction is very poor.

Water and nutrients are the major inputs for crop production. It is predicted that most of the Asian countries will face severe problem related to water availability by 2025 (Singh, 1999). With the rapid growth in industrialization in different countries, the pollution problem is also on the increase, which have resulted in utilization of significant quantities of fresh water available for agriculture. Application of Sewage water to agricultural soil is a useful source of plant nutrients, particularly nitrogen and phosphorous and also organic matter that can potentially improve soil fertility and physical properties. However, in addition to these beneficial effects, effluents often contain appreciable amounts of both organic and inorganic toxic materials. Many organic pollutants, being biodegradable, are less persistent, and presumably have transient and less serious effects as they are eventually metabolized to carbon dioxide and other inorganic substances.

In many areas of developing countries, untreated wastewater flows through channels into rivers where it is diverted by subsistence farmers to small plots of vegetables including tomato, cabbage, beetroot and others which are easily consumed as salad. The public health risks of using such contaminated streams for irrigation are obvious (Mead and Griffin, 1998; WHO 2004). However, treated effluents can be used for irrigation under controlled conditions to minimize the transfer of pathogenic and toxic contaminants into agricultural products, soil surfaces and groundwater (Batarseh *et al.* 1989).

Since, industrial effluent is a pool of various elements, it may be beneficial in enhancing the fertility status of soil as it contains appreciable amount of essential plant nutrients that are taken up by crops during plant growth and development. Amount of available nitrogen, phosphorus and other nutrients also increase in the soil but excess of them can leach and pollute groundwater under continuous effluent use for long periods (Chaney, 1990).

Materials and methods

In the materials and methods part, it lacks a lot of information; It does not indicate how the sample was taken, from how many hectares, from how many agricultural farms, whether there are repetitions or not, what methodology was used for the measurements in the laboratory, how many repetitions were made, in addition to the fact that it is necessary to clarify what statistical methods were used. .

It is also not clear how long the soils have been irrigated, what type of water is used or what its origin is since they mention different sources of different types of water and in which places; The information they provide is very poor, with this information it is impossible to repeat this study, the materials and methods part is extremely vague. On the other hand, it is not noted if there are some samples that have not been watered and that serve as a comparison for the results section. In the materials and methods part, it lacks a lot of information; It does not indicate how the sample was taken, from how many hectares, from how many agricultural farms, whether there are repetitions or not, what methodology was used for the measurements in the laboratory, how many repetitions were made, in addition to the fact that it is necessary to clarify what statistical methods were used.

It is also not clear how long the soils have been irrigated, what type of water is used or what its origin is since they mention different sources of different types of water and in which places; The information they provide is very poor, with this information it is impossible to repeat this study, the materials and methods part is extremely vague. On the other hand, it is not noted if there are some samples that have not been watered and that serve as a comparison for the results section.

One of the main industrial cities near Kanpur is Unnao, which is home to the majority of the cotton, leather, pharmaceutical, steel, and other industries. More than 50 industrial facilities, mostly tanneries, may be found in the Unnao industrial area, which is close to Kanpur on the northern bank of the Ganga River. The effluents released by the industries are ultimately dumped into the Ganga River after passing through a common effluent treatment plant with approximately 70% treating capacity. Unnao is a second township across the Ganga, located between the rivers Ganga and Sai, roughly 20 km from Jammu. The District is located between Latitudes 26°8' and 27°2' North and 80°3' and 81°3' east. It is divided from Kanpur and Fatehpur by the Ganga on the west, Lucknow on the east, Rae Bareli on the south, and Hardoi on the north. It covers 4589 sq. km of land.

In the present investigation total 250 soil samples (0-20cm) were collected from farmer's field with the help of tube auger randomly from field during *rabi* season nearby sewage outlet who engage in irrigating their crops with industrial effluents.

Result and discussion:

In the results section they cannot point out that something went down or up because there is no way to compare.

The bulk density of soils irrigated with sewage was decreased due to addition of organic matter through sewage irrigation. Particle density was not changed in sewage irrigated soil (Mathan, 1994). This can be attributed to improvement in total porosity and aggregate stability in the sewage irrigated soils due to addition of organic matter which plays an important role in improving soil physical environment. Rattan *et al.* (2001) observed enhanced available water content in the soils due to continuous application of sewage. Rai *et al.* (2011).

1.24	1.13	1.17	1.09	1.17	2.62	2.65	2.64	2.63	2.61	54.28	55.23	56.78	56.28	53.45
1.37	1.28	1.25	1.17	1.25	2.78	2.82	2.72	2.77	2.83	56.75	56.78	58.32	61.85	58.63
1.29	1.15	1.38	1.13	1.13	2.65	2.68	2.86	2.82	2.77	55.39	54.35	55.89	58.63	54.89
1.33	1.32	1.22	1.08	1.28	2.71	2.77	2.77	2.69	2.85	54.71	55.78	56.43	59.58	55.36
1.26	1.17	1.35	1.15	1.16	2.83	2.63	2.69	2.85	2.81	55.84	58.42	58.97	57.25	53.92
1.38	1.24	1.19	1.04	1.21	2.77	2.85	2.74	2.71	2.68	57.28	56.29	56.45	60.69	57.18
1.42	1.37	1.24	1.11	1.18	2.85	2.79	2.81	2.88	2.72	56.19	58.85	57.83	56.93	53.88
1.31	1.29	1.32	1.08	1.24	2.81	2.81	2.78	2.64	2.85	54.23	54.78	56.79	61.44	56.25
1.38	1.25	1.18	1.23	1.19	2.78	2.74	2.75	2.82	2.69	57.85	57.63	55.43	58.29	58.54
1.35	1.18	1.26	1.16	1.23	2.69	2.86	2.68	2.85	2.65	58.03	56.45	57.28	57.65	57.92

Table-2: Effect of sewage on chemical properties of soil

pH					Electrical Conductivity (dS/m)					Organic Carbon (%)				
E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅
7.24	7.42	7.06	7.04	6.76	0.92	0.73	0.52	0.48	0.63	54.39	53.28	55.39	54.78	53.85
7.38	7.55	7.13	7.13	6.92	1.15	0.88	0.68	0.55	0.77	56.82	55.42	58.74	58.35	58.62
7.33	7.63	7.08	7.09	6.85	0.98	0.76	0.56	0.43	0.64	55.27	53.85	56.45	61.54	55.48
7.29	7.49	7.11	7.22	6.98	1.06	0.91	0.63	0.68	0.71	58.43	54.69	59.62	59.37	57.95
7.42	7.52	7.15	7.17	6.83	1.13	0.85	0.59	0.52	0.69	54.78	55.74	57.28	55.82	54.23
7.38	7.68	7.09	7.08	6.95	0.98	0.79	0.64	0.49	0.75	56.19	56.22	55.83	60.45	56.89
7.27	7.47	7.18	7.21	6.88	1.09	0.84	0.77	0.61	0.78	58.95	54.85	57.58	56.28	58.26
7.35	7.54	7.05	7.16	6.94	1.12	0.93	0.73	0.57	0.65	57.63	56.69	56.94	59.84	57.35
7.43	7.61	7.13	7.25	6.79	0.97	0.89	0.58	0.54	0.73	54.78	53.95	58.26	61.53	54.87
7.37	7.58	7.17	7.19	6.87	1.14	0.95	0.65	0.66	0.71	56.25	54.33	59.44	60.79	55.63

Table-3: Effect of sewage on chemical properties of soil

Available Nitrogen	Available Phosphorus	Available potassium
---------------------------	-----------------------------	----------------------------

(kg/ha)					(kg/ha)					(kg/ha)				
E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅	E ₁	E ₂	E ₃	E ₄	E ₅
653	565	613	876	945	22.8	24.6	17.4	14.4	18.8	222	213	174	183	198
678	574	687	915	1027	28.3	29.4	22.7	20.9	21.5	228	217	188	198	205
694	637	807	943	1056	27.5	31.9	21.5	17.3	24.9	225	225	183	185	199
686	702	698	967	978	24.9	32.5	19.3	18.8	19.3	224	219	179	193	203
702	757	836	989	1013	26.4	28.3	24.8	21.5	22.7	229	221	185	189	201
679	782	723	992	1105	28.8	32.8	19.5	19.2	28.1	234	215	178	197	194
713	795	756	1023	1178	25.5	27.2	20.7	21.6	26.8	231	228	181	191	208
708	803	654	978	1094	29.2	33.5	23.2	18.4	25.4	227	232	186	188	211
725	727	845	1015	1203	27.8	25.8	18.8	16.7	27.7	233	235	175	192	197
668	764	789	984	1217	23.54	30.7	21.5	20.5	23.5	227	229	188	198	204

Conclusion:

Continuous use of sewage irrigation recorded improvement in soil physical properties like bulk density, water retention, organic carbon and build-up of soil available N, P, K, and micronutrient status. The industrial sewage irrigated soils had higher electrical conductivity and organic carbon as compared to the ground water alternate irrigated soils. The electrical conductivity although increased due to effluent/sewage, it was within the tolerance limit to cause any soil salinity hazard. This warrants the potential hazard to soil and plant health suggesting necessity of their safe use after pre-treatment in order to make use of sewage as a cheap potential alternative source of plant nutrients in agriculture.

Reference:

There are only 10 citations in the manuscript.

Only the references marked in yellow are citations, the rest are not.

Of the 19 references used, 11 are from the year 2000 or more and only two are from the year 2020 or more.

- Akan, J.C., Abdulrahman, F.I., Sodipo, O.A. and Lange, A.G. (2010).** Physico-chemical Parameters in Soil and Vegetable Samples from Gongulon Agricultural Site, Maiduguri, Borno State, Nigeria., *Journal of American Science* 6(12).
- Amusan, A.A., Ige, D.V. and Olawale, R. (2005).** Characteristics of soils and crops uptake of metals in municipal waste dump sites in Nigeria. *Journal of Human Ecology*, 17: 167-171.
- Awomesoa, J.A., Ufoegbunea, G.C., Oluwasanyaa, G.O. and Ademola-Aremua, O.O. (2009).** Impact of industrial effluents on water, soils and plants in the Alakia industrial area of Ibadan, South West Nigeria. 91: 5-15.
- Batarseh, L. I. O. A., Rimavi, E. Salameh, (1989).** Treated wastewater reuse in agriculture. Part1. Hussein Medical Center Project. The Water Research and Study Center, University of Jordan, Issue no. 12.
- Bhise, P. M., Cholla, O. and Venugopalan, M. V. (2007).** Effect of waste water irrigation on soil properties under different land system. *Journal of the Indian Society Soil Science*, 55: 254-258.
- Brar, M.S., Maltti, S.S., Singh, A.P., Arora, C.L. and Gill, K.S. (2000).** Sewage water irrigation effects on some potentially toxic trace elements in soil and potato plants in Norton western India. *Canadian Journal of Soil Science*, 80: 465-471.
- Chaney, R. L., (1990).** Twenty years of land application research. *Biocycle*, 31:54-59.
- Chojnacka, K., Witek-Krowiak, A., Moustakas, K., Skrzypczak, D., Mikula, K., Loizidou, M. (2020).** A transition from conventional irrigation to fertigation with reclaimed wastewater: Prospects and challenges. *Renewable and Sustainable Energy Reviews* 130 109959. <http://www.elsevier.com/locate/rser>.
- Gupta, A.P., Narwal, R.P., Antil, R.S. (1999).** Sewer water composition and its effect on soil properties. *Bioresource Technology*, 65: 171-173.
- Jackson, M. (1973).** "Soil chemical analysis Prentice hall Inc. Englewood Cliffs" New Jersey, U.S.S.
- Kharche, V.K., Desai, V.N. and Pharande, A.L. (2011).** Effect of Sewage Irrigation on Soil Properties, Essential Nutrient and Pollutant Element Status of Soils and Plants in a Vegetable Growing Area around Ahmednagar City in Maharashtra. *Journal of the Indian Society of Soil Science*, 59:177-184.

Kumar, Vipin and Nagar, Nidhi.(2012). Effect of industrial effluent on content of nutrient and heavy metals in plant and soil. Proc. International Conference on Chemical, Bio-chemical and Environmental Sciences (ICBEE'2012) December 14-15, 2012 Singapore.

Kurana 1994?

El-Bassi, L., Azzaz, A. A., Jellali, S., Akrou, H., Marks, E. A., Ghimbeu, C. M., & Jeguirim, M. (2021). Application of olive mill waste-based biochar's in agriculture: Impact on soil properties, enzymatic activities and tomato growth. *Science of the total environment*, 755, 142531.

Malla, R. and Totawat, K. L. (2006). Effect of municipal sewage on soil properties and chemical build up in vegetable crops grown on Haplusteps of sub humid southern plains of Rajasthan. *Journal of the Indian society of Soil Science*, 54: 226-231.

Mathan 1994?

Mead, P. S., and P. M. Griffin, (1998). Escherichia coli 0157: H7. *Lancet* 352, 1207-1212.

Mitra, A. and Gupta, S.K. (1999). Effect of sewage irrigation on essential plant nutrient and pollutant status in a vegetable growing area around Calcutta. *Journal of the Indian Society of Soil Science*, 47: 99-105.

Rai 2001?

Rattan 2001?

Rattan. R. K., Datta, S. P. Chonkar, P. K. Suribabu, K. and Singh, A. K. (2005). Long term impact of irrigation with sewage effluents on heavy metal content in soils, crops and ground water - a case study, *Journal of Agric. Ecosystems Environ.*, 109 : 310-322.

Singh, P.K., Deshbhratar, P.B., Ramteke, D.S. (2012). Effects of sewage wastewater irrigation on soil properties, crop yield and environment. *Agricultural Water Management*, 103: 100-104.

Singh, R. Sand, R. P. Singh, (1999). Distribution of DTPA extractable Cd, Pb, Cr, Cu, Zn, Mn and Fe in soil profiles contaminated by sewage and industrial effluent. *J. Indian Soc. Soil. Sci.* 42(3):466-468.

WHO 2004?