

## **Effect of pre-harvest fruit bagging on fruit quality characteristics of different guava cultivars during rainy season**

### **ABSTRACT**

The present investigation was carried out on 6-7 years old grafted guava plants at Guava Demonstration Centre, Bhuna, Haryana, India during the rainy season (2019-20). During the experiment, the fruits of different guava cultivars viz. Hisar Safeda, Hisar Surkha, Allahabad Safeda, and Shweta were bagged with different bagging materials immediately after the fruit set. Results showed that the fruits bagged with blue polythene exhibited maximum fruit volume (113.96 ml) and organoleptic score (8.8), while the minimum (95.30 ml and 6.3, respectively) values were observed in the control. The highest fruit firmness (6.62 Kg/cm<sup>2</sup>) was observed with green polythene, while the minimum (5.03 Kg/cm<sup>2</sup>) was in control. Additionally, bagged fruits showed better peel colour development. The fruit drop was also reduced in all treatments except for the control (20%), yellow paper (12.78%), and green paper (11.66%). Fruit drop was minimum in transparent polythene (5.55%) and green polythene. On the other hand, Decay loss was eliminated in yellow paper, green paper, blue paper, cotton cloth, and muslin cloth bags, and significantly reduced in all the treatments except the control (8.88%). Among the cultivars, the maximum fruit firmness (5.98 Kg/cm<sup>2</sup>) was recorded in Hisar Safeda, whereas the least firmness (5.64 Kg/cm<sup>2</sup>) was in Allahabad Safeda. In conclusion, pre-harvest bagging of fruits was found promising in enhancing the quality attributes of various guava cultivars which can also be applied in real farm situations to enhance the yield as well as the quality of guava crop.

**Keywords:** Cloth bags; Infected fruits; Polythene bags; Paper bags

### **1. Introduction**

Guava (*Psidium guajava* L.), a member of the family **Myrtaceae**, is also known as ‘Apple of

tropics' as its nutritive value is equivalent to that of an apple. It is widely grown in the tropical and subtropical regions of the world. It is one of the most common and well-distributed fruit crops in the Indian subcontinent. Concerning its area and production, it ranks next to mango, banana and citrus. During its growth and developmental periods, the fruits became highly susceptible to insect-pest infestations, diseases, birds as well as mechanical damage, resulting in the reduction of their commercial value and ultimately causing significant economic losses (Sharma et al., 2014).

Under North-Indian conditions, guava produces two crops *i.e.*, rainy and winter season crops annually. The winter season crop is generally, superior in quality to the rainy season crop as it is severely infested with the fruit fly (*Bactrocera correcta* Bezzi), anthracnose (*Colletotrichum gloeosporioides* Penz.) disease, and bird's attacks (Satarkar et al., 2009; Vargas et al., 2015). In the month of April, the population of fruit flies starts to build up in the orchards and continues up to November in different fruit crops. Due to the presence of a wide range of hosts, a large number of generations per year, high mobility of adults, and high fecundity; their management become very difficult in the orchards. Moreover, all the developmental stages of fruit flies are unexposed *viz.* eggs and larvae are present inside the fruits, whereas, pupae are buried in the soil (Brar et al., 2019; Parkash et al., 2021). Therefore, the application of insecticides does not prove effective against fruit flies. Hence, bagging is considered an eco-friendly alternative, which can act as a physical barrier between fruit and the environment, subsequently protecting the fruit from fruit flies, diseases, mechanical damage, sunburning as well as bird damage (Hofman et al., 1997; Sharma et al., 2014; Bishnoi et al., 2023). This technique generally applied to many fruits improves the visual quality of fruits not only through the promotion of colour, however, also through the reduction of the incidence of cracking and russeting (Sharma et al., 2014). This technique confers the bagging of individual fruit or a fruit bunch on the tree for a specific

period. Bagging beneficially changes the micro-environment inside the bag which also exerts a beneficial impact on fruit size and internal fruit quality (Fan and Mattheis, 1998; Sharma et al., 2017, Sharma et al., 2020). However, the selection of bagging material and time of fruit bagging is of utmost importance for getting better results (Liu et al., 2004; Lu et al., 2008). Thus, the present investigation was conducted to evaluate the use of different bagging materials on various quality parameters of guava fruits.

## 2. MATERIAL AND METHODS

### 2.1 Experimental site

The present study was carried out at the farms of the Guava Demonstration Centre, Bhuna, Haryana, India. Which is situated at 29° 32' latitude, 75° 42' and longitudes 222 m above mean sea level. The temperature varies from 40 °C to 48 °C during summer to as low as to freezing point accompanied by chilling frost in winter. The approximate average rainfall is 450 mm, most of which is received mainly during the South-West monsoon (i.e., July to September), while few showers also occur from December to February due to western disturbances. The physiological analysis of fruits was done at PG-Lab (Department of Horticulture), CCS Haryana Agricultural University, Hisar.

### 2.2 Experimental details

In the present investigation, well-grown grafted plants of 4 different cultivars viz. **Hisar Safeda, Hisar Surkha, Allahabad Safeda** and **Shweta** were selected for the experiment. All the selected plants were of the same size and vigour with an age of 6 to 7 years old, planted at a geometry of 3 x 6 m on raised bunds with drip irrigation facilities. Trees were well maintained, regularly pruned and subjected to the recommended cultural practices i.e., irrigation, fertilization, weeding, insect pest and disease management.

Fifteen uniform-sized fruits were bagged immediately after fruit set each on three plants (replications) with different materials as per treatment schedule (Table 1). Five fruits from

each plant, *i.e.*, 15 fruits (from 3 replications) were harvested from each cultivar for each treatment and subsequently analysed for different parameters as mentioned in the next section. The harvesting time of fruits among the treatments was same, however, varied among cultivars.

**Table 1: Treatments details**

<b>Treatments</b>	<b>Description</b>
T <sub>1</sub>	White paper bag
T <sub>2</sub>	Red paper bag
T <sub>3</sub>	Yellow paper bag
T <sub>4</sub>	Green paper bag
T <sub>5</sub>	Blue paper bag
T <sub>6</sub>	Brown paper bag
T <sub>7</sub>	Newspaper bag
T <sub>8</sub>	Transparent polythene bag
T <sub>9</sub>	White polythene bag
T <sub>10</sub>	Pink polythene bag
T <sub>11</sub>	Yellow polythene bag
T <sub>12</sub>	Green polythene bag
T <sub>13</sub>	Blue polythene bag
T <sub>14</sub>	Cotton cloth bag
T <sub>15</sub>	Muslin cloth bag
T <sub>16</sub>	Control (unbagged)

## **2.3 Evaluation of physiological parameters**

### **2.3.1 Fruit Volume (ml)**

Fruit volume was measured by the water displacement method in which the fruits were

dipped in a calibrated measuring cylinder filled with water and an increase in the water level after dipping of the fruit was recorded and the values were expressed as milliliters.

### 2.3.2 Fruit firmness (kg/cm<sup>2</sup>)

A penetrometer with a cylindrical plunger probe of 4 mm was used for measuring the fruit's firmness. It is measured by making a puncture on the fruit's stem end and expressed in kg/cm<sup>2</sup>.

### 2.3.3 Decay loss (%)

The decay of fruits was decided based on visual observations. Decay loss was evaluated by simply counting the number of spoiled fruits displaying fungal mycelia or sporulation and expressed as per cent spoilage of fruits.

$$\text{Decay loss (\%)} = \frac{\text{Number of spoiled fruits}}{\text{Total number of fruits}} \times 100$$

### 2.3.4 Fruit drop (%)

The fruit drop percentage was calculated by counting the number of bagged fruit present on the plant at the time of harvest. Fruit drop (%) was calculated by the following formula:

$$\text{Fruit drop (\%)} = \frac{\text{Initial number of fruits bagged} - \text{Bagged fruit remained on plant}}{\text{Initial number of fruits bagged}} \times 100$$

### 2.3.5 Organoleptic quality

The fruits were evaluated for organoleptic quality based on their appearance, taste, mouthfeel, aroma, and overall acceptability by a panel of 10 semi-trained members (5 males and 5 females, 23-45 years old) from the Department of Horticulture, CCSHAU, Hisar, India using a 9-point hedonic scale. On the scale, 9 stands for Liked extremely and 1 stands for dislike extremely (Ranganna, 1977).

**Table 2: The ranking based on 9 points hedonic rating scale is given below:**

Acceptability	Marks
Extremely desirable	9
Very much desirable	8
Moderately desirable	7
Slightly desirable	6
Neither desirable nor undesirable	5
Slightly undesirable	4

### 2.3.6 Fruit colour

Colour of the fruit was observed visually by a team of 10 judges. The colour observations *i.e.*, greenish yellow, yellowish green, light yellow and light green were recorded.

### 2.4 Statistical analysis

The data were statistically analyzed in Randomized Block Design using SPSS software (IBM, SPSS Inc., USA). The p-values were calculated and the results were expressed as CD at 5% level of significance. Data were subjected to two-way ANOVA.

## 3. RESULTS AND DISCUSSION

### 3.1 Fruit volume (ml)

It is evident from Table 3 that different treatments influenced the fruit volume significantly. The maximum volume was observed in the fruits bagged with blue polythene, it was statistically at par with fruits bagged in green polythene (112.63 ml), cotton cloth (111.7 ml) and yellow polythene (109.32 ml), whereas the minimum fruit volume (95.3 ml) was in unbagged (control) fruits, it was statistically at par with fruit bagged in pink polythene (96.52 ml), white polythene (99.98 ml) and white paper (100.38 ml). The maximum fruit volume (127.81 ml) among the cultivars was observed in fruits of Shweta as compared to Hisar Surkha which had minimum fruit volume (92.28 ml). The improved fruit volume might be due to increased relative humidity and development of a favorable microclimate inside the

bags, which resulted in better growth and development of fruits (Jat, 2019). These results were in close conformity with the earlier findings of (Khalil Omar et al., 2014) in bagged date palm fruits.

### **3.2 Fruit Firmness (Kg/cm<sup>2</sup>)**

The data pertaining to fruit firmness is presented in Table 3 showed that the maximum fruit firmness was observed in fruits bagged in green polythene bags (6.62 Kg/cm<sup>2</sup>) as compared to the control (5.03 Kg/cm<sup>2</sup>), it was statistically at par with fruits bagged in blue polythene bag (5.15 Kg/cm<sup>2</sup>). Among the cultivars, the maximum firmness was reported in fruits of Hisar Safeda (5.98 Kg/cm<sup>2</sup>), while the minimum firmness was in fruits of Allahabad Safeda (5.64 Kg/cm<sup>2</sup>). Breakdown of starch, cell wall or reduction in the middle lamella by solubilization of pectic substances leads to a reduction in fruit firmness. Whereas, increased fruit firmness might be due to the better stability of the cell wall, middle lamella and increase in pectin content under bagging conditions (Jat, 2019). The results of the present study were in line with (Bentley and Viveros, 1992) in bagged 'Granny Smith' apples and (Sharma et al., 2013) in bagged 'Delicious' apples.

### **3.3 Decay Loss (%)**

The data on infected fruits, as influenced by different bagging treatments in different cultivars, have been presented in Table 4. The revealed that the infected fruits (decay loss) were observed at maximum (8.88%) in unbagged fruits, whereas it had been completely controlled in the fruits bagged with yellow paper or green paper or blue paper or cotton cloth and muslin cloth, it was statistically at par with fruits bagged in other bagging materials *i.e.*, white paper (0.56%), red paper (0.56%), brown paper (0.56%), newspaper (0.56%), yellow polythene (0.56%), blue polythene (1.11%) and pink polythene (1.67%). As per previous documentation, bagging maintains a physical barrier between fruits and the environment, leading the fruits to be free from damage by fruit flies, birds, pests and

diseases, which results in the reduction of infected fruits (Sharma et al., 2014). Similar results were obtained by (Behera and Pathak, 2015) and (Sharma et al., 2020) in guava fruits.

### **3.4 Fruit drop (%)**

The fruit drop varied significantly with different treatments as shown in Table 4. The maximum fruit drop (20%) was observed in unbagged fruits, whereas the minimum fruit drop was observed in fruits bagged with transparent polythene and green polythene (5.55%), it was at par with fruits bagged in blue polythene bags (6.11%), muslin cloth bags (6.66%), cotton cloth bags (7.22%), newspaper bags (8.33%), red paper bags (8.88%), white paper bags (9.44%), blue paper bags (9.44%), brown paper bags (9.44%), white polythene bags (9.44%), pink polythene bags (9.44%) and yellow polythene bags (10%). It might be due to perforations made in these bags allowing free circulation of air and did not allow to build up an excess temperature and relative humidity in bags (Sharma et al., 2020). The results of the present study were in agreement with (Islam et al., 2017; Kireeti et al., 2018) in bagged mango fruits.

### **3.5 Organoleptic score**

Organoleptic score varied significantly among the treatments and the highest organoleptic score was assigned to fruits bagged in blue polythene bags (8.8), which was statistically at par with the fruits bagged in cotton cloth (8.5) and green polythene (8.5) as compared to the minimum in fruits of control (6.3) (Table 5). Among various cultivars, the organoleptic score was found non-significant, while the interaction was found significant. It might be due to the fact that pre-harvest bagging improves the internal fruit quality by reducing blemishes and micro-environment alterations within fruits during fruit development phase (Sharma et al., 2014). Similar results were also obtained by (Abbasi et al., 2014) in bagged guava and (Islam et al., 2019) in mango fruits.

### **3.6 Fruit colour**

Data presented in Table 5 showed that fruits bagged with blue paper, yellow polythene, green polythene, blue polythene, and cotton cloth showed better peel colour (Light green and light yellow) development as compared to the other treatments. The peel colour of fruit has a close relationship with the sensitivity of fruits to light and the rate of anthocyanin biosynthesis in fruit peel during fruit development phase. Bagging modifies the light quality and quantity inside the bag which might result in better peel colour development (Zhang et al., 2015). Similar results were obtained by (Singh et al., 2007) in bagged guava, (Hofman et al., 1997; Watanawan et al., 2008) in mango and (Signes et al., 2007) in 'Perla' grapes.

### **4. Conclusion**

The study demonstrated that pre-harvest bagging of guava fruits with different materials can significantly improve the quality attributes of various guava cultivars. The use of blue polythene bags resulted in the highest fruit volume and organoleptic score, while green polythene bags exhibited the highest fruit firmness. The bagged fruits also showed better peel color development, reduced fruit drop, and decay loss. These findings suggest that pre-harvest bagging can be an effective technique to enhance the yield and quality of guava crops, and can be practically applied in real farm situations. However, further research is needed to explore the long-term effects of pre-harvest bagging on the guava crop and its economic feasibility.

### **References**

- Abbasi NA, Chaudhary MA, Ali MI, Hussain A, and Ali I 2014. On tree fruit bagging influences quality of guava harvested at different maturity stages during summer. *International Journal of Agriculture and Biology* **16**, 543–549.
- Behera SD and Pathak S 2015. Pre-Harvest Treatments for Fruit Quality Improvement in

Rainy Season Guava (*Psidium guajava*). *International Journal of Science and Research* **6**, 1358–1361.

Bentley W and Viveros M 1992. Brown-bagging Granny Smith apples on trees stops codling moth damage. *California Agriculture* **46**, 30–32.

Bishnoi M, Goyal RK, Kumar A, Vats C, Bhawna, Kumar M and Kumar A 2023. Effect of Preharvest Fruit Bagging on Physical Parameters of different Guava Cultivars in Rainy Season Crop. *Indian Journal of Ecology* **50**(5), 1452-1456

Brar JS, Arora NK, Kaur K, Kaur G, Gill KS, and Gill MIS 2019. Fruit Bagging for Improving Quality of Rainy Season Guava Under Punjab Conditions. *Agricultural Research Journal* **56**, 475–479.

Fan X and Mattheis J 1998. Bagging “Fuji” Apples during Fruit Development Affects Color Development and Storage Quality. *HortScience* **33**, 1235–1238.

Hofman PJ, Smith LG, Joyce DC, Johnson GI, and Meiburg GF 1997. Bagging of mango (*Mangifera indica* cv. 'Keitt') fruit influences fruit quality and mineral composition. *Postharvest Biology and Technology* **12**, 83–91.

Islam MT, Rahman MS, and Shamsuzzoha M 2017. Influence of pre-harvest bagging on fruit quality of Mango (*Mangifera indica* L.) cv. Mishribhog. *International Journal of Biosciences* **11**, 59–68.

Islam MT, Shamsuzzoha M, Bari MA, Rahman MS, Aktar M, Islam M, and Rahman MA 2019. Influence of bagging time on fruit quality and shelf life of mango (*Mangifera indica* L.) Cv. Amrapali in Bangladesh. *International Journal of Agriculture and Environmental Research* **5**, 412–423.

Jat ML 2019. Effect of pre-harvest fruit bagging on maturity and physico-chemical properties of guava fruits cv.VNR Bihi. G. B. Pant University of Agriculture and Technology, Pantnagar.

- Khalil Omar AED, Al-Saif AM, and El-Aziz Ahmed MA 2014. Bagging of bunches with different materials influences yield and quality of Rothana date palm fruit. *Journal of Food, Agriculture and Environment* **12**, 520–522.
- Kireeti A, Haldankar P, and Parulekar Y 2018. Studies on effect of types of bag on mango fruit (Cv. Kesar) at egg stage. *International Journal of Chemical Studies* **6**, 1–4.
- Liu YJ, Xu JH, Zhang ZH, Jiang JM, and Yu D 2004. Effects of different paper bags on fruit quality of loquat. *Acta Agriculturae Universitatis Jiangxiensis* **26**, 334–337.
- Lu X, Zeng L, and Lu YY 2008. Study on optimal time for controlling *Bactrocera dorsalis* Hendel in carambola orchard. *Journal of Environmental Entomology* **30**, 214–219.
- Parkash J, Singh B, and Yadav SS 2021. Bio-ecological management of melon fruit fly, *Bactrocera cucurbitae* Coquillett on bottle gourd. *Journal of Entomological Research* **45**(suppl), 937-939.
- Ranganna S 1977. Manual analysis of fruit and vegetable product. Tata McGraw-Hill, New Delhi, New Delhi.
- Satarkar VR, Krishnamurthy S V, Faleiro JR, and Verghese A 2009. Spatial distribution of major *Bactrocera* fruit flies attracted to methyl eugenol in different ecological zones of Goa, India. *International Journal of Tropical Insect Science* **29**, 195–201.
- Sharma R, Ram A, Sagar V, and Sethi S 2017. Influence of pre-harvest fruit bagging on Golden Delicious apple (*Malus × domestica*). *Indian Journal of Agricultural Sciences* **87**, 1170–1173.
- Sharma RR, Nagaraja A, Goswami AK, Thakre M, Kumar R, and Varghese E 2020. Influence of on-the-tree fruit bagging on biotic stresses and postharvest quality of rainy-season crop of ‘Allahabad Safeda’ guava (*Psidium guajava* L.). *Crop Protection* **135**.
- Sharma RR, Pal RK, Asrey R, Sagar VR, and Rana MR 2013. Pre-harvest fruit bagging influences fruit color and quality of apple cv. Delicious. *Agricultural Sciences* **4**, 443–448.

- Sharma RR, Reddy SVR, and Jhalegar MJ 2014. Pre-harvest fruit bagging: A useful approach for plant protection and improved post-harvest fruit quality - A review. *Journal of Horticultural Science and Biotechnology* **89**, 101–113.
- Signes AJ, Burló F, Martinez-Sanchez F, and Carbonell-Barrachina AA 2007. Effects of preharvest bagging on quality of black table grapes. *World Journal of Agricultural Sciences* **3**, 32–38.
- Singh BP, Singh RA, Singh G, and Killadi B 2007. Response of bagging on maturity, ripening and storage behaviour of “winter guava.” *Acta Horticulturae* **735**, 597–601.
- Vargas RI, Piñero JC, and Leblanc L 2015. An overview of pest species of *Bactrocera* fruit flies (Diptera: Tephritidae) and the integration of biopesticides with other biological approaches for their management with a focus on the pacific region. *Insects* **6**, 297–318.
- Watanawan A, Watanawan C, and Jarunate J 2008. Bagging “Nam Dok Mai #4” mango during development affects color and fruit quality. *Acta Horticulturae* **787**, 325–328.
- Zhang BB, Guo JY, Ma RJ, Cai ZX, Yan J, and Zhang CH 2015. Relationship between the bagging microenvironment and fruit quality in ‘guibao’ peach [*Prunus persica* (L.) batsch]. *Journal of Horticultural Science and Biotechnology* **90**, 303–310.

**Table 3: Effect of pre-harvest fruit bagging on volume (ml) and firmness (Kg/cm<sup>2</sup>) of different guava cultivars in rainy season crop (2019-20)**

Sr. No	Treatments	Volume (ml)					Firmness (Kg/cm <sup>2</sup> )				
		Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta	Mean	Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta	Mean
T <sub>1</sub>	White paper bag	95.88	87.13	92.58	125.93	100.38	5.83	5.68	5.55	5.66	5.68
T <sub>2</sub>	Red paper bag	105.84	93.30	97.88	129.99	106.75	6.42	6.23	6.08	6.22	6.24
T <sub>3</sub>	Yellow paper bag	103.06	91.99	96.82	129.45	105.33	5.67	5.56	5.46	5.53	5.56
T <sub>4</sub>	Green paper bag	107.41	95.79	98.74	132.65	108.65	6.47	6.31	6.23	6.38	6.35
T <sub>5</sub>	Blue paper bag	109.96	97.56	100.26	127.27	108.76	5.98	5.75	5.61	5.74	5.77
T <sub>6</sub>	Brown paper bag	98.78	86.64	90.74	126.54	100.68	6.11	5.78	5.74	5.75	5.85
T <sub>7</sub>	Newspaper bag	98.37	88.11	93.78	122.08	100.59	5.47	5.29	5.18	5.35	5.32
T <sub>8</sub>	Transparent polythene bag	100.21	92.33	91.47	123.02	101.76	6.38	6.15	5.95	6.03	6.13
T <sub>9</sub>	White polythene bag	97.52	89.21	90.36	122.83	99.98	6.31	5.89	5.83	5.89	5.98
T <sub>10</sub>	Pink polythene bag	93.86	84.45	87.73	120.04	96.52	5.54	5.43	5.27	5.48	5.43
T <sub>11</sub>	Yellow polythene bag	108.85	97.33	99.47	131.63	109.32	6.61	6.44	6.29	6.49	6.46
T <sub>12</sub>	Green polythene bag	111.32	100.72	102.20	136.27	112.63	6.82	6.56	6.49	6.59	6.62
T <sub>13</sub>	Blue polythene bag	112.13	99.92	105.76	138.02	113.96	5.27	5.13	4.96	5.23	5.15
T <sub>14</sub>	Cotton cloth bag	111.94	97.45	103.56	133.84	111.70	5.42	5.23	5.06	5.11	5.21
T <sub>15</sub>	Muslin cloth bag	101.14	91.62	95.65	126.89	103.82	6.24	5.91	5.73	5.77	5.91
T <sub>16</sub>	Control (unbagged)	92.40	83.00	86.29	117.51	95.30	5.21	4.99	4.88	5.02	5.03
Treatment Mean		103.04	92.28	95.89	127.81		5.98	5.77	5.64	5.77	
C.D. (p=0.05)		Treatments (T) = 5.08, Cultivars (C) = 2.54, T x C = NS					Treatments (T) = 0.12, Cultivars (C) = 0.06, T x C = NS				

**Table 4: Effect of pre-harvest fruit bagging on decay loss (%) and fruit drop (%) of different guava cultivars in rainy season crop (2019-20)**

Sr. No	Treatments	Decay loss (%)					Fruit drop (%)				
		Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta	Mean	Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta	Mean
T <sub>1</sub>	White paper bag	2.22	0.00	0.00	0.00	0.56	8.88	8.89	8.88	11.11	9.44
T <sub>2</sub>	Red paper bag	0.00	0.00	2.22	0.00	0.56	8.88	11.11	6.66	8.88	8.88
T <sub>3</sub>	Yellow paper bag	0.00	0.00	0.00	0.00	0.00	13.33	13.33	11.11	13.33	12.78
T <sub>4</sub>	Green paper bag	0.00	0.00	0.00	0.00	0.00	13.33	11.11	11.11	11.11	11.66
T <sub>5</sub>	Blue paper bag	0.00	0.00	0.00	0.00	0.00	8.88	13.33	6.66	8.88	9.44
T <sub>6</sub>	Brown paper bag	0.00	0.00	2.22	0.00	0.56	6.66	11.11	6.66	13.33	9.44
T <sub>7</sub>	Newspaper bag	0.00	2.22	0.00	0.00	0.56	6.66	6.66	13.33	6.66	8.33
T <sub>8</sub>	Transparent polythene bag	2.22	2.22	2.22	2.22	2.22	4.44	4.44	6.66	6.66	5.55
T <sub>9</sub>	White polythene bag	4.44	6.66	2.22	2.22	3.89	13.33	6.66	8.88	8.88	9.44
T <sub>10</sub>	Pink polythene bag	2.22	0.00	0.00	4.44	1.67	11.11	4.44	13.33	8.88	9.44
T <sub>11</sub>	Yellow polythene bag	0.00	0.00	2.22	0.00	0.56	13.33	6.66	11.11	8.88	10.00
T <sub>12</sub>	Green polythene bag	0.00	2.22	4.44	2.22	2.22	6.66	4.44	4.44	6.66	5.55
T <sub>13</sub>	Blue polythene bag	2.22	2.22	0.00	0.00	1.11	4.44	6.66	6.66	6.66	6.11
T <sub>14</sub>	Cotton cloth bag	0.00	0.00	0.00	0.00	0.00	8.89	8.89	6.66	4.44	7.22
T <sub>15</sub>	Muslin cloth bag	0.00	0.00	0.00	0.00	0.00	4.44	6.67	8.88	6.66	6.66
T <sub>16</sub>	Control (unbagged)	8.88	11.11	8.88	6.66	8.88	20.00	15.55	22.22	22.22	20.00
Treatment Mean		1.39	1.67	1.53	1.11			8.75	9.58	9.58	
C.D. (p=0.05)		Treatments (T) = 2.18, Cultivars (C) = NS, T x C = NS					Treatments (T) = 4.68, Cultivars (C) = NS, T x C = NS				

**Table 5: Effect of pre-harvest fruit bagging on the organoleptic score and fruit colour of different guava cultivars in rainy season crop (2019-20)**

Sr. No	Treatments	Organoleptic score					Fruit Colour			
		Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta	Mean	Hisar Safeda	Hisar Surkha	Allahabad Safeda	Shweta
T <sub>1</sub>	White paper bag	6.8	6.9	6.5	7.2	6.9	Yellowish green	Yellowish green	Yellowish green	Yellowish green
T <sub>2</sub>	Red paper bag	7.5	7.9	7.9	7.8	7.8	Yellowish green	Yellowish green	Yellowish green	Yellowish green
T <sub>3</sub>	Yellow paperbag	7.8	7.5	7.9	7.8	7.8	Greenish Yellow	Yellowish green	Greenish Yellow	Yellowish green
T <sub>4</sub>	Green paper bag	8.1	7.9	7.2	8.1	7.8	Greenish Yellow	Yellowish green	Greenish Yellow	Yellowish green
T <sub>5</sub>	Blue paper bag	8.1	8.2	8.2	7.5	8.0	Light Yellow	Light green	Light Green	Greenish Yellow
T <sub>6</sub>	Brown paper bag	7.1	6.9	6.9	7.3	7.0	Greenish Yellow	Yellowish Green	Yellowish green	Yellowish green
T <sub>7</sub>	Newspaper bag	6.8	6.9	7.5	6.9	7.0	Yellowish green	Yellowish green	Yellowish green	Yellowish green
T <sub>8</sub>	Transparent polythene bag	7.5	7.5	7.2	6.5	7.2	Light Yellow	Yellowish green	Yellowish green	Yellowish green
T <sub>9</sub>	White polythene bag	6.8	7.1	6.9	6.6	6.9	Yellowish green	Yellowish green	Yellowish green	Yellowish green
T <sub>10</sub>	Pink polythene bag	6.8	6.9	6.9	6.2	6.7	Yellowish green	Greenish Yellow	Yellowish green	Yellowish green
T <sub>11</sub>	Yellow polythene bag	8.1	8.1	7.5	8.1	8.0	Light Yellow	Light green	Light Green	Greenish Yellow
T <sub>12</sub>	Green polythene bag	8.5	8.5	8.2	8.6	8.5	Light Green	Light Yellow	Light Green	Light Yellow
T <sub>13</sub>	Blue polythene bag	8.8	8.9	8.9	8.5	8.8	Light Green	Light Green	Light Green	Light Green
T <sub>14</sub>	Cotton cloth bag	8.5	8.9	7.9	8.9	8.5	Light Green	Light Yellow	Light Green	Light Yellow
T <sub>15</sub>	Muslin cloth bag	7.8	7.8	7.9	7.3	7.7	Greenish Yellow	Yellowish green	Greenish Yellow	Yellowish green
T <sub>16</sub>	Control (unbagged)	6.5	6.5	6.2	6.1	6.3	Greenish Yellow	Greenish Yellow	Greenish Yellow	Greenish Yellow
<b>Treatment Mean</b>		7.6	7.6	7.5	7.5					
<b>C.D. (p=0.05)</b>		<b>Treatments (T) = 0.35, Cultivars (C) = NS, T x C = 0.70</b>								