

## Original Research Article

# Prevalence, Trends, and Sociodemographic Determinants of Sexually Transmitted Diseases in Bangladesh

### ABSTRACT

**Background:** Sexually Transmitted Diseases (STDs) pose a significant public health challenge in Bangladesh, with implications spanning health, society, and economics. This research, conducted with a sample size of 2,794 participants, delves into the prevalence, trends, and sociodemographic determinants of STDs within the Khulna Division.

**Methods:** A stratified random sampling technique was employed, and data collection involved structured questionnaires, medical examinations, and laboratory tests. Descriptive and inferential statistical analyses, including chi-square tests and logistic regression, were used to interpret the data.

**Results:** Chlamydia and Gonorrhea exhibited concerning upward trends, while Syphilis and HIV remained relatively stable. Young adults (20-29 years), unmarried individuals, and those with secondary education displayed higher STD prevalence.

**Conclusion:** The findings underscore the urgency of targeted interventions to address rising STD rates, particularly among high-risk groups. Public health strategies should focus on awareness, safe sexual practices, and improved access to screening and treatment services. Addressing disparities in sexual health knowledge and services among different educational groups is essential.

*Keywords: Sexually Transmitted Diseases (STDs), Prevalence, Sociodemographic determinants, Bangladesh, Public health.*

### 1. INTRODUCTION

Sexually Transmitted Diseases (STDs) represent a significant global public health challenge, with their impact extending far beyond individual health to encompass social, economic, and demographic dimensions [1]. In Bangladesh, a densely populated South Asian country, the burden of STDs has garnered increasing attention due to its potential to disrupt the overall health and well-being of the population [2]. With a burgeoning population and diverse sociodemographic factors, Bangladesh faces unique challenges in addressing the prevalence and trends of STDs [3]. The high prevalence of STDs in Bangladesh is closely linked to a range of factors, including sexual behaviors, access to healthcare, and sociodemographic determinants [4]. Understanding these factors is crucial for designing effective prevention and intervention strategies tailored to the specific needs of the Bangladeshi population [5]. Moreover, as the global health landscape evolves, it is imperative to continually assess the prevalence and trends of STDs to adapt public health policies and programs accordingly [6].

This comprehensive analysis aims to provide a thorough examination of the current prevalence and trends of STDs in Bangladesh, taking into account the sociodemographic determinants that influence their spread [7]. By synthesizing existing data and employing advanced epidemiological methods, this study seeks to offer valuable insights into the

dynamics of STDs in the country [8]. This research is driven by the recognition that addressing STDs in Bangladesh necessitates a multidisciplinary approach, encompassing not only the medical aspects of disease control but also sociocultural, economic, and healthcare infrastructure considerations [9]. Therefore, the findings of this analysis will inform evidence-based policies and interventions that can contribute to the reduction of STD prevalence in Bangladesh and promote the overall health and well-being of its population [10].

### **Objective**

The objective of this research is to analyze the prevalence and trends of Sexually Transmitted Diseases (STDs) in Bangladesh, identify sociodemographic determinants of STD transmission, assess the impact of behavioral factors, evaluate healthcare accessibility, and provide evidence-based policy recommendations to enhance STD prevention and control efforts in the country.

## **2. METHODS**

### **2.1 Study Area, Sampling Technique, and Sample Size**

The research was conducted within the Khulna Division of Bangladesh, a region characterized by its diverse sociodemographic and geographic characteristics. Khulna Division was selected as the study area due to its significant population density and varied socioeconomic factors, making it representative of broader conditions in Bangladesh. A stratified random sampling technique was employed to ensure a representative sample of the population. Khulna Division was divided into distinct strata based on geographical zones, including urban and rural areas, to account for regional disparities. In each stratum, clusters of sub-districts were randomly selected.

The sample size for this study was determined using the following formula:

$$n = \frac{Z^2 pq}{E^2}$$

where:

- n is the sample size,
- Z is the z-value corresponding to the desired confidence level (1.96 for 95% confidence),
- p is the estimated prevalence of UTIs among the target population (derived from preliminary studies),
- q is 1-p,
- E is the margin of error.

### **2.2 Variables**

The variables considered in this study include:

- Dependent Variable: Prevalence of Sexually Transmitted Diseases (STDs) within the Khulna Division.
- Independent Variables: Sociodemographic factors such as age, gender, education level, marital status, and socioeconomic status, as well as behavioral factors including condom usage, number of sexual partners, and healthcare-seeking behavior.

### **2.3 Statistical Analysis:**

Data analysis was conducted using standard statistical methods. Descriptive statistics, such as means and percentages, were employed to describe the prevalence and distribution of STDs within the region. Inferential statistics, including chi-square tests and logistic

regression, were used to explore associations between independent variables and STD prevalence.

- Chi-square ( $\chi^2$ ) test was used to examine the association between categorical variables.
- Logistic regression was employed to model the relationship between the independent variables and the binary outcome variable (presence or absence of STDs).

The significance level for all statistical tests was set at  $\alpha=0.05$  to determine statistical significance.

### 3. RESULTS

Our comprehensive analysis was based on a substantial sample size of 2,794 participants, representative of various sociodemographic groups in Bangladesh. The results are presented under two main subsections: the prevalence and trends of sexually transmitted diseases (STDs), and the sociodemographic determinants influencing the occurrence of these diseases.

#### 3.1 Prevalence and Trends of STDs

Table 1 illustrates the prevalence and trends of STDs among the study participants. We observed a prevalence rate of 12.3% for Chlamydia, 8.7% for Gonorrhea, 3.2% for Syphilis, and 2.1% for HIV. The chi-square test results showed significant trends over time, with Chlamydia ( $\chi^2 = 15.23$ ,  $p < 0.05$ ) and Gonorrhea ( $\chi^2 = 10.78$ ,  $p < 0.05$ ) showing an upward trend, whereas Syphilis and HIV remained relatively stable.

**Table 1: Prevalence and Trends of STDs**

STD Type	Prevalence (%)	Chi-Square Test	P-Value
Chlamydia	12.3	15.23	<0.05
Gonorrhea	8.7	10.78	<0.05
Syphilis	3.2	2.11	>0.05
HIV	2.1	1.98	>0.05

#### 3.2 Sociodemographic Determinants of STDs

Table 2 reveals the associations between various sociodemographic factors and the prevalence of STDs. Age, marital status, and level of education were all found to have a significant association with STD prevalence. Individuals aged 20-29 years had the highest prevalence (14.2%,  $\chi^2 = 13.47$ ,  $p < 0.05$ ), followed by those who were unmarried (11.9%,  $\chi^2 = 12.03$ ,  $p < 0.05$ ) and those with a secondary level of education (10.4%,  $\chi^2 = 8.76$ ,  $p < 0.05$ ).

**Table 2: Sociodemographic Determinants of STDs**

Sociodemographic Factor	Prevalence (%)	Chi-Square Test	P-Value
Age (20-29 years)	14.2	13.47	<0.05
Marital Status (Unmarried)	11.9	12.03	<0.05
Education (Secondary Level)	10.4	8.76	<0.05
Rural/Urban Residence	9.1/7.3	3.54	>0.05
Income Level (Low)	9.8	4.67	>0.05

### 4. DISCUSSION

The findings regarding the prevalence and trends of STDs are noteworthy. Chlamydia and Gonorrhea displayed a significant upward trend over time, which is concerning. These trends could be attributed to a variety of factors, including changes in sexual behavior, increased access to healthcare services, and improved diagnostic methods [11]. Effective public health strategies are needed to curb the rising rates of these infections. Syphilis and HIV, on the other hand, showed relative stability in prevalence. While this is a positive sign, it underscores the importance of maintaining robust prevention and control programs to prevent any resurgence in these diseases. Additionally, continued monitoring and surveillance are essential to promptly identify any shifts in the epidemiological landscape [12].

The associations between sociodemographic factors and STD prevalence shed light on vulnerable populations. Young adults between 20 and 29 years of age exhibited the highest STD prevalence, a trend that aligns with global patterns [13]. This underscores the need for targeted sexual health education and interventions for this age group. Marital status emerged as another significant determinant, with unmarried individuals displaying a higher prevalence of STDs. This finding underscores the importance of promoting safe sexual practices, including consistent condom use, among individuals regardless of their marital status. Education level also played a role, with those with a secondary level of education showing higher STD prevalence. This could be indicative of disparities in access to sexual health information and services among different educational groups [14]. Addressing these disparities through educational outreach and accessible healthcare services is crucial.

The implications of our findings are clear. To combat the rising trends of Chlamydia and Gonorrhea, public health initiatives should focus on increasing awareness, promoting safe sexual behaviors, and enhancing access to screening and treatment services. Furthermore, targeted interventions for young adults, unmarried individuals, and those with lower educational attainment are essential to reduce the burden of STDs in these vulnerable populations. Our research contributes to the understanding of STD epidemiology in Bangladesh, providing a foundation for evidence-based public health policies and interventions. Continued surveillance and research are vital to adapt strategies to the evolving landscape of STDs and safeguard the sexual health of the population.

#### **4. CONCLUSION**

Our examination of STD prevalence trends revealed a concerning increase in Chlamydia and Gonorrhea cases. These upward trends emphasize the urgency of targeted interventions to address these infections. Understanding the drivers of this increase, including changes in sexual behaviors and healthcare-seeking patterns, is essential for tailoring effective prevention and control strategies. Syphilis and HIV exhibited relative stability in prevalence. However, complacency is not an option. Sustaining efforts in prevention, diagnosis, and treatment remains crucial to prevent any resurgence in these diseases.

The associations we identified between sociodemographic factors and STD prevalence underscore the importance of targeted approaches. Young adults, particularly those aged 20 to 29, emerged as a high-risk group. Implementing sexual health education programs and promoting safer sexual behaviors among this demographic is imperative. Marital status was also a significant determinant, highlighting the need to address sexual health within both married and unmarried populations. Education level played a role, with those having secondary education showing higher prevalence. This points to disparities in access to sexual health information and services that must be addressed through educational outreach and improved healthcare accessibility.

Our research carries clear implications for public health policies and interventions in Bangladesh. To combat rising trends in Chlamydia and Gonorrhoea, efforts should focus on raising awareness, promoting safe sexual practices, and enhancing access to screening and treatment services. Targeted interventions for high-risk groups, including young adults and unmarried individuals, are essential components of a comprehensive strategy. Addressing disparities in sexual health knowledge and access to services among different educational groups should be a priority. Public health initiatives must work to ensure that information and resources are accessible to all segments of the population. As the landscape of STDs evolves, it is imperative to remain vigilant through continued surveillance, research, and the adaptation of strategies. By doing so, we can safeguard the sexual health and well-being of the population and work toward a future with reduced STD prevalence and associated health burdens.

### **ETHICAL APPROVAL**

The ethical approval for this study was considered by the Ministry of Health, Government of Peoples Republic of Bangladesh

### **REFERENCES**

1. Newman L, Rowley J, Vander Hoorn S, et al. Global Estimates of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting. *PLoS ONE*. 2015;10(12):e0143304.
2. Ahmed SM, Ahmed S. Reducing the Burden of Sexually Transmitted Infections in Resource-Poor Countries: The Role of Improved Diagnostics. *Nature Reviews Microbiology*. 2006;4(12):S23-S33.
3. Bangladesh Bureau of Statistics. Population and Housing Census 2011: Preliminary Results. Dhaka: Bangladesh Bureau of Statistics; 2012.
4. Hawkes S, Morison L, Foster S, et al. Reproductive-tract Infections in Women in Low-income, Low-prevalence Situations: Assessment of Syndromic Management in Matlab, Bangladesh. *The Lancet*. 1999;354(9192):1776-1781.
5. Blanchard JF, O'Neil J, Ramesh BM, Bhattacharjee P, Orchard T, Moses S. Understanding the Social and Cultural Contexts of Female Sex Workers in Karnataka, India: Implications for Prevention of HIV Infection. *Journal of Infectious Diseases*. 2005;191(Suppl 1):S139-S146.
6. World Health Organization. Global Health Sector Strategy on Sexually Transmitted Infections 2016-2021: Towards Ending STIs. Geneva: World Health Organization; 2016.
7. Alam N, Streatfield PK, Khan SI, Momtaz D, Kristensen S, Vermund SH. Factors Associated with Condom Use Among Persons with Multiple Sexual Partners in Two Rural Areas of Bangladesh. *International Journal of STD & AIDS*. 2007;18(1):48-53.
8. Sarker MA, Sanou AK, Snow R, Ganame J, Gondos A. Determinants of HIV Counseling and Testing Participation in a Prevention of Mother-to-Child Transmission Program in Rural Burkina Faso. *African Journal of AIDS Research*. 2007;6(3):271-279.
9. Ahmed SM, Adams AM, Chowdhury M, Bhuiya A. Gender, Socioeconomic Development and Health-seeking Behaviour in Bangladesh. *Social Science & Medicine*. 2000;51(3):361-371.
10. Government of the People's Republic of Bangladesh. National Health Policy 2011. Dhaka: Ministry of Health and Family Welfare; 2011.

11. Mercer, C. H., Aicken, C. R., Tanton, C., Estcourt, C. S., Brook, M. G., Keane, F., ... & Johnson, A. M. (2016). Serial monogamy and biologic concurrency: measurement of the gaps between sexual partners to inform targeted strategies. *AIDS*, 30(8), 1333-1342.
12. Cohen, M. S., Smith, M. K., Muessig, K. E., Hallett, T. B., Powers, K. A., & Kashuba, A. D. (2013). Antiretroviral treatment of HIV-1 prevents transmission of HIV-1: where do we go from here?. *The Lancet*, 382(9903), 1515-1524.
13. World Health Organization. (2016). Report on global sexually transmitted infection surveillance 2018. World Health Organization.
14. Maticka-Tyndale, E., & Tenkorang, E. Y. (2010). A multi-level model of condom use among male and female upper primary school students in Nyanza, Kenya. *Social Science & Medicine*, 71(3), 616-625.

UNDER PEER REVIEW