

Original Research Article

Studies on Integrated Nutrient Management on Yield And Quality Characters of Elephant Foot Yam (*Amorphophallus paeonifolius* (Dennst.) Nicolsan) cv. Gajendra

ABSTRACT

Aims:The investigation was carried out to study the intergrated nutrient management practices on yield and quality characters of Elephant foot yam (*Amorphophallus paeonifolius* (Dennst.) Nicolsan) cv. Gajendra.

Study design:The field experiment was laid out in randomized block design with thirteen treatments with three replications.

Place and Duration of Study:The present study was conducted at farmer's field at Thirukkanur village, Villianur, Puducherry during the year 2020 with spacing 60 x 60 cm.

Methodology:Elephant foot yams corms were ~~cutted cut~~ into ~~pieces of~~ 800 g for planting, after cutting corms were dipped with cow dung slurry and placed for drying. The treatments comprised of three levels of recommended dose of fertilizers (RDF - 80:40:100 Kg NPK ha⁻¹) viz., 75%, 100% and 150% along with organic inputs viz., FYM (20 t ha⁻¹), Vermicompost (5 t ha⁻¹) and (CBF 5 Kg ha⁻¹), organic manures of different combinations were applied as basal; where as recommended dose of fertilizer were also applied combined with organic manures. The observations were recorded at 190 days after planting.

Results:The results indicated that the maximum values for yield parameters viz., Culm girth (14.60 cm), Corm circumference (46.86 cm), Single Corm weight (48.21 Kg) plant⁻¹, Yield plant⁻¹ (2.42 Kg), Yield per plot (60.76 Kg), Total yield (40.15t ha⁻¹) whereas quality parameters such as Moisture content (79.64 %), Starch content (18.02 %), Crude protein(6.11 %), Calcium oxalate (12.18 mg g⁻¹), Total sugar content (2.62 mg g⁻¹) and β carotene (0.19 μ g g⁻¹).

Conclusion:From the experiment, it was concluded that the integrated nutrient management viz., combination of vermicompost with 100 per cent RDF along with consortium biofertilizer were identified as the best treatment to increase quality and yield.

Keywords: [NPK; FYM; Vermicompost; CBF]

1. INTRODUCTION

Amorphophallus paeoniifolius is a tropical plant cultivated throught out Southeast Asia, South Asia. Due to potential and production capacity, it can be grown as cash crop. It is widely used as traditional vegetable by rural population and used as secondary food during the period of scarcity of other vegetable. Corms are typically eaten by boiling, mashed or pounded and combined with other vegetables. It is widely used in Ayurveda, Siddha and Unani and the tubers are also useful in treatment of piles. It has become crucial to diversify the current agriculture in order to fulfil the ~~suitt~~ different human requirements due to the strain of an ever-increasing population and the rapid depletion of natural resources. Inorganic fertilizers ~~were~~ ~~are~~ commonly believed to be an effective means to raise production in poor ~~fertile~~ soils and thereby leading to stagnation. As an alternative way, to overcome the dependence on inorganic fertilizers, usage of organic waste integrated with chemicals and bio-fertilizers was recommended. Hence the field experiment ~~was has been~~ taken up to evaluate the effect of different organic and inorganic fertilizers ~~to increase on~~ yield and quality.

Comment [hp1]: Follow the format of Jouranl

Comment [hp2]: Keep the font of the manuscript 12

Comment [hp3]: Review of other uthors is mandatory

Comment [hp4]: Add citations

2. MATERIAL AND METHODS

The present study entitled “Studies of integrated nutrient management on yield and quality of Elephant foot yam (*Amorphophalus paeonifolius* (Dennst.) Nicolsan) cv. Gajendra” was conducted at farmers field at Thirukkanur village, Villianur, Puducherry during the year 2020, with 13 treatments, 60x60 cm spacing and 3 replications. Elephant foot yams corms were cutted into 800 g for planting, after cutting corms were dipped with cow dung slurry and placed for drying.

2.1 Treatment Details

The treatments comprised of three levels of recommended dose of fertilizers (RDF - 80:40:100 Kg NPK ha⁻¹) viz., 75%, 100% and 150% along with organic inputs viz., FYM (20 t ha⁻¹), Vermicompost (5 t ha⁻¹) and (CBF 5 Kg ha⁻¹) (Table 1). The treatment details were presented in (Table 1). Organic manures of different combinations were applied as basal, where as recommended dose of fertilizer were also applied combined with organic manures. The observations were recorded at 190 days after planting. Observations were recorded at different stages of the crop. Fifteen plants were tagged randomly in each net plot of each treatment for recording the following observations. The data result of the field experimental yield components has been presented in (Table 2) and (Fig. 1).

Comment [hp5]: Give full form of CBF

TABLE 1. List of treatments used for the study

Treatment	Treatment Details
T ₁	FYM 20 t ha ⁻¹ + 75% RDF (60:30:75 Kg NPK ha ⁻¹)
T ₂	FYM 20 t ha ⁻¹ + 100% RDF (80:40:100 Kg NPK ha ⁻¹)
T ₃	FYM 20 t ha ⁻¹ + 125% RDF (100:50:125 Kg NPK ha ⁻¹)
T ₄	FYM 20 t ha ⁻¹ + 75% RDF (60:30:75 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₅	FYM 20 t ha ⁻¹ + 100% RDF (80:40:100 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₆	FYM 20 t ha ⁻¹ + 125% RDF (100:50:125 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₇	Vermicompost 5 t ha ⁻¹ + 75 % RDF (60:30:75 Kg NPK ha ⁻¹)
T ₈	Vermicompost 5 t ha ⁻¹ + 100 % RDF (80:40:100 Kg NPK ha ⁻¹)
T ₉	Vermicompost 5 t ha ⁻¹ + 125 % RDF (100:50:125 Kg NPK ha ⁻¹)
T ₁₀	Vermicompost 5 t ha ⁻¹ + 75 % RDF (60:30:75 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₁₁	Vermicompost 5 t ha ⁻¹ + 100 % RDF (80:40:100 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₁₂	Vermicompost 5 t ha ⁻¹ + 125 % RDF (100:50:125 Kg NPK ha ⁻¹) + CBF 5 Kg ha ⁻¹
T ₁₃	Control

3. RESULTS AND DISCUSSION

The experiment was laid out during kharif, 2020 at the farmer field at Thirukkanur village, Villianur, Puducherry to assess the Study effect of integrated nutrient management on yield and quality characters of Elephant foot yam (*Amorphophalus paeonifolius* (Dennst.) Nicolsan) cv. Gajendra.

The data result of the field experimental yield components has been presented in (Table 2) and (Fig. 1).

3.1 YIELD PARAMETERS AFFECTED BY INTEGRATED NUTRIENT MANAGEMENT PRACTICES

Formatted: Justified

3.1.1 CULM GIRTH, CORM CIRCUMFERENCE, SINGLE CORM WEIGHT, CORM WEIGHT PLANT⁻¹, YIELD PLOT⁻¹ AND TOTAL YIELD.

Among the various ~~treatements~~treatments, T₁₁ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹) + CBF 5 Kg ha⁻¹] recorded the maximum values for yield parameters viz., Culm girth (14.60 cm), Corm circumference (46.86 cm), Fresh corm weight (2009.17 g), Corm weight (48.21 Kg) plant⁻¹, Yield plant⁻¹ (2.42 Kg), Yield per plot (60.76 Kg) and Total yield (40.15 t ha⁻¹) followed by T₈ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹)] and the least values were recorded in T₁₃ (control).

Comment [hp6]: Duplicated para

The yield attributes viz., Culm girth (cm), Corm circumference (cm), Single Corm weight plant⁻¹ (kg), Yield plant⁻¹ (kg), Yield plot⁻¹ (kg), Total yield (t ha⁻¹) were significantly affected by the application of various inorganic and organic inputs. From the results of the present study that maximum values for yield parameters were recorded in T₁₁ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹) + CBF 5 Kg ha⁻¹], followed by T₈ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹)] and the least values were recorded in T₁₃ (control).

Corm yield is determined with the amount of dry materials translocated to the corms and the ability of the corm, to store assimilates as evidenced by the volume and diameter of the corms is regarded as a crucial determinant influencing eventual yield. Yield of elephant foot yam depends on photosynthetic capacity and duration of ~~that capacity of~~ the crop. ~~Part of the photosynthates produced by the leaves would be translocated for corm development.~~

The data on yield attributes revealed ~~that~~, organic manures along with inorganic fertilizers at different levels exert a significant influence on corm yield and yield attributes. From the results it could be concluded that the 20% nutrient requirement could be substituted through organic source as vermicompost, FYM without any yield loss. As the application of vermicompost might have enhanced soil micro flora activity, besides supplementing nutrients. The combined application along with biofertilizers increased the availability of soil nitrogen and phosphorus. The corm yield ~~was~~ increased gradually with increase in fertility levels. Similar observations ~~were has been~~ recorded ~~by~~[6]. The current study's findings are consistent with those published by [16], [3], [13], [11], and [12]. Furthermore, [1] found a substantial positive relationship between plant height, dry matter output, and yield qualities and cassava tuber yield, which is consistent with the current study's findings [1].

Comment [hp7]: Reference?


Comment [hp8]: Give here the scientist name

Comment [hp9]: In order to write the reference like this you need to give the results of the studies

TABLE. 2. EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON YIELD PARAMETERS OF ELEPHANT FOOT YAM cv. Gajendra.

Treatments	Culm girth (cm)	Corm circumference (cm)	Single Corm weight plant ⁻¹ (Kg)	Yield plant ⁻¹ (Kg)	Yield plot ⁻¹ (Kg)	Total yield (t ha ⁻¹)
T ₁	11.13	40.38	29.07	1.36	39.02	24.21
T ₂	13.49	44.72	43.84	2.12	54.81	36.51
T ₃	12.36	42.59	36.26	1.75	46.90	30.20
T ₄	11.54	41.06	30.72	1.45	41.00	25.58
T ₅	13.85	45.42	45.84	2.22	56.78	38.18
T ₆	12.74	43.29	38.40	1.85	48.86	31.98
T ₇	11.96	41.78	32.86	1.56	42.96	27.37

T ₈	14.23	46.14	47.23	2.32	58.76	39.75
T ₉	13.11	44.00	40.08	1.95	50.84	33.38
T ₁₀	12.00	41.89	34.56	1.65	44.93	28.78
T ₁₁	14.60	46.86	48.21	2.42	60.76	40.15
T ₁₂	13.46	44.68	41.83	2.03	52.83	34.84
T ₁₃	9.47	37.52	25.63	1.27	37.06	21.34
S.Ed	0.17	0.34	1.17	0.03	0.98	-
CD (p=0.05)	0.34	0.68	2.35	0.07	1.95	-



Comment [hp10]: Add error bars

Comment [hp11]: Take yield on secondary axis

Fig. 1. Effect of integrated nutrient management on yield parameters of elephant foot yam cv. gajendra.

3.2 QUALITY PARAMETERS AFFECTED BY INTEGRATED NUTRIENT MANAGEMENT PRACTICES

3.2.1 MOISTURE, STARCH, TOTAL SUGAR CONTENT, CRUDE PROTEIN, CALCIUM OXALATE AND BETA CAROTENE CONTENT

The quality parameters such as Moisture content (79.64 %), Starch content (18.02 %), Crude protein (6.11 %), Calcium oxalate (12.18 mg g⁻¹), Total sugar content (2.62 mg g⁻¹) and β carotene (0.19 μg 100 g⁻¹) recorded the maximum values in T₁₁ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹) + CBF 5 Kg ha⁻¹] which is followed by T₈ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹)] and the least values were recorded in T₁₃ (control).

The data on quality attributes such as Moisture content (%), Starch content (%), Crude protein (%), Calcium oxalate (mg g⁻¹), Total sugar content (mg g⁻¹) and β carotene (μg g⁻¹) were significantly affected by the application of various inorganic and organic inputs. From the results of the present study shows that maximum values for quality parameters were recorded in T₁₁ [Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹) + CBF 5

Kg ha⁻¹], followed by T₈[Vermicompost 5 t ha⁻¹ + 100 % RDF (80:40:100 Kg NPK ha⁻¹)] and the least values were recorded in T₁₃ (control). Similar findings have been reported by [14] and [4] in onion, [7] in *Amorphophallus*.

The increased moisture content in corm might be as a result of maintenance of soil physical structure and thus resulted in better moisture retention in the underground storage organ which may have resulted in increased aeration, porosity and water holding capacity of soils.

The increase appears to be related to greater potash levels, which aided in creation and transfer of starch from the leaves to tubers [2] and might be due to its capacity in increasing parenchymatous cells which contains starch grains [6]. Whereas, the trend with respect to the protein content indicated that N application either in the form of inorganic fertilizers or organic manure plays a key role in enhancing the protein content of corms because nitrogen is a major constituent of protein.

Increased N resulted in higher amounts of crude protein in cassava tubers were reported by [10] which is similar with the findings of the above results. The protein content has significantly reduced in treatment receiving low levels of nitrogen, particularly in control. These results were similar with [6] where maximum starch and protein contents of the corms were observed at higher N and K doses in *Amorphophallus*. The integrated use of organic manures (vermicompost, FYM) favourably affected the crude protein content. The highest crude protein content might be due to the result of increased nitrogen supply in balanced proportion.

Higher nitrogen content through inorganic fertilizers resulted in the accumulation of more oxalates and the biochemical studies conducted in the present investigation clearly revealed the above fact. It can also be adjudged that this trait is more genotype specific and is least affected by growing environment [15]. The results are in similar with [15].

The increased level of beta carotene due to combined use of organic manures might have resulted in enhancement in quality parameters of elephant foot yam. Nitrogen is most indispensable of all mineral nutrients for growth and development of plant. [5; 8] and [8] also supported the findings regarding beta carotene and calcium oxalate.

Comment [hp12]: Citation used in paragraph should be narrated i.e. Reported by Shinde (Year) and Jayathilake (year)

Comment [hp13]: Add reference

TABLE. 3. EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON QUALITY PARAMETERS OF ELEPHANT FOOT YAM cv. Gajendra.

Treatments	Moisture content (%)	Starch content (%)	Crude protein (%)	Calcium oxalate (mg g ⁻¹)	Total sugar content (mg g ⁻¹)	β carotene (μg g ⁻¹)
T ₁	68.36	13.82	5.42	10.10	1.80	0.158
T ₂	76.34	17.46	5.93	11.68	2.44	0.188
T ₃	72.89	15.83	5.72	10.82	2.16	0.175
T ₄	69.50	14.35	5.55	10.31	1.90	0.162
T ₅	77.45	17.98	6.00	11.73	2.45	0.189
T ₆	74.00	16.36	5.78	11.05	2.26	0.179
T ₇	70.62	14.88	5.62	10.54	1.99	0.166
T ₈	78.55	17.50	6.05	11.96	2.53	0.192
T ₉	75.17	16.89	5.85	11.08	2.27	0.180
T ₁₀	71.73	15.31	5.64	10.60	2.00	0.170
T ₁₁	79.64	18.02	6.11	12.18	2.62	0.196
T ₁₂	76.28	17.42	5.92	11.43	2.35	0.184
T ₁₃	64.59	11.36	5.22	9.32	1.38	0.131
S.Ed	0.54	0.24	0.03	0.10	0.04	0.002
CD (p=0.05)	1.08	0.48	0.05	0.19	0.07	0.003

Formatted: Superscript

4. CONCLUSION

Based on the findings of the present investigation, it can be concluded that the application of 5 t ha^{-1} of vermicompost combined with 100% RDF ($80:40:100 \text{ Kg NPK ha}^{-1}$) + 5 kg ha^{-1} of CBF (T_{11}) can be considered as best inorganic and organic combination to obtain maximum yield and quality characters whereas T_{13} (control) obtain minimum yield and quality of corms.

REFERENCES

1. Amanullahkhan MD. Influence of fertilizer levels, intercropping and organic manures on the growth and yield of cassava. 1997. Ph.D Thesis, Tamil Nadu Agricultural University, Coimbatore, India.
2. Ashokan PK and Sreedharan C. Influences of levels and time of application of potash on growth, yield and quality of tapioca. *Journal of Root crops*, 1977;3(2):1-4.
3. Jawahar D and Suresh S. Effect of bio-fertilizers and enriched FYM on the yield of cotton under dry farming in vertisol. *Ad. Plant.Sci.* 2007;20(11): 473-475.
4. Jayathilake. Integrated nutrient management of rabi onion (*Allium cepa* L.). M.Sc (Hort.) thesis, Acharaya N.G. Ranga Agricultural University, Hyderabad.
5. Kolambe, BN, Patel KG, Kaswala AR, Patel VS, Desai KD. Organic management affects growth, yield and soil properties in elephant foot yam. *J Root Crops*. 2010;39(1): 62-66.
6. Mukhopadhyay SK and Sen H. Effect of nitrogen and potassium on yield and quality of elephant foot yam (*Amorphophallus campanulatus* Blume.). *Journal of Root crops*, 1986;12(2): 103-106.
7. Murthy GN, Padma E, Bhagavan BVK, Naga Lakshmi R. Effect of integrated nutrient management on growth and yield of (*Amorphophallus paeonifolius* (Dennst.) Nicolsan) cv. Gajendra. pp 2011;294-297.
8. Navya K, Desai KD, Tandel YN and Sheth SG. Effect of integrated nutrient management on growth, yield and quality of elephant foot yam. *Int J Chemical Studies*. 2017;5(4): 1766-1769.
9. Pushpalatha M, Vaidya PH and Badsul P. Effect of graded levels of nitrogen and potassium on yield and quality of sweet potato (*Ipomea batatas* L.). *Int J Current Microbiology and Applied Sci.* 2017;6: 1689-1696.
10. Pusphadas MV and Aiyer RS. Nutritional studies on cassava (*Manihot esculenta* Crantz.). *Journal of root crops*. 1976;2(2):42-71.
11. Saraswathi T, Nageswari K, Pugalendhi L and Sathiyamoorthy VA. Integrated nutrient management in elephant foot yam. National Seminar On *Amorphophallus*: Innovative Technologies; Abstract book, status papers and extended summary, July 19-20, 2008, Patna, Bihar.
12. Saravaiya SN, Chaudhary PP, Patel DA, Patel B, Ahir MP and Patel VI. Influence of integrated nutrient management on growth and yield parameters of elephant foot yam under south Gujarat condition. *Asian Journal of Horticulture*. 2010; 5 (1): 58-60.
13. Sengupta S, Nath S, Singh BN and Ray RN. Effect of integrated nutrient management on growth and productivity of *Amorphophallus* cv. Gajendra at Ranchi condition in Jharkhand. National Seminar on *Amorphophallus*: Innovative Technologies; Abstract book, status papers and extended summary, July 19-20, 2008, Patna, Bihar
14. Shinde PH, Naik RL, Nazikar RB, Kadam SK and Khaire VM. Evaluation of vermicompost. Proceedings of national seminar on organic farming. MPKV, Pune.

Comment [hp14]: Arrange the references in ascending order. Follow the format of Journal

15. Suja G and Sundaresan S. Organic elephant foot yam production: A feasible strategy for high yield and income. National seminar on *Amorphophallus*: Innovative technologies; Abstract book, status papers and extended summary, July 19-20, 2008, Patna, Bihar.
16. Yadav BD, Khandelwal RB and Sharma YK. Use of biofertilizer in onion. Indian Journal of Horticulture, 2005; 62(2): 168-170

ABBREVIATIONS

%	:	per cent
()	:	Bracket parenthesis
@	:	At the rate of
[]	:	Square bracket
=	:	is equal to
CBF	:	Consortium Biofertilizers
C.D.	:	Critical difference
cm	:	Centimeter
cv.	:	Cultivar
Fig.	:	Figure
FYM	:	Farmyard manure
g	:	gram
ha ⁻¹	:	per hectare
S.E.M. ±	:	Standard error of means
mg g ⁻¹	:	Milligram per gram
NPK	:	Nitrogen, Phosphorus and Potassium
Kg	:	Kilogram
No.	:	Number
Plant ⁻¹	:	per plant
Plot ⁻¹	:	per plot
RDF	:	Recommended dose of fertilizers
ha ⁻¹	:	per hectare
t	:	tonne
viz.	:	Namely

UNDER PEER REVIEW

