

Nutritional Quality of Complementary food blended with Moringa Leaf Powder.

ABSTRACT

Nutritional Quality of Complementary porridge made from Pearl millet, Orange-Fleshed Sweet Potato, Dates and blended with Moringa Leaf Powder was evaluated using standard methods. The objective of this study was to formulate and assess highly nutritious food for young children. A total of 7 formulations were made, whose compositional proportions were calculated based on the Recommended Daily Allowances (RDA) for children aged 1-2 years. The Pearl millet grain flour, Moringa leaf powder and dates were blended in ratios of 90:2.5:7.5 [PmMD1], 95:2.5:2.5 [PmMD2] and 88:3:9 [PmMD3], forming three formulations, respectively. Also, Pearl millet grain flour, Moringa leaf powder and Orange fleshed sweet potato were blended in ratios of 95.5:2:2.5 [PmMP1], 88:2:10 [PmMP2], and 95:2.5:2.5 [PmMP3] forming three formulations, respectively. A control sample did not contain Moringa powder and was composed of Pearl millet, Dates and Orange fleshed sweet potato in a ratio of 50:15:35 [PmDP], respectively. The nutritional composition of the flour on dry weight basis was; crude fiber: 2.6-3.3 g/100g, ash: 1.9-2.13 g/100g, Crude fat: 0.66-1.38 g/100g, crude protein: 8.34-11.07 g/100g, Carbohydrates: 73-77 g/100g and Energy of 346-352g/100g. The content of iron, zinc and calcium was 4.5-6.3, 1.46-1.81, 183-240mg/100g, respectively and Vitamin A content was 3082-4975 µg/110g. The incorporation of Moringa leaf powder improved the nutritional quality of the complementary food.

Key words: Complementary food, Micronutrients, Moringa leaf, Recommended Daily Allowance.

INTRODUCTION

Childhood under nutrition remains a pressing issue worldwide, particularly affecting children under the age of 5, with a significant number of affected children residing in developing countries (UNICEF, 2009). The problem of malnutrition often arises in infants during or after the introduction of complementary foods, contributing significantly to the high prevalence of malnutrition in children below the age of 5 (Muhimbula and Zacharia, 2010; Mosha *et al.*, 2000). According to these studies, a majority of infants are introduced to cereal-based complementary foods well before the recommended age of 6 months or, in some cases; they do not receive these foods until they reach their second year of age.

One of the main causes of death among children and pregnant mothers or nursing mothers is malnutrition, which has permanent impacts on health in underdeveloped nations. Iron, zinc, iodine and vitamin A deficiency-related protein-energy malnutrition disorders have also been a problem to young children (Haidar *et al.*, 2003).

According to Mosha *et al.* (2000), Tanzanian traditional supplementary foods are based on starchy staples, typically cereals such as maize, rice, sorghum, and finger millet as well as non-cereals such as cassava, sweet potatoes, yams, bananas, and plantains. Unfortunately however, such diets are typically provided without enough supplementation with high quality protein sources (Kikafunda *et al.* 2006). These foods are widely available, inexpensive, and therefore accessible to most rural residents who are lysine deficient (FAO *et al.*, 2015). The primary cause of the widespread protein energy malnutrition that affects babies and young children is an overdependence on these starchy sources of protein.

Children need nutrient-rich diet that is high in vitamins, minerals, proteins, and carbohydrates for a healthy growth. These are essential for newborns and young children's healthy development and for the treatment of disorders brought on by a poor diet, such as malnutrition, which has long been recognized as a significant nutritional issue in developing nations (FAO *et al.*, 2015). After six months, it is appropriate to introduce complementary foods, which provide additional nutrition to meet all the growing child's needs, when breast milk alone is no longer sufficient to provide enough nutrients. In recent years inadequate child feeding procedures, a shortage of complementary foods of sufficient quantity and quality, and high rates of infections, have contributed to health and growth problems among children (FAO *et al.*, 2015).

Food-based strategies greatly improve the nutritional quality of foods while posing little risk in many developing countries (Omar and Michael, 2008; WHO/FAO, 2006), making them effective, acceptable, and sustainable approaches to combating the devastating effects of malnutrition. One potential approach is the creation of affordable, nutrient-dense complementary/supplementary foods employing underutilized local cereal, legume, root, and tuber crops (Forsido *et al.*, 2013; Omar and Michael, 2008; WHO/FAO, 2006).

Complementary food refers to any solid foods or nutritious beverages, aside from breast milk, that are introduced to infants to supplement their diet when mother's milk alone is no longer

sufficient. These foods are introduced to infants once they reach the age of six months, following a period of exclusive breastfeeding. Inadequate amounts and quality of complementary foods can result in elevated infection rates and other health complications in young children (FAO *et al.*, 2015). Research has indicated that more than 85% of the complementary foods given to infants fail to meet the nutrient density levels recommended by the World Health Organization (WHO), thereby contributing to malnutrition problems (Tenagashaw *et al.*, 2017).

The aim of current research was to produce and investigate the potential nutrient-rich and appetizing complementary food products from readily available, wholesome, reasonably priced, and locally grown underutilized crops such as the *Moringa oleifera* mixed with cereals and other non-cereal starchy foods.

Materials and Methods

Samples

Pearl millet and Orange fleshed sweet potato were obtained from Mawenzi market, in Morogoro, Moringa leaves were obtained from Frida homestead, Morogoro. Dry dates were obtained from the Kilombero market, Arusha Tanzania.

Sample preparation

The processing steps for the preparation of Pearl millet, Orange fleshed sweet potato, Dates and Moringa leaf flour are shown in Fig. 1, and their respective flour sample are shown in Fig.2.

Preparation of Moringa leaf powder

The fresh Moringa leaves from the farm were sorted and the young and fresh leaves were selected. Damaged and diseased leaves were discarded. The leaves were washed with clean water and soaked in 1% NaCl for 5 minutes solution to kill microbes. The excess water was drained and leaves were spread out on racks for 20 minutes before being shade dried at room temperature for 4 days. A high-speed multifunctional crusher model 750A was used to ground the leaves to powder. A 500 μ m sieve was used to obtain a fine powder (Olaitan *et al.*, 2014; Malla *et al.*, 2021).

Preparation of Orange-Fleshed Sweet Potato Flour.

Sweet potatoes were washed, peeled, and chopped into (3mm thick) slices. The slices were oven dried at 50°C overnight and later milled using a high-speed multifunctional crusher model 750A. A 500 μ m sieve was used to obtain a fine powder (Falade *et al.*, 2010).

Germination and Preparation of pearl millet

Pearl millet was sorted to remove extraneous matter and then washed to remove dust and mud. The sorted and cleaned millet was soaked in a 5 L bucket containing cold water for 10 hours at

room temperature. Water was drained from the millet grains and spread individually on wet muslin cloth where water was sprinkled at 6h break to stimulate the germination process. The millet grains germinated for 48 hours). The germinated millet was oven dried at 60°C for 10 h and then ground using a hammer mill into flour. A 500µm was used to sieve germinated flour (Nefale *et al.*, 2018). All the flour samples were kept in air-tight polyethylene bags and kept in a freezer at -10°C.

Fig 3: Pearl millet flour

Fig 4: Date powder




Fig. 5. Flowchart for the preparation of complementary food powder

Preparation of Date Powder

Dried dates from the market were washed with clean water to remove dust. Inner seeds were also removed and the dates were kept in the oven for 12 hours at 60°C. They were then left to dry for one hour. A high-speed multifunctional crusher model 750A was used to crush the dates so as to obtain a powder that was then sieved using a 500 µm (Hasan *et al.*, 2019).



Figure 2. Flour and Powder for the complementary food ingredients.

Formulation of potential complementary food

The linear regression method was used to combine the samples into different formulations to meet the FAO/WHO/UNICEF (1985) requirement for micronutrients for young children and infants. Seven samples (including the control sample) were formulated as indicated in Table 1.

Table 1: Composition of complemented food from pearl millet, orange-fleshed sweet potato, date, and Moringa leaf (g/100 g)

Ingredients	Formulation Name	Ratios
Pearl Millet+ Moringa leaf powder+ Dates	PmMD1	90:2.5:7.5
	PmMD2	95:2.5:2.5
	PmMD3	88:03:09
Pearl Millet+ Moringa leaf powder+ Orange-fleshed sweet potato	PmMP1	95.5:2:2.5
	PmMP2	88:02:10
	PmMP3	95:2.5:2.5
Pearl Millet+ Dates + Orange- fleshed + sweet potato (Control sample)	PmDP	50:15:35

Key: Pm = Pearl Millet, M= Moringa Leaf, D= Dates, P= Orange fleshed sweet potatoes

Proximate and mineral composition

The proximate composition of the product was determined according to official AOAC methods as shown below. Results were presented as means of duplicate determinations.

Crude fat

The total fat content was analyzed by using the official method 945.87 (AOAC, 1999) for Soxhlet ether extraction. A dry sample weighing 5 g was placed into the extraction thimble and assembled in the Soxhlet apparatus. Reflux was conducted using 60 mL of petroleum ether in three distinct phases: a boiling phase of 15 minutes, a fat extraction phase of 30 minutes, and a petroleum ether recovery phase of 10 minutes. The recovered petroleum ether was evaporated, and the resulting fat was collected in pre-weighed cups. To remove any remaining petroleum ether, the cups were dried in an oven at 105°C for 30 minutes. After cooling in a desiccator for 20 minutes, the cups were weighed.

Percentage fat was calculated by using the formula:

$$\% \text{ Crude fat} = \frac{\text{weight of crude fat (g)}}{\text{weight of dry sample (g)}} \times 100$$

Ash content

The determination of ash content was conducted following the procedure outlined in AOAC (1999), method 923.03. Initially, a dry sample weighing five grams was subjected to oven drying at 105°C for duration of 24 hours. The weight of the crucible and the dried sample was accurately recorded. Subsequently, the dried samples contained within the crucibles were incinerated in a muffle furnace at a temperature of 550°C for a period of three hours, resulting in the formation of grey ash. The ash content was calculated as the variance between the weight of the sample before and after the incineration process.

Percentage ash was calculated from the relationship:

$$\text{Ash (\%DM)} = \frac{\text{weight of ash (g)}}{\text{weight of dry sample (g)}} \times 100$$

Crude protein

The determination of crude protein content in the samples was performed using the micro-Kjeldahl method 920.87 as specified in AOAC (1999). Initially, dried samples weighing 0.5 g were accurately weighed and transferred into digestion tubes. To each tube, 0.6 g of a catalyst mixture comprising 10 g of K₂SO₄ and 0.5 g of CuSO₄, along with 6 mL of concentrated H₂SO₄, were added. The samples were then digested using a Tecator digestion system 40 (Model 1016 digester, Sweden) for a duration of 3 hours, resulting in a clear greenish solution. After cooling, the digested solution was transferred to a distillation unit (Foss Tecator, Model 2200 Kjeltec auto distilling unit, Sweden). A total of 70 mL of distilled water was added to the digested solution, followed by 70 mL of 40% NaOH, and the mixture was steam distilled for 4 minutes. The distillate, measuring 50 mL, was collected in a conical Erlenmeyer flask containing 25 mL of 4% boric acid. The collected distillate was then titrated with 0.105 g/100 mL hydrochloric acid. A blank volume was also determined, and a value of 0.04 mL was obtained.

$$\% \text{ Crude Protein} = \frac{14.01 \times (\text{Title} - \text{Blank}) \times 6.25 \times \text{Concentration of acid (n/mol)}}{\text{weight of dry sample (g)} \times 10} \times 100$$

Where 6.25 is the protein-nitrogen conversion factor

Crude fiber

The determination of crude fiber content was conducted using the official method 920.86 outlined in AOAC (1999). The Ankom fiber analyzer (Model ANKOM 220, USA) was employed for this purpose. A sample weighing 1.0 g was subjected to digestion in the fiber analyzer using dilute sulfuric acid (0.125 M H₂SO₄) for a duration of 30 minutes. The resulting residues were then washed with hot water. Subsequently, the residues were digested using dilute alkali (0.125

M KOH) for 30 minutes and again washed with hot water. The digested residues were dried in an oven at 105°C for 5 hours, cooled, and weighed. The residues were further subjected to incineration in a muffle furnace at 550°C for 2 hours, followed by cooling and weighing. The difference in weight between the residues before and after incineration was considered as the total fiber content.

$$\% \text{ Crude Fiber} = \frac{(\text{weight of sample residues before incineration} - \text{weight after}) (g)}{\text{weight of dry sample} (g)} \times 100$$

Carbohydrate

Carbohydrate was calculated as a percentage difference by the formula:

$$\% \text{ Carbohydrate} = 100 \% - (\% \text{ protein} + \% \text{ crude fiber} + \% \text{ crude fat} + \% \text{ Ash}).$$

Energy

The energy content was calculated using the Atwater's conversion factors. Thus energy values were obtained by multiplying % fat by factor 9, and % protein and % carbohydrate by factor 4 each (AOAC, 1999).

$$\text{Energy content} = [(\% \text{Carbohydrate} \times 4) + (\% \text{Fat} \times 9) + (\% \text{protein} \times 4)]$$

Mineral Composition

The mineral composition analysis of the samples was conducted with a slight modification following the method described by Jachimowicz *et al.* (2021). Initially, the samples were sieved using a 1.18mm sieve. Then, 0.5g of each sample was accurately weighed and placed in vials. For digestion, 2ml of nitric acid (HNO₃) and water in a 1:1 ratio were added to each sample. Additionally, 5ml of hydrochloric acid (HCl) in a 1:4 ratio with distilled water was also included. The vials were covered with watch glasses.

Samples in the vials were heated in a hot block (Hot block 150 model: SC-154-240) at 95°C and boiled for 30 minutes until it reached 85°C. After this process, the vials were removed and allowed to cool down to room temperature. Subsequently, distilled water was added to each vial, making the total volume 50ml. The content of each vial was transferred to test tubes and then analyzed using the Inductively Coupled Plasma Optical Emission Spectrometry machine (ICP-OES-5900 Agilent) with the model number (Agilent 5900 SVDV ICP-OES), Serial number MY2215CP04, Software version 7.6.0.12121, and firmware version 5590). The results were displayed on the computer.

Vitamin A

The Vitamin A content in the samples was determined through the UV-VIS Spectrophotometry method by measuring β -carotene, its precursor in plants. To extract β -carotene, a mixture of 50 mL acetone-hexane with 0.1% BHT was added to 5g of the sample. After shaking for 10 minutes, the mixture was centrifuged and the supernatant was separated using a funnel. The resulting solution was saponified with 25 mL of 0.5M methanolic potassium hydroxide, followed by shaking and settling for 30 minutes, with subsequent washing using 100 mL portions of distilled water. The aqueous layer was continuously discarded, and the extract was dried by filtering over anhydrous sodium sulfate.

The filtrate was concentrated at 45°C using a rotary evaporator and then reconstituted in methanol to a final volume of 50 mL. Standard solutions with various concentrations were prepared using 95% UV β -carotene. A stock solution of 100 $\mu\text{g}/\text{mL}$ was prepared by dissolving 0.01 g of β -carotene standard in 10 mL of hexane and then increased to 100 mL. The absorbance (A) of each concentration was measured at a wavelength of 450 nm using UV-VIS Spectrophotometry (Malla *et al.*, 2022).

Data analysis

All data was analyzed by Statistical Package for Social Sciences (SPSS) version 25. One way analysis of variance (ANOVA) test was performed following a post-hoc test with significant differences being determined at 5% level ($p < 0.05$). All results were expressed as mean \pm SD of duplicate values.

Results and Discussion

The chemical composition of complementary food blended with Moringa Leaf Powder (g/100 g dry weight) is summarized in Table 2. The composition was compared to the recommended daily intake for children aged 12-24 months by WHO (2005) and WHO/UNICEF (1998) (Table 3). The composition was also compared to the Tanzania standard for processed cereal based foods for infants and young children (TZS 180:2013) indicated by the Tanzania Bureau of Standards (Table 3).

Table 2: Proximate composition (g/100 g DM), energy (kcal/ 100 g) of Complementary food blended with Moringa Leaf Powder

Sample	Protein	Crude Fat	Fiber	Ash	Energy	Carbohydrates
PmDP	8.34±0.01 ^a	1.10±0.07 ^f	3.33±0.00 ^g	2.13±0.19 ^h	352±0.76 ⁱ	77±0.04 ^j
PmMD1	10.09±0.05 ^{bc}	1.05±0.07 ^f	2.82±0.25 ^g	1.99±0.25 ^h	347±2.60 ⁱ	74±0.87 ^k
PmMD2	10.19±0.00 ^{cd}	0.66±0.49 ^f	3.23±0.32 ^g	2.00±0.30 ^h	346±0.57 ⁱ	74±0.96 ^{jk}
PmMD3	11.07±0.28 ^e	0.98±0.01 ^f	3.01±0.02 ^g	2.23±0.25 ^h	347±1.42 ⁱ	73±0.60 ^k
PmMP1	9.92±0.06 ^c	1.29±0.04 ^f	2.85±0.34 ^g	2.03±0.09 ^h	348±2.22 ⁱ	74±0.51 ^k
PmMP2	9.72±0.03 ^b	1.25±0.03 ^f	3.10±0.31 ^g	2.04±0.08 ^h	348±0.28 ⁱ	74±0.10 ^k
PmMP3	10.58±0.00 ^d	1.38±0.06 ^f	2.69±0.58 ^g	1.99±0.21 ^h	350±0.39 ⁱ	74±0.24 ^k

Means ± SD, values within the same column with different superscript letters are significantly different from each other (p<0.05).

Fat

There was no significance difference (p>0.05) in crude fat among the samples. The values ranged from 0.66g/100g DM in PmMD2 to 1.38g/100g DM in PmMP3. Similar observations were made by Gebretsadikan *et al.* (2015) where fat content of OFSP-Soybean-Moringa leaf porridge was not significantly altered. The results were due to inclusion of high levels of OFSP (60%-85%) and low levels of Soybean (10%-30%). A study by Bello *et al.* (2022) reported that the fat content of the Pearl millet-OFSP-bean cookies was mainly from margarine that was added during preparations of the cookies. Pearl millet and OFSP have been reported to have low lipid content of 2.25% and 0.37% respectively (Adebiyi *et al.*, 2017; Kolawole *et al.*, 2020). This could also be the reason for low fat content observed in this study. A study by Haile *et al.* (2016) on composite flour of OFSP and bulla (*Enset ventricosum*) flours had fat content of 0.2% to 0.8% which is also within the set standard although it was not sufficient for the daily requirements for adults who were the targeted group. Although the values obtained in the current study were much lower than those given in Table 3 (Tanzania standard, TZS180:2013), they all met the set standard, and hence may be incorporated into infants food formulations complied.

Table 3: Recommended daily nutrient requirements for children aged 12-23 months

Nutrient	Requirement/day
Energy	900kcal
Protein	13g
Vitamin A	300µg
Iron	7mg
Zinc	3mg

Sources: WHO, (2005), WHO/UNICEF, (1998)

Ash

Ash refers to any inorganic material present in food or a residue that remains after heating and removing water. In current study, the ash content ranged between 1.99 to 2.13. There was no significance difference ($p>0.05$) in ash content among all samples in this study. According to Gebretsadikan *et al.* (2015) total ash content of OFSP-Moringa-Soybean porridge was positively influenced by OFSP and Moringa due to their high mineral content. A Study by Xu *et al.* (2020) support that potatoes have high mineral content. A study by Mohammed *et al.* (2016) on cookies reported that OFSP had higher contribution in mineral content of the cookies. Contrary to the present study, a study by Haile *et al.* (2016) on porridge made from composite flour of OFSP and bulla reported a significant difference in ash among the samples. The study also indicated an increase in crude ash content as the amount of OFSP increased. Another study by Olaitan *et al.* (2014) on quality evaluation of complementary food made from Pearl millet and Moringa leaf powder indicated an increase in the ash content as the amount of Moringa increased. All formulated samples under the current study were within the set levels as per TZS 180: 2013 indicated in Table 4.

Protein

Protein ranged between 8.34g/100g DM (52.1%) to 11.07g/100g DM (68.7%). There was a significant difference in protein content ($p<0.05$) among studied samples. This was from the samples with no Moringa to the sample that had the highest amount of Moringa of 3%. It was observed in this study that amount of increased protein content was directly proportional to Moringa Leaf Powder added This could be due to considerable amounts of protein that are found in Moringa. A study by Malla *et al.* 2021 reported 20% protein in the Moringa leaf. Other studies by Ntila (2017) and Kayi (2013) reported 20.47% and 30.3% of protein in dried Moringa leaf, respectively. Increased protein level in Moringa supplemented porridge could be due to significant amounts of protein found in Moringa leaves (Abuye *et al.*, 2003; Mellese *et al.*, 2009). A study by Olaitan *et al.* (2014) reported a significant difference in protein between the porridge samples upon addition of Moringa leaf powder. Increased protein content as the amount of Moringa was added in Moringa supplemented biscuits samples was reported by Hedhili *et al.* (2021).

The protein content in this study is within the acceptable limit set by TZS 180:2013 but was below the limits set by WHO/UNICEF.

Table 4: Requirements for processed cereal-based foods for infants and young children.

Characteristics	Requirements
Moisture content, % by mass, max.	Products for further processing 8.0
Total protein (quality at least 70% that of casein) by mass, min	14.0
Fat, % by mass, max.	8.5
Total carbohydrates, %by mass, min	60.0
Total ash, % by mass, max.	5.0
Ash insoluble in HCL, % by mass, max	0.05
Crude fiber (on dry basis), % by mass, max.	5
Vitamin A, IU/100g. min	500
Vitamin C mg/ 100g, min	25
Added Vitamin D, IU/100g.	25
Thiamine (as hydrochloride) mg/100g. min	300 to 800
Nicotine acid, mg/100g.min.	0.5
Calcium, mg/100g.max	1.0
Phosphorus mg/100g. min	25.0
Iron mg/100g, min	10

Source: TZS 180: 2013.

Crude Fiber

There was no significance difference ($p > 0.05$) in crude fiber content among the samples in this study. The fiber content in the samples was contributed by Moringa and OFSP. A study by Gebretsadikan *et al.* (2015) reported direct association of crude fiber with Moringa leaf powder and soybean. Moringa leaf powder is reported to have high fiber content than OFSP (Abuye *et al.*, 2003; Senanayake *et al.*, 2013). A study by Haile *et al.* (2016) reported an increase in fiber content as OFSP was added to Bulla flour. A study by Olaitan *et al.* (2014) on Pearl millet and

Moringa leaf powder porridge reported a significant difference in fiber content. There was an increase in fiber content among the porridge samples as Moringa leaf powder was added. As per standards sets by TZS 180:2013 for crude fiber content all samples in this study complied with the specification.

Carbohydrates

Although the carbohydrate content ranged from 73 g/100g DM to 77 g/100g DM, there was a significant difference ($p < 0.05$) in carbohydrates content between the control and all other samples except for sample PmMD2 among the samples in this study. The amount of carbohydrates from the product is contributed by both OFSP and pearl millet. Pearl millet contains 63-78% of carbohydrates (Taylor and Kruger, 2016). The amount of carbohydrates increased significantly in the product as OFSP amounts were added, this is due to its high starch content (Faber *et al.*, 2010). A study by Gebretsadikan *et al.* (2015) reported a limited impact on the changes of carbohydrates due to a narrow range of OFSP used in the mixture. Olaitan *et al.* (2014) reported a decrease in the carbohydrate content in the porridge samples as the more Moringa leaf powder was added in the formulation. This is because Moringa has less carbohydrate content compared to pearl millet and hence reduced carbohydrates content in the sample. The Tanzania Bureau of Standards has set a minimum of 60% for carbohydrate content and all samples in this study have met the standard.

Energy

The energy content of the samples ranged between 346 kcal/100g DM and 352 kcal/100g DM. There was no significant difference ($p > 0.05$) in energy among the samples in this study. This could have been attributed by limited variation in the amount of ingredients used to develop the products. A study by Haile *et al.* (2016) reported a significant difference in energy among the samples. The energy content was reported to increase as the amount of OFSP increased. A higher energy content of OFSP than that of bulla was reported to be responsible for the increased energy. All the samples in the study were below the energy set limits set by WHO/ UNICEF and Tanzania Bureau of Standards under TZS 180:2013.

Mineral Composition

The minerals and Vitamin A composition of Complementary food blended with Moringa Leaf Powder (g/100 g dry weight) is summarized in Table 5.

Iron

The Iron content of formulated products ranged from 4.5mg/100g to 6.3 mg/100g. There was no significant difference ($p > 0.05$) in iron among the samples in this study. The amount of Iron increased as the amount of Moringa increased in the sample. Similar observations were made by Gebretsadikan *et al.* (2015) in porridge samples where iron content increased as the amount of soybean and Moringa increased in the samples. Other studies by Abuye *et al.* (2003) and Mellese *et al.* (2009) had similar observations on the increase of iron content in the products as concentration of Moringa increased. Moringa leaves are a rich source of iron and other essential

minerals (Gandji *et al.*, 2018). A study by Govender & Siwela (2020) reported an increase in iron content from bread fortified with 5% and 10% Moringa compared to the unfortified bread.

According to WHO/UNICEF the set standard for iron content in complementary food for a 12–24-month child is 7mg/100g. CODEX (2006) has set a minimum of 4.8mg/100g while Tanzania Bureau of Standards under TZS 180:2013 has set a minimum of 10.87 mg/100g. All the samples in this study were below the limits set by WHO/UNICEF and TZS 180:2013. Six out of seven samples in the study reached the set limits made by CODEX (2006).

Zinc

The zinc content ranged from 1.46mg/100g to 1.81mg/100g among the samples. There was no significance difference ($p>0.05$) in zinc among samples in this study. An increase in zinc content in the samples that had more amount of Moringa was observed. This implicates that addition of Moringa leaf powder boost the amount of zinc in food. Similar observations were made by Roni *et al.* (2021) in the Moringa fortified cakes. According to CODEX (1991) amount of zinc in the weaning food should have a minimum of 2.42mg/100g. On the other hand, WHO/ UNICEF have set a standard of 3mg/day of zinc for complementary foods for children of 12-23 months of age. All the samples in this study were below the requirements set by both CODEX and WHO/UNICEF.

Calcium

The calcium content of the porridge samples increased with the increase in the concentration of Moringa. There was a significant difference ($p<0.05$) in calcium content among the samples. The maximum calcium content of 240g/100g was recorded from the sample PmMD3 and the minimum calcium content of 183g/100g was recorded from the control sample PmDP. The increase in calcium levels in the samples is supported by the presence of calcium levels in Moringa leaves (Kasolo *et al.*, 2010). A study by Sengev *et al.* 2013 reported a significant difference in calcium between bread samples fortified with Moringa and unfortified samples. According to FAO/WHO (1991) the minimum calcium content of weaning foods should be 435mg/100g.

Table 5: Mineral Composition (mg/100g DM), Vitamin A ($\mu\text{g}/100\text{g}$) of Complementary food blended with Moringa Leaf Powder

Samples	Iron	Zinc	Calcium	Vitamin A
PmDP	4.56±0.22 ^a	1.51±0.01 ^b	183±2.12 ^c	441±380 ^{ef}
PmMD1	5.77±1.09 ^a	1.73±0.04 ^b	237±2.12 ^d	308±535 ^e
PmMD2	5.85±0.63 ^a	1.81±0.16 ^b	236±2.82 ^d	357±368 ^{ef}
PmMD3	6.33±0.66 ^a	1.81±0.02 ^b	240±0.70 ^d	434±117 ^{ef}
PmMP1	5.7±0.14 ^a	1.47±0.67 ^b	202±1.41 ^e	381±680 ^{ef}
PmMP2	5.40±0.57 ^a	1.46±0.57 ^b	210±0.70 ^f	497±797 ^f

PmMP3	6.19±0.14 ^a	1.54±0.73 ^b	237±2.12 ^d	373±815 ^{ef}
-------	------------------------	------------------------	-----------------------	-----------------------

Means ± SD, values within the same column with different superscript letters are significantly different from each other (p<0.05)

Vitamin A

The β- carotene content among the samples ranged from 308µg to 497 µg per 100g. There was a significant difference (p< 0.05) in β- carotene content among samples. The increase in β- carotene is due to addition of both Moringa leaf powder and Orange fleshed sweet potato. OFSP contains a significant amount of Pro vitamin A (Bechoff, 2010). A study by Malla *et al.* (2021) reported that Moringa leaf powder contains higher contents of β- carotene than Millet. The study also reported that β- carotene content improved in the Moringa fortified flour compared to the unfortified samples. It was furthermore reported in this study that β- carotene is one of the target nutrient that increased due to addition of Moringa. A study by Boateng *et al.* (2019) supports that Moringa is a good fortificant to obtain enough amounts of Vitamin A. There were significant amounts of β- carotene in all the samples in this study due to addition of OFSP and Moringa, both of which have significant amounts of β- carotene. A study by Bello *et al.* (2022) reported that β- carotene content increased in cookies samples as OFSP increased. According to the Recommended Daily Intake of Vitamin A set by WHO/UNICEF for a 12–23-month-old, all the samples under this study reached the set limits.

Contribution of the nutrient's composition of the formulated porridge to the nutrient

Requirement of children aged 12-23 months.

Based on recommendations from UNICEF and WHO, children aged 12-23 months should be given three-quarter to a full cup (250mls) of complementary food per serving, with additional nutritious snacks offered 1-2 times per day as desired (WHO, 2009). However, according to FAO (2011), UNICEF (2010), and Brown (2003), children in this age group should ideally be given 3-4 servings a day due to their increased rate of growth and energy requirements as they continue to grow. If the porridge formulated in this study is provided to these children, it is assumed that children aged 12-23 months will consume a total of 1000mls per day (250mls per serving, 4 servings in a day).

The formulation labeled as "PmMP2," which was the most acceptable from the sensory evaluation panelists, could provide the following nutrients for children aged 12-23 months: 1,392 Kcal Energy, 38.8g crude protein, 1491 µg vitamin A, 21.6 mg Iron, and 5.6 mg Zinc. These nutrient levels exceed the recommended allowances for some nutrients targeted in this study. This indicates that the formulated porridge has the potential to provide beneficial nutritional contributions beyond the recommended amounts for children in this age group.

Another formulation that can be given to children aged 12-23 months is "PmMP3" which was the second most accepted formulation. If provided in 3 servings, a child will consume a total of 750mls per day. A child will obtain 1,050kcal Energy, 30g crude protein, 1119 µg vitamin A,

18.5mg Iron and 4.6mg Zinc from such diet. This formulation could also provide beneficial nutritional contributions beyond the recommended amounts for children in this age group. The Table 6 indicates amount that can be obtained in each nutrient and from every formulated mix when administered in 3 servings in a day.

Table 6: Distribution summary of nutrients in each sample obtained from 3 servings per day.

Amount of nutrients obtained from 3 servings per day						
Sample	Energy (Kcal)	Iron (mg)	Zinc (mg)	Calcium(mg)	Protein (g)	Vitamin A (µg)
PmDP	1056	13.5	4.5	549	24	1323
PmMD1	1041	17.1	5.1	711	30	924
PmMD2	1038	17.4	5.4	708	30	1071
PmMD3	1041	18.9	5.4	720	33	1302
PmMP1	1044	17.1	4.2	606	30	1143
PmMP2	1044	16.2	4.2	630	29	1491
PmMP3	1050	18.5	4.6	711	30	1119

All the samples in this study have shown to meet the stipulated standard set by WHO/UNICEF. Among all the samples the control sample PmDP provides the highest amount of energy. The sample PmMP2 provides the highest amount of Vitamin A. Iron, zinc, protein and calcium are obtained at a higher concentration in the porridge sample PmMD3. Although the sample PmMD3 has the highest amount of most nutrients it is among the least accepted sample in the sensorial aspect and therefore has the least chance to be taken.

Conclusion and Recommendation

The current study revealed that incorporating Moringa into the complementary food resulted in increased nutrient levels. For children aged 12-23 months to receive all the recommended daily nutrients, they would need to be provided with the amounts of meals as suggested by FAO/UNICEF. This research demonstrated that Moringa effectively improved the nutrient content of the complementary food. However, using more than 3% Moringa could further

enhance the nutrient levels. To address its bitter taste, leafy aroma, and deep green color, other foods can be added to mask these undesirable aspects of Moringa. In this way, children can consume the food without sensory issues while still benefiting from the nutritional advantages of Moringa.

REFERENCES

- Abuye, C., Urga, K., Knapp, H., Selmar, K., Omwega, A.M., Imungi, J.K. and Winterhalter, P. (2003), "A compositional study of *Moringa stenopetala* leaves", *East African Medical Journal*, Vol. 80 No. 5, pp. 247-250.
- Adebiyi, J.A., Obadina, A.O., Adebo, O.A., Kayitesi, E. 2017. Comparison of nutritional quality and sensory acceptability of biscuits obtained from native, fermented, and malted pearl millet (*Pennisetum glaucum*) flour. *Food Chemistry*, 232, 210-217
- AOAC (1999). *Official Methods of Analysis*. 16th Edition. Association of Official Analytical Chemists, Washington, D. C, USA. 54pp.
- Bechoff, A. (2010), "Investigating carotenoid loss after drying and storage of orange fleshed sweetpotato", PhD thesis, Greenwich University, pp. 75-102.
- Bello, F. A., Folademi, M. A., & Iwok, L. J. (2022). Development of pearl millet flour-based cookies supplemented with mung bean and orange fleshed sweet potato flours. *The Annals of the University Dunarea de Jos of Galati. Fascicle VI-Food Technology*, 46(1), 155-168.
- Bello, F. A., Folademi, M. A., & Iwok, L. J. (2022). Development of pearl millet flour-based cookies supplemented with mung bean and orange fleshed sweet potato flours. *The Annals of the University Dunarea de Jos of Galati. Fascicle VI-Food Technology*, 46(1), 155-168.
- Boateng, L., Nortey, E., Ohemeng, A. N., Asante, M., & Steiner-Asiedu, M. (2019). Sensory attributes and acceptability of complementary foods fortified with Moringa oleifera leaf powder. *Nutri Food Sci.*, 49(3), 393-406.
- CODEX (2006). Codex Alimentarius Commission: CODEX STAN 074-1981, REV.1-2006-Standards for processed cereal-based foods for infants and young children. 27pp.
- Faber, M., Laurie, S., Ball, A. and Andrade, M. (2010), "A crop-based approach to address vitamin a deficiency in Southern Africa – a regional manual for SADC", ARC-Roodeplaas, Pretoria, pp. 65-84.

- Falade, K. O., & Solademi, O. J. (2010). Modelling of air drying of fresh and blanched sweet potato slices. *International journal of food science & technology*, 45(2), 278-288.
- FAO (1985). Guidelines for development of supplementary foods for older infants and young children. Report of the 14th Session of the Codex Committee on Foods for Special Dietary Uses.
- FAO, IFAD and WFP (2015). The State of Food Insecurity in the World. FAO, Rome. 62pp.
- FAO, Rome. 10pp. Kikafunda, J. K., Abenakyo, L. and Lukwago, F. B. (2006). Nutritional and sensory properties of high energy/nutrient dense composite flour porridges from germinated maize and roasted beans for child-weaning in developing countries: a case for Uganda. *Ecology of Food and Nutrition* 45: 279-294
- Forsido, S.F., Rupasinghe, H.P.V. and Astatkie, T. (2013), "Antioxidant capacity, total phenolics and nutritional content in selected Ethiopian staple food ingredients", *International Journal of Food Sciences and Nutrition*, 64(8), 915-920.
- Gandji, K., Chadare, F. J., Idohou, R., Salako, V. K., Assogbadjo, A. E., & Kakai, R. G. (2018). Status and utilisation of Moringa oleifera Lam: A review. *African Crop Science Journal*, 26(1), 137-156.
- Gebretsadikan, T. M., Bultosa, G., Forsido, S. F., & Astatkie, T. (2015). Nutritional quality and acceptability of sweet potato–soybean–moringa composite porridge. *Nutrition & Food Science*, 45(6), 845-858.
- Govender, L., & Siwela, M. (2020). The effect of moringa oleifera leaf powder on the physical quality, nutritional composition and consumer acceptability of white and brown breads. *Foods*, 9(12), 1–17.
- Haidar, J., Muroki, N.M., Omwega, A.M. and Ayana, G. (2003), "Malnutrition and deficiency in breast feeding women in urban slum communities from Addis Ababa, Ethiopia", *East African Medical Journal*, 80(4), 191-194.
- Haile, A., Geribo, M., & Kinfu, E. (2016). Evaluation of porridge made from composite flour of orange-fleshed sweet potato and Enset (Bulla) flours. *Agriculture and Food Sciences Research*, 3(1), 37-44.
- Hasan, M. U., Malik, A. U., Ali, S., Imtiaz, A., Munir, A., Amjad, W., & Anwar, R. (2019). Modern drying techniques in fruits and vegetables to overcome postharvest losses: A review. *Journal of Food Processing and Preservation*, 43(12), e14280.
- Hedhili, A., Lubbers, S., Bou-Maroun, E., Griffon, F., Akinyemi, B. E., Husson, F., & Valentin, D. (2021). Moringa Oleifera supplemented biscuits: Nutritional values and consumer segmentation. *South African Journal of Botany*, 138, 406-414.

- Jachimowicz, K., Winiarska-Mieczan, A., Baranowska-Wójcik, E., & Bąkowski, M. (2021). Pasta as a Source of Minerals in the Diets of Poles; Effect of Culinary Processing of Pasta on the Content of Minerals. *Foods*, 10(9), 2131.
- Joint, F. A. O., WHO Expert Committee on Nutrition, & World Health Organization. (1971). *Joint FAO/WHO Expert Committee on Nutrition: eighth report, food fortification, protein-calorie malnutrition, Geneva, Switzerland, 9-18 November 1970*. World Health Organization.
- Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*, 4(9), 753–757.
- Kayi, K. K. (2013). *A study on Moringa oleifera leaves as a supplement to West African weaning foods*. Hamburg University of Applied Sciences. Incomplete ref
- Kolawole, F.L., Balogun, M.A., Oyeyinka, S.A., Adejumo, R.O., Sanni-Olayiwola, H.O. 2020. Effect of processing methods on the chemical composition and bio-accessibility of beta-carotene in orange-fleshed sweet potato. *Journal of Food Processing & Preservation*, e14538. Volume, issue, pages?
- Malla, J. K., Ochola, S., Ogada, I., & Munyaka, A. (2021). Nutritional Value and Sensory Acceptability of *M. oleif* Sengeev, A. I., Abu, J. O., & Gernah, D. I. (2013). Effect of *Moringa oleifera* leaf powder supplementation on some quality characteristics of wheat bread. *Food and nutrition sciences*, 4(3), 270.era Fortified Finger Millet Porridge for Children with Cerebral Palsy in Nairobi County, Kenya. *Journal of Food Research*, 10(5), 1-36
- Melesse, A., Bulang, M. and Kluth, H. (2009), "Evaluating the nutritive values and in vitro degradability characteristics of leaves, seeds and seedpods from *Moringa stenopetala*", *Journal of the Science of Food and Agriculture*, 89(2): 281-287
- Mosha T., Laswai S, Tetens I (2000). Nutritional composition and micronutrient status of home made and commercial weaning foods consumed in Tanzania. *Plant Foods Human Nutrition*, 55(3): 185-205.
- Muhimbula, HS. and Issa-Zacharia, A. (2010). Persistent child malnutrition in Tanzania: Risks associated with traditional complementary foods (A review). *African Journal of Food Science* 4 (11): 679 – 692.
- Nefale, F. E., & Mashau, M. E. (2018). Effect of germination period on the physicochemical, functional and sensory properties of finger millet flour and porridge. *Asian Journal of Applied Sciences*, 6(5).
- Ntila, S. L. (2017). *The potential of Moringa oleifera (Lam.) leaves for use in complementary foods to combat child food and nutrition insecurity among South African rural and peri-urban communities* [PhD Thesis]. University of KwaZulu-Natal.

- Olaitan, N. I., Eke, M. O., Uja, E. M., & Oja, E. A. (2014). Quality evaluation of complementary food formulated from Moringa Oleifera leaf powder and pearl millet (*Pennisetum Glaucum*) flour. *International Journal of Engineering and Science*, 3(11), 59-63.
- Omar, D. and Michael, H. (2008), "The food fortification formulator: technical determination of fortification levels and standards for mass fortification", Academy for Educational Development and United States Agency for International Development (USAID).
- Roni, R. A., Sani, M. N. H., Munira, S., Wazed, M. A., & Siddiquee, S. (2021). Nutritional Composition and Sensory Evaluation of Cake Fortified with Moringa oleifera Leaf Powder and Ripe Banana Flour. *Applied Sciences*, 11(18), 8474.
- Senanayake, S.A., Ranaweera, K.K.D.S., Gunaratne, A. and Bamunuarachchi, A. (2013), "Comparative analysis of nutritional quality of five different cultivars of sweet potatoes (*Ipomea batatas* (L) Lam) in Sri Lanka", *Food Science and Nutrition*,1(4):284-291
- Sengev, A. I., Abu, J. O., & Gernah, D. I. (2013). Effect of Moringa oleifera leaf powder supplementation on some quality characteristics of wheat bread. *Food and nutrition sciences*, 4(3), 270.
- Taylor, J. R. N., Kruger, J. (2016). Millets. *Encyclopedia of Food and Health* (pp. 748–757). Oxford: Academic Press.
- Tenagashaw, M. W., Kinyuru, J. N., Kenji, G. M., Melaku, E. T., & Susanne, H.- K. (2017). Nutrient density of complementary foods formulated from a blend of Teff, Soybean and Orange-fleshed Sweet Potato. *International Journal of Food Science and Nutrition Engineering*, 7(4),61-69.
- TZS 180 (2013).Tanzania Bureau of Standards: Specification for processed cereal-based foods for infants and young children-Specification, Tanzania Standards. 20pp
- UNICEF, (2009).Tracking Progress on Child and Maternal Nutrition. A survival and development priority: New York, NY 10017, USA
- WHO/FAO (2006), *Guidelines on Food Fortification with Micronutrients*, Geneva, pp. 3-328.