

## Original Research Article

# Identification of Ectoparasites and Endoparasites in Snakehead Fish (*Channa striata*) and Climbing Perch Fish (*Anabas testudineus*) Caught in Sebangau River

### ABSTRACT

This study aims to determine the types of ectoparasites and endoparasites that infect snakehead fish (*Channa striata*) and climbing perch fish (*Anabas testudineus*). Sampling was carried out at the Sebangau River while the identification was carried out at the Fish Quarantine Station, Quality Control and Safety of Fishery Products in Palangka Raya. Examination of ectoparasites includes mucus, gills, scales, and fish fins. Examination of endoparasites by observing the organs of the fish such as liver, meat, intestines and stomach. There are eight types of ectoparasites in snakehead fish, namely *Trichodina* sp, *Dactylogyrus* sp, *Gyrodactylus* sp, *Apiosoma* sp, *Tetrahymena* sp, *Aelosoma* sp, *Lamproglena* sp, and *Henneguya* sp, while there are four types of endoparasites in snakehead fish, namely *Camallanus* sp, *Procamallanus* sp, *Neoechynorynchus* sp, and *Senga* sp. There are five types of ectoparasites in the climbing perch fish caught in the Sebangau River, namely *Trichodina* sp, *Dactylogyrus* sp, *Aelosoma* sp, *Myxobolus* sp, and *Epistylis* sp. Meanwhile, there is only one type of endoparasite that attack snakehead fish, namely *Procamallanus* sp. The total prevalence of snakehead fish's ectoparasites was 81.7%, with an intensity level of 6.1 individual parasites per fish. The total prevalence of ectoparasites in climbing perch fish is 45%, with an intensity level of 5.4 individual parasites per fish. The total prevalence of snakehead fish endoparasites is 65%, with an intensity level of 9.8 individual parasites per fish. The prevalence of climbing perch fish's endoparasites is 31.7%, with an intensity level of 19 individual parasites per fish.

*Keywords: ectoparasites; endoparasite, snakehead fish, climbing perch fish, Sebangau river*

### 1. INTRODUCTION

The Sebangau River is a peat swamp river in Central Kalimantan which produces the black waters characteristics [14]. The habitat which influences the types of available fish. Snakehead fish (*Channa striata*) and climbing perch fish (*Anabas testudineus*) are the mostly fish caught in the Sebangau River. However, the surrounding area still has many settlements, agriculture, ponds or aquaculture, while the estuary area becomes a place for catching the fish. Disposal of liquid and solid waste is expected to cause the changes in quality or increase in environmental pollution and sources of disease [13]. Besides being obtained from catches, these two fish have also begun to be cultivated [3,8]. Taking fish directly in the waters for cultivation purposes certainly needs to consider the adaptation of these fish to be placed in a new environment. One of the things that determine the success of a fish farming business is the control of pests and diseases. The emergence of infections in the fish's body which will slowly affect the growth and productivity of the fish [5].

In general, diseases are caused by parasites. Parasites can cause serious problems to the point of causing mass death in cultivated fish [16]. In addition, parasites can cause economic losses and have an impact on human health when consumed raw or cooked imperfectly. Prevention of disease is the first thing of fish farmers need to do. Several studies on parasites in fish caught in the Sebangau River have been conducted [1,9,11]. However, overall there are no reports regarding to the type and distribution of ectoparasites and endoparasites that attack snakehead fish and climbing perch fish caught in the

Sebangau River, it causes the further research need to be conducted so the data can provide information of early disease control and further development of aquaculture.

## 2. MATERIALS AND METHODS

### 2.1 Materials

The tools used in this study consisted of scales, rulers, microscopes, dissecting sets, glass objects, cover glasses, pipettes, microscopes, preparations, trays, digital cameras and stationery. The materials used consisted of test fish and chemicals. The fish used in this study were snakehead fish (60 samples) and climbing perch fish (60 samples). The chemicals used were distilled water/aquadest (500 ml) and NaCl (500 ml).

### 2.2 Methods

The numbers of snakehead fish and climbing perch fish used were 60 samples for each. Samples were taken directly from the Sebangau river and brought to the laboratory for parasite examination. Sample fish were measured for the total of length (cm) and weight (g). Examination of ectoparasite infection, included: body surface (mucus), gills, scales, and fins of fish taken with a sterile scalpel. Then, after taking the fish's organs, it was placed on a glass object and dripped with distilled water/aquadest or NaCl, then covered with a cover glass and then to be observed under a microscope.

Examination of endoparasites was carried out by observing the organs of the fish such as liver, meat, intestines and stomach. Observation of fish endoparasites was carried out by dissecting parts of the fish to extract the internal organs, then taking and placing them on a glass object. dripped with distilled water/aquadest or NaCl solution and covered with a cover glass. The observation of parasites was carried out using a microscope and identification of parasites using a guidebook [7].

### 2.3 Statistical analysis

The data of ectoparasites and endoparasites that found from the results of examination sample were then recorded so that the number and types of ectoparasites and endoparasites were then counted. And to determine the level of parasites attack on fish, it could be analyzed by calculating the prevalence and intensity according to Kabata (1985):

$$P = N / n \times 100\%$$

Descriptions :

Q : Prevalence

N : Number of infected samples

n : Number of samples observed

while the intensity is calculated using the formula:

$$I = P / N$$

Descriptions :

I : Intensity of parasite attack (ind/tail)

P : The number of infecting ectoparasites (ind)

N : Number of infected samples (tails)

Parasite prevalence and intensity levels were analyzed using criteria that referred to [20], which are presented in the following Table 1.

**Table 1. Prevalence criteria of parasites attack**


No	Levels of Attack	Descriptions	Prevalences
1	Constantly	Very severe infection	100 - 99%
2	Almost constantly	Severe infection	90 - 99%
3	Ordinarily	Moderate infection	89 - 70%
4	Very often	Very frequent infection	69 - 50%
5	Normally	Common infection	49 - 30%
6	Often	Frequent infection	29 - 10%
7	Occasionally	Sometimes infection	9 - 1 %
8	Seldom	Rare infection	>1 - 0,1%
9	Very Rare	Very rare infection	>0,1 - 0,01%
10	Almost never	Never Infection	>P0,01%

**Table 2. Intensity criteria of parasites attack**

No	Level of Attack	Intensity
1	Very low	<1
2	Low	1 - 5
3	Moderate	6 - 55
4	Severe	51 - 100
5	Very severe	>100
6	Super Infection	>1000

### 3. RESULT AND DISCUSSION

Based on the identification results that carried out for 3 months (90 days) there were 60 snakehead fish and 60 climbing perch fish with a total of 120 fish observed. Then be found as many as 8 types of ectoparasite species and 4 types of endoparasite species in snakehead fish caught in the Sabangau river namely, *Trichodina* sp, *Dactylogyrus* sp, *Henneguya* sp, *Lamproglena* sp, *Tetrahymena* sp, *Epistylis* sp, *Apiosoma* sp, and *Gyrodactylus* sp ectoparasites, with the target of scales and mucus, fins, and gills were then carried out to identify the endoparasites obtained, namely *Camallanus* sp, *Procamallanus* sp, *Neoechinorhynchus* sp, and *Senga* sp targeting the liver, intestines, and meat. Then the identification of the climbing perch fish was carried out, there were 5 types of ectoparasitic species, namely *Trichodina* sp, *Dactylogyrus* sp, *Aelosoma* sp, *Myxobolus* sp, and *Epistylis* sp with the target of scales and mucus, fins, and gills [Fig. 1].



**Fig 1. Ectoparasites and endoparasites which found in snakehead fish and climbing perch fish: a. *Trichodina* sp.; b. *Epistylis* sp.; c. *Dactylogyrus* sp.; d. *Gyrodactylus* sp.; e. *Henneguya* sp.; f. *Lamproglene* sp.; g. *Tetrahymena* sp.; h. *Myxobolus* sp.; i. *Aelosoma* sp.; j. *Apiosoma* sp.; k. *Camallanus* sp.; l. *Procamallanus* sp.; m. *Neoechinorhynchus* sp.**

Henneguya sp. became the most dominant ectoparasite infecting snakehead fish in the Sebangau River, with gills being the most frequently infected organ. This can happen because the gill organ is an organ that often comes into direct contact with the environment, so it is susceptible to parasites. Stated that the high presence of parasites in the gills was caused by the direct contact with the water [21]. According to [18] stated that the infected fish by the parasite did not show severe clinical symptoms in the infected organs. This can happen because the fish's body had strong immune response and the influence of water quality in the location could still be tolerated by snakehead fish. Furthermore, *Trichodina* sp became the most dominant ectoparasite infecting climbing perch fish, with the gills being the most frequently infected organ. Infected fish got skin irritation, excessed mucus production, pale gills, gasping so that the fish often hangs on the surface of the water, decreased appetite, weak fish movements, damaged tail fin and reddish color due to broken capillaries in the fins [2]. *Trichodina* sp was very often found in the Sebangau River and was most commonly found in the gills. After the gills that were most often infected, the next part were the scales, mucus, and fins. Scales and mucus became the most numerous organs after gills that were attacked by parasites because scales and mucus became innate body protectors and become a separator between the internal and external environment of fish [4].

In the activity of identifying endoparasites in snakehead fish and climbing perch fish, the types of endoparasites that were found in snakehead fish were dominated by 4 types of endoparasites, namely *Camallanus* sp, *Procamallanus* sp, *Neoechinorhynchus* sp, and *Senga* sp.

*Procamallanus* sp became a parasite that often infects endoparasites in snakehead fish and climbing perch fish in the Sebangau River, the intestine being the organ that was often infected by this type of parasite. This can happen because the intestine is an organ that digests the foods consumed by fish from their environment. According to [12] that the intestine provides a source of nutrition for nematodes including blood, tissue cells, body fluids and food essences contained in the lumen of the small intestine.

Snakehead fish infected with the ectoparasite species which were identified in this study had an average length of 18.7 cm and a weight of 99.8 grams, and the type of endoparasite that attacked snakehead fish in this study had average of length 16.7 cm and weight of 97.3 grams. In climbing perch fish, ectoparasites attacked this fish with an average length of 13.6 cm and weight of 37.7, while endoparasites attacked the climbing perch fish in this study had an average length of 12.9 cm and weight of 36.2 grams. In some fish species, the larger the size/weight of the host, the higher the infection by certain parasites. The older hosts may contain larger numbers of parasites, although if becomes an adaptation, the host will tolerant to the parasites [15].

The results of the number of ectoparasites and endoparasites in snakehead fish and climbing perch fish caught in the Sebangau river (Table 3,4).

**Table 3. Results of the number of ectoparasites in snakehead fish and climbing perch fish**

Fish types	Parasites type	$\Sigma$ Parasite in the target organ (Ind/sel)			Total
		Fins	Scale/mucus	Gill	
Snakehead Fish	<i>Trichodina sp</i>		455	331	786
	<i>Dactylogyrus sp</i>			422	422
	<i>Henneguya sp</i>			2.250	2.250
	<i>Lamproglena sp</i>			63	63
	<i>Apiosoma sp</i>	68	14	47	119
	<i>Tetrahymena sp</i>		100		100
	<i>Epistylis sp</i>		9		9
	<i>Gyrodactylus sp</i>	1			1
TOTAL	8	69	578	3.113	3.750
Climbing Perch Fish	<i>Trichodina sp</i>	13	130	411	554
	<i>Dactylogyrus sp</i>			152	152
	<i>Aelosoma sp</i>		1		1
	<i>Mixobolus sp</i>			11	11
	<i>Epistylis sp</i>			7	7
TOTAL	5	13	131	581	725

Based on the table above, as many as 8 types of parasitic ectoparasites are found in snakehead fish in the Sebangau River with the number of parasites of 3.750 ind/cell, *Henneguya sp* being the ectoparasite type that most infects these fish, with a total of 2.250 ind/cell. Whereas *Epistylis sp* with 9 ind/cell and *Gyrodactylus sp* with 1 ind/cell, become the fewest ectoparasites that infect the fish. Ectoparasites are the most found in the gill organ, then followed by mucus, and then scales.

Then, as many as 5 types of ectoparasites are obtained from climbing perch fish in the Sebangau River with a total of 725 ind/cell, *Trichodina sp* being the most common type of ectoparasites found in the area, namely with a total of 554 ind/cell. *Epistylis sp* with 8 ind/cell and *Aelosoma sp* with 1 ind/cell are the fewest ectoparasites that infect the fish.

**Table 4. Results of the number of endoparasites in snakehead fish and climbing perch fish**

Fish types	Parasite types	$\Sigma$ Parasite in the target organ (Ind/sel)				Total
		Live r	Intestine	Meat	Stomach	
Snakehead Fish	<i>Camallanus sp</i>		14		1	15
	<i>Procamallanus sp</i>		12	1	3	16
	<i>Neoechynorynchus sp</i>		10	2		12
	<i>Senga sp</i>	1	4			5
TOTAL	4	1	40	3	4	48
Climbing Perch Fish	<i>Procamallanus sp</i>		41		1	42
TOTAL	1		41		1	42

Based on the table above, as many as 4 types of parasitic endoparasites are found in snakehead fish in the Sebangau River with the number of parasites are 48 ind/cell, *Procalamallanus sp* being the most common endoparasite that attacks snakehead fish, with the number of 16 ind/cell, followed by *Camallanus sp* with the number of parasites 15 ind/cell and *Neoechynorynchus sp* with the number of 12 ind/cell. Research conducted by [9] showed that the parasites of *Procalamallanus sp* and *Camallanus sp* mostly attacked snakehead fish in the intestinal organs. *Senga sp* is the least endoparasite that infected snakehead fish.

In the climbing perch fish, there is only 1 type of endoparasite that infects this fish in the Sebangau River area, namely *Procamallanus sp* with the number of parasites of 42 ind/cell.

The results of the prevalence (%) and intensity of ectoparasites and endoparasites in snakehead fish and climbing perch fish caught in the Sebangau river (Table 5,6).

**Table 5. Prevalence (%) and intensity of ectoparasites of snakehead fish and climbing perch fish**

Fish Types	Parasite Types	$\Sigma$ Observed fish	$\Sigma$ Infected Fish (tail)	Prevalence (%)	Intensity
Snakehead Fish	<i>Trichodina sp</i>	60	14	23,3	56,1
	<i>Dactylogyrus sp</i>	60	26	43,3	15,2
	<i>Henneguya sp</i>	60	10	16,7	225
	<i>Lamproglena sp</i>	60	16	26,7	3,9
	<i>Apiosoma sp</i>	60	13	21,7	9,2
	<i>Tetrahymena sp</i>	60	1	1,7	100
	<i>Epistylis sp</i>	60	2	3,3	4,5
	<i>Gyrodactylus sp</i>	60	1	1,7	1
Climbing Perch Fish	<i>Trichodina sp</i>	60	15	25	36,9
	<i>Dactylogyrus sp</i>	60	18	30	8,4
	<i>Aelosoma sp</i>	60	2	3,3	0,5
	<i>Mixobolus sp</i>	60	2	3,3	5,5
	<i>Epistylis sp</i>	60	20	33,3	0,4

From the table above, it can be seen that *Dactylogyrus sp* is the parasite that has the highest prevalence in snakehead fish and climbing perch fish ectoparasites with a prevalence of snakehead fish of 43.3% and climbing perch fish of 30%. *Dactylogyrus sp* is a parasite that has the highest prevalence of ectoparasites of snakehead fish and climbing perch fish, with a prevalence of 43.3% for snakehead fish and 30% for climbing perch fish, which is include in the common infection category, but the highest intensity of snakehead fish is *Henneguya sp*. with 225 individuals parasites per fish which makes this parasite into the super infection category, while the highest intensity in climbing perch fish is *Trichodina sp*, with 36.9 individual parasites per fish, makes this parasite into the moderate infection category. *Dactylogyrus sp*. parasite is an ectoparasites that attacks fish in peat swamp water such as javan combtail fish; climbing perch fish, red snakehead fish, ocellated fish, and catfish [1,6,10,11]. However, the highest intensity of snakehead fish is *Henneguya sp*, with 225 individual parasites per fish, while the highest intensity of climbing perch fish is *Trichodina sp*, with 36.9 individual parasites per fish.

**Table 6. Prevalence (%) and intensity of endoparasites of snakehead fish and climbing perch fish**

Fish Types	Parasites Types	$\Sigma$ Fish Observed	$\Sigma$ Infected Fish (tail)	Prevalence (%)	Intensity
Snakehead Fish	<i>Camallanus sp</i>	60	9	15	1
	<i>Procamallanus sp</i>	60	8	13,3	1,1
	<i>Neoechynorynchus sp</i>	60	9	15	0,5
	<i>Senga sp</i>	60	5	8,3	0,3

Climbing Perch Fish	<i>Procamallanus sp</i>	60	20	33,3	3,3
---------------------	-------------------------	----	----	------	-----

Camallanus sp and Neoechynorynchus sp are parasites that have the highest prevalence in snakehead fish endoparasites, which are 15% that makes these parasites into the frequent infections category. Procamallanus sp is the only endoparasite that attacks climbing perch fish which has a prevalence of 33.3%, include in the common infection category, but the highest intensity of snakehead fish is Camallanus sp, with 1.1 individual parasites per fish which makes this parasite in the category of low infection, while the intensity of Procamallanus sp in climbing perch fish is 3.3 individual parasites per fish which is also in the low infection category.

The mode of transmission of Neoechynorynchus sp is the same as other parasites of the Acanthopcephala phylum, that is through a definitive host of a crustacean containing Acanthella. Acanthella will attach to the surface of the intestine with the proboscis and will develop into adulthood [19].

Camallanus worm infection does not show clinical symptoms, but if it is heavily infected it can cause fish to become weak, have injuries to the intestines, anemia, and emaciation (thin and dry body) [17]. Camallanus sp. parasite is not specific because it is found in various types of fish with different water conditions such as seawater fish and freshwater fish [10].

The total ectoparasites of snakehead fish are 81.7% included in the moderate infection category, with an intensity level of 6.1 individual parasites per fish which is also in the moderate infection category. The prevalence of total ectoparasites in climbing perch fish is 45% in the common infection category, with an intensity level of 5.4 individual parasites per fish which is included in the low infection category.

Prevalence and total intensity of endoparasites of snakehead and parrot fish. Snakehead fish endoparasites are 65% include in the very frequent infection category, with an intensity level of 9.8 individual parasites per fish, include in the moderate infection category. The prevalence of parrot fish endoparasites was 31.7%, include in the common infection category, with an intensity level of 19 individual parasites per fish, include in the moderate infection category.

#### 4. CONCLUSION

There were eight types of ectoparasites in snakehead fish caught in the Sebangau River, namely Trichodina sp, Dactylogyrus sp, Gyrodactylus sp, Apiosoma sp, Tetrahymena sp, Aelosoma sp, Lamproglana sp, and Henneguia sp. Meanwhile, there were four types of endoparasites in snakehead fish, namely Camallanus sp, Procamallanus sp, Neoechynorynchus sp, and Senga sp. There were five types of ectoparasites in the climbing perch fish caught in the Sebangau River, namely Trichodina sp, Dactylogyrus sp, Aelosoma sp, Myxobolus sp, and Epistylis sp. Meanwhile, there was only 1 type of endoparasite that attacked the snakehead fish, namely Procamallanus sp. Henneguia sp became the most dominant ectoparasite infecting snakehead fish in the Sabangau River and Trichodina sp became the most dominant ectoparasite infecting climbing perch fish. Procamallanus sp was a parasite that often infected endoparasites in snakehead fish and climbing perch fish in the Sebangau River. The total prevalence of snakehead fish ectoparasites was 81.7%, with an intensity level of 6.1 individual parasites per fish. The total prevalence of ectoparasites in betok fish was 45%, with an intensity level of 5.4 individual parasites per fish. The total prevalence of endoparasites of snakehead fish was 65%, with an intensity level of 9.8 individual parasites per fish. The prevalence of climbing perch fish's endoparasites was 31.7%, with an intensity level of 19 individual parasites per fish.

#### REFERENCES

1. Agustinus F, Gusliany. Identification of Ectoparasites in Kapar Fish (*Belontia hasselti*) Reared in Plastic Lined Pond. *Ziraa'ah Journal*. 2020;45(2):103-110.
2. Ali SK, Koniyo Y, Mulis. Identification of Ectoparasites in Nile Tilapia fish (*Oreochromis nilotica*) in Limboto Lake of Gorontalo Province. *NIKE. Scientific Journal of Fisheries and Marine*. 2013;1(3):114-125.

3. Augusta TS, Pernando R. Technique of Snakehead (*Channa striata*) Spawning in Peatland Fish Farming Installation in Garung Village Pulang Pisau District. *Journal of Tropical Animal Science*. 2019;8 (1):13-18.
4. Dash S, Das SS, Samal J, Thatoi HN. Epidermal Mucus, a Major Determinant in Fish Health: a review. *Iranian Journal of Veterinary Research*. 2018;19(2):72-81.
5. Hasnidar H. Identification and Prevalence of Ectoparasites and Endoparasites in Mud Groper Fish (*Epinephelus tauvina*) of Talawi District, Batu Bara. *Regency. Aquaculture Journal*. 2021;17-25.
6. Herlina S. Ectoparasite Infection Rate in Toman Fish (*Channa micropeltes*). *Ziraa'ah Journal*. 2021;46(3):393-397.
7. Kabata Z. *Parasites and Disease of Fish Cultured in The Tropics*. Taylor & Francis. London & Philadelphia; 1985.
8. Maidie A, Sumaharjo SW, Asra M, Ramadhan, Hidayanto DW. Climbing Perch Fish Hatchery Development (*Anabas testudineus*) for Household Scale. *Aquaculture Media*. 2015;10(1):31-37.
9. Maryani, Monalisa SS, Sembiring IRB, Fransisco T. Identification of Endoparasites in Snakehead Fish (*Channa striata*) in Sebangau River Palangka Raya Central Kalimantan. *Journa of Aquaculture of River and Lake*. 2022;7(1):8-12.
10. Maulana DM, Muchlisin ZA, Sugito S. Intensity and Prevelency of Parasites on Climbing Perch Fish (*Anabas testudineus*) from Inland Waters of Northern Region of Aceh Province. *Scientific Journal of Fisheries and Marine Students of Unsyiah*. 2017;2(1):1-11.
11. Minggawati I, Agustinus F, Augusta TS, Pahawang CP, Francisco T. Identification and Prevalence of Ectoparasites and Endoparasites in Kerandang Fish *Channa pleuroptalma* and Catfish *Clarias batrachus* Captured from Sebangau River. *Indonesia Aquaculture Journal*. 2022;21 (2):152-160.
12. Muttaqin MZ, Abdul GN. The Prevalence dan Infection degree of *Anisakis* sp. in Digestive Tract of Red Snapper Fish (*Lutjanus malabaricus*) on Fish Auction of Brondong Lamongan. *Scientific and Arts Pomits Journal*. 2013;2(1):30-33.
13. Nirmalasari R. The Sebangau Water Quality Analysis at Kereng Bangkiray Harbor based on the Diversity and Composition of Phytoplankton. *Journal of Natural and Environmental Sciences*. 2018;9(17):48-58.
14. Nurseptiani, S, Kamal MM, Taryono, Surjanto D. Rights-Based Inland Water Fishery Management in Sebangau River, Sebangau National Park Central Kalimantan. *J. Kebijakan Sosek KP*. 2021;11(2):91-102.
15. Noble ER, Noble G.A. *Parasitologi*. Yogyakarta: Gadjah Mada University; 1989.
16. Purwanti R, Susanti R, Martuti NKT. Effect of Ginger Extract on Reducing the Number of Protozoan Ectoparasites in Tiger Grouper Fish. *Unnes J Life Sci*. 2012;1(2):70-77.
17. Rigby MC, Font WF, Deardorff TL. Redescription of *Camallanus Cotti* Fujita, 1927 (Nematoda: Camallanidae) from Hawai'i. *J. Parasitologi*. 1997;1161-1164.
18. Saputra Y, Rosita, Wirabakti MC, Djauhari R, Monalisa SS. Prevalence and Intensity of Ectoparasite Protozoa on Snakehead Fish (*Channa striata*) That Caught in Kahayan River. *AGRIENVI*. 2022;16(1):68-74.
19. Sobocka LE, Szostakowska B, MacKenzie K, Hemmingsen W, Prajsnar S, Eydal M. Genetic and Morphological Variation in *Echinorhynchus gadi* Zoega In Muller, 1976 (Acanthocephala: Echinorhynchidae) from Atlantic Cod *Gadus Morhua* L. *Journal of Helminthology*. 2012;86(1):16-25.
20. Williams EH, Williams LB. *Parasites of Offshore Big Game Fishes of Puerto Rico and The Western Atlantic*. Mayaguez: Antillean College Press;1996.

21. Yuli S, Harris H. Level of Ectoparasite Attack on Catfish (*Pangasius hypophthalmus*) Cultivated in Floating Net Cage at Musi River Palembang. *Journal of Fisheries Science and Aquaculture*. 2017;12 (2):50-57.

UNDER PEER REVIEW