

Original Research Article

Estimation of Magnitude of Heterosis for Grain yield and Related Traits in Sesame (*Sesamum indicum* L.)”

Abstract

Sesame (*Sesamum indicum* L.) is an important oilseed crop which is native to India and cultivated in around 2.3 million ha across its several states. India stands way below in terms of productivity compared to several other countries across the globe. Being an often cross pollinated crop, heterosis even in a smaller magnitude could be exploited in sesame in varietal development. Hence, the present study was performed with the objective of estimating the extent of relative heterosis, heterobeltiosis and standard heterosis in eight cross combinations for important yield attributing traits in sesame in order to select promising cross combinations for utilization in heterosis breeding and/or selection of superior segregants in segregating generations. Eight cross combinations derived out of crossing high yielding parents viz., TMV (SV) 7 and VRI (SV) 2 as female and E8, DSS 9, D 5 and Prachi as male in L X T fashion were utilized for the analysis keeping TMV (SV) 7 as the Standard parent. The analysis showed significant differences between the genotypes for all the characters studied. The hybrids showed Relative/ Mid parental heterosis ranging from 2.02 to 26.05 per cent; heterobeltiosis from -4.55 to 15.85 per cent and standard heterosis from -11.27 to 15.85 per cent for grain yield /ha. Mid Parental Heterosis (RH) was found to be positive and significant for the characters no. of branches/plant, no. of capsules / plant, grain yield (kg/ha) in all the hybrid combinations except VRI (Sv) 2 / Prachi. Based on heterobeltiosis, the hybrids TMV (SV) 7 / E 8, TMV (SV) 7 / DSS 9, TMV (SV) 7 / DS 5, VRI (SV) 2 / DS 5 could be selected since they showed significant heterobeltiosis for important yield contributing characters viz., no. of branches / plant, no. of capsules /plant and grain yield. With regards to the Standard heterosis for grain yield and contributing characters like no. of branches/plant, no. of capsules/plant and grain yield/ha the hybrids viz., TMV (SV) 7 / E 8, TMV (SV) 7 / DS 5 and TMV (SV) 7 / DSS 9 showed positive and significant heterosis. Considering all the three types of heterosis, the hybrids TMV (SV) 7 / E 8, TMV (SV) 7 / DS 5 and TMV (SV) 7 / DSS 9 may be selected for yield enhancement in Sesame.

Key words: Heterobeltiosis, Productivity, Relative Heterosis, Standard Heterosis, Yield enhancement.

Introduction

Sesame is described as the “Queen of oilseeds” due to its high oil content (38-54%), protein (18-25%), calcium, phosphorus, oxalic acid and excellent qualities of seed oil and meal (Prasad, 2002). *Sesamum* oil has long shelf life due to the presence of lignans (sesamin, sesamol and sesaminol), which have remarkable antioxidant function (Gauthaman and Saleem, 2009). Whole sesame seeds are rich source of mammalian lignin precursors (sesamin, sesamol and sesaminol) and have been suggested to have potential anticancer effects (Coulman *et al.*, 2005). Sesame oil, is also used as a solvent, oleaginous vehicle for drugs, skin softener, and used in the manufacture of margarine and soap (Morris, 2002). It is resistant to drought, short in maturity duration and suitable for wide range of cropping systems. Sesame seed is an important source of edible oil and is also widely used as a spice because of its ease of extraction and its great stability. In India sesame occupies a place of prominence among oilseeds.

The annual area put under Sesame in India is about 2.3million hectares (13.1 % per cent of the world hectareage) and the total production is nearly 81lakh tonnes (Agricultural Statistics at a glance,2021). India ranks third in sesame production and second in area after Sudan in the world with productivity of 502 kg/ha (<https://aicrp.icar.gov.in/sesame/about-us/>). Though the progress achieved in sesame in terms of cultivated area and production is quite high, its productivity in India is the lowest when compared to the world average. The major constraints identified for most of the countries including India are, instability in yield, lack of wider adaptability, drought, non synchronous maturity *etc.* There are few studies on the genetics of sesame as well as on the improvement of genetic variation and root system. In-depth research on sesame is still necessary to further increase the yield and quality of sesame and to improve the related traits of sesame (Wei *et al.*, 2022). Heterosis breeding is a potential technique to improve yields in sesame (Nayak *et al.*, 2017). Hybrid vigor in sesamum even a small magnitude for individual component, have additive or synergistic effects on end product (Sasikumar and Sardana, 1990). The magnitude of heterosis provides a basis for genetic diversity and a guide to the choice of parents for developing superior F₁ hybrids, so as to exploit hybrid vigour and/or for building better gene pools to be employed in population improvement (Jatothu *et al.*, 2013). Besides, heterotic crosses may be amenable for selection of high yielding transgressive segregants in F₂ and follow up selfing generations (Tripathy *et al.*, 2016). Hence the present investigation was

undertaken to study the extent of heterosis present in select sesame hybrids developed for yield enhancement.

Materials and Methods

The present investigation is carried out in the Research farm at Oilseeds Research Station, Tindivanam during Summer, 2020. Oilseeds Research Station, Tindivanam lies in the North Eastern Zone of Tamil Nadu at an altitude of 45.6 above MSL at lies between 12.5°N latitude and 79.5°E longitude. The average annual rainfall is 1158 mm and the temperature ranges from 19-21 degrees as minimum to 33-42 degrees as maximum. The soil is categorized as sandy loam with a pH of 7-7.5. The. The cropping pattern followed in ORS, Tindivanam is Groundnut-Groundnut/Sesame-Sesame or Castor-Sesame with the main emphasis to develop high yielding, biotic/abiotic stress resistant groundnut and sesame varieties. The experimental material included eight hybrids evolved by crossing two female and 4 male parents in line x tester fashion. The female parents taken for the study were the high yielding varieties developed by TNAU viz., TMV (Sv) 7 and VRI (Sv) 2 while the male parents were E8, DSS 9, DS 5 and Prachi which showed good general combining ability with the female parents selected from the All India Coordinated Research trials. The crosses were performed in L x T fashion during Rabi and Summer, 2019 giving suitable isolation distance.

Laying out of experimental plot

The resultant eight hybrids along with the 6 parents were sown directly in plot size of 4 x 3 m in a Randomized Block Design replicated thrice. All the package of practices was followed to raise a good crop. Observations other than days to 50% flowering (days) were recorded at the time of harvest in 10 plants selected at random per replication in each cross combination and parent viz., plant height (cm), no. of branches/plant (No.) and no. of capsules/plant (No.) were recorded. The number of days taken from sowing for 50% of the plants in a plot to flower is recorded as days to 50% flowering. The height from the base to tip of the plant is measured as plant height (cm). The number of primary branches originating from the main stem is recorded as no. of branches/plant (No.) and the number of capsules is counted as no. of capsules/plant (No.). The plot yield (kg/plot) of the hybrids and the parents recorded in each replication was converted to per hectare yield (kg/ha). In order to manage the grain shattering and reduction in yield, the individual plots were harvested at physiological maturity where 75% of the plants in a plot is

mature. The mean of characters recorded in all the ten plants selected at random per replication were utilized for L x T statistical analysis as per Kempthorne (1957) in the statistical package TNAUSTAT. The per cent of heterosis of all the three categories viz., relative or mid parent heterosis, heterobeltiosis or better parent heterosis (Fonseca and Patterson, 1968) and standard heterosis (Meredith and Bridge, 1972) were worked out based on the mean values as suggested by. For working of standard heterosis, the cosmopolitan high yielding variety released by TNAU in 2009 and one of the parents used in the study viz., TMV (Sv) 7 was utilized.

Results and Discussion

The analysis of variance for different characters showed significant differences between the genotypes for all the characters studied (Table 1). The mean performance of the parents and hybrids were tabulated in Table 2.

Table 1. Analysis of Variance for yield parameters in Sesame

Sl. No.	Source	df	Mean Squares				
			Days to 50% flowering (days)	Plant Height (cm)	No. of Branches/ plant (no.)	No. of Capsules/ plant (no.)	Yield (kg/ha)
1.	Genotypes	13	10.53*	405.41*	5.95*	703.24*	23514.13*
2.	Replication	2	1.45	6.88	0.02	3.24	280.27
3.	Error	26	1.84	23.69	0.20	27.45	902.70

Table 2. Mean performance of the cross combinations and parents

Genotypes	Days to 50 % flowering (days)	Plant height (no.)	No. of branches / plant (no.)	No. of capsules / plant (no.)	Yield/ha (kg)
Hybrids					
TMV (Sv) 7 / E8	32	154.0	7.5	101.3	913.9
TMV (Sv) 7 / DSS 9	35	153.1	8.6	79	847.2
TMV (Sv) 7 /	33	141.8	6.5	75.3	880.6

DS 5					
TMV (Sv) 7 / Prachi	34	137.2	5.7	62	794.5
VRI (Sv) 2 / E8	33	141.6	5.9	73	800.0
VRI (Sv) 2 / DSS 9	34.7	138.3	5.1	61.9	705.6
VRI (Sv) 2 / DS 5	32.7	142.3	5.7	84.5	802.8
VRI (Sv) 2 / Prachi	31	152.2	5.3	58.2	700.0
Parents					
TMV (Sv) 7	33	145.2	5.3	63.3	788.9
VRI (Sv) 2	32.3	129.6	4.9	64.7	733.3
E8	30	140.4	5.1	55.3	661.1
DSS 9	28	123.4	3.5	47.3	647.2
DS 5	32	130.6	3.3	59.3	680.5
Prachi	30.7	112.8	4.4	42.7	638.9
SEd	1.1	4.0	0.37	4.3	24.5
CD (0.05)	2.3	8.2	0.76	8.8	50.5

The per cent heterosis under the three categories viz., relative heterosis, heterobeltiosis and standard heterosis is presented in Table 3. Relative/Mid parental heterosis ranged from -1.59 (VRI (Sv) 2 / Prachi) to 14.92 per cent (VRI (Sv) 2 / DSS 9) in days to 50% flowering; from -5.49 (TMV (Sv) 7 / Prachi) to 17.43 per cent (VRI (Sv) 2 / Prachi) in plant height; 15.11 (VRI (Sv) 2 / Prachi) to 96.95 per cent (TMV (Sv) 7 / DSS 9) in no. of branches / plant; 8.51 (VRI (Sv) 2 / Prachi) to 70.79 per cent (TMV (Sv) 7 / E8) in no. of capsules/plant and from 2.02 (VRI (Sv) 2 / Prachi) to 26.05 per cent (TMV (Sv) 7 / E8) in grain yield. Heterobeltiosis ranged between -4.12 (VRI (Sv) 2 / Prachi) and 7.22 per cent (VRI (Sv) 2 / DSS 9) for days to 50% flowering, -5.47 (TMV (Sv) 7 / Prachi) and 17.43 per cent (VRI (Sv) 2 / Prachi) for plant height, 5.48 (VRI (Sv) 2 / DSS 9) and 63.29 per cent (TMV (Sv) 7 / DSS 9) for No. of branches/plant, -9.95 (VRI (Sv) 2 / Prachi) and 60.0 per cent (TMV (Sv) 7 / E8) for no. of capsules/plant and between -4.55 (VRI (Sv) 2 / Prachi) and 15.85 per cent (TMV (Sv) 7 / E8) in grain yield / ha.

Wide range of relative heterosis and heterobeltiosis also existed in the study conducted by Rajput *et al.* 2017 for yield, no. of capsules /plant, no. of branches and no. of seeds/capsule. (Table 3)

Table 3. Relative heterosis, Heterobeltiosis and Standard heterosis for yield parameters in Sesame

Genotypes/ Hybrids	Days to 50% flowering			Plant Height (cm)			No. of Branches/plant			No. of Capsules/plant			Yield (kg/ha)		
	RH	HB	SH	RH	HB	SH	RH	HB	SH	RH	HB	SH	RH	HB	SH
TMV (Sv) 7 / E8	1.59	-3.03	-3.03	7.87**	6.11*	6.11*	45.34**	43.04**	43.04**	70.79**	60.00**	60.00**	26.05**	15.85**	15.85**
TMV (Sv) 7 / DSS 9	14.75**	6.06	6.06	14.03**	5.46	5.46	96.95**	63.29**	63.29**	42.77**	24.74**	24.74**	17.99**	7.39*	7.39*
TMV (Sv) 7 / DS 5	1.54	0.00	0.00	-2.30	-2.30	-2.30	50.39**	22.78**	22.78**	22.83**	18.95**	18.95**	19.85**	11.62**	11.62**
TMV (Sv) 7 / Prachi	6.81*	3.03	3.03	-5.49	-5.49	-5.49	17.24*	7.59	7.59	16.98*	-2.11	-2.11	11.28**	0.70	0.70
VRI (Sv) 2 / E8	5.88	2.06	0.00	9.23**	0.83	-2.46	19.06**	16.34*	12.66	21.61**	12.84	-2.11	14.74**	9.09*	1.41
VRI (Sv) 2 / DSS 9	14.92**	7.22*	5.05	6.69*	6.69*	-5.49	23.20**	5.48	-2.53	10.54	-4.28	-2.26	2.21	-3.79	-10.56**
VRI (Sv) 2 / DS 5	1.55	1.03	-1.01	9.40**	9.01**	-1.95	39.84**	17.81*	8.86	36.29**	30.67**	33.42**	13.56**	9.47**	1.76
VRI (Sv) 2 / Prachi	-1.59	-4.12	-6.06	17.43**	17.43**	4.87	15.11*	9.59	1.27	8.51	-9.95	-8.05	2.02	-4.55	-11.27**

RH- Relative heterosis; HB- Heterobeltiosis; SH- Standard heterosis

Standard heterosis varied between -6.06 (VRI (Sv) 2 / Prachi) and 6.06 per cent (TMV (Sv) 7 / DSS 9) in days to 50% flowering, -5.49 (TMV (Sv) 7 / Prachi and VRI (Sv) 2 / DSS 9) and 6.11 per cent (TMV (Sv) 7 / E 8) in plant height, -2.53 (VRI (Sv) 2 / DSS 9) and 63.29 per cent (TMV (Sv) 7 / DSS 9) in no. of branches / plant, -8.05 (VRI (Sv) 2 / Prachi) and 60.0 per cent (TMV (Sv) 7 / E 8) in no. of capsules / plant and between -11.27 (VRI (Sv) 2 / Prachi) and 15.85 per cent (TMV (Sv) 7 / E 8) in yield / ha. Significant standard heterosis has been reported for days to 50% flowering, plant height, no. of branches/plant, no. of capsules/plant and grain yield by Virani *et al.* 2017 and Shekawat *et al.* 2014.

Earliness is preferred for varieties and hybrids in general in sesame. Hence for the character days to 50% flowering alone negative heterosis is considered and for all other characters positive heterosis is considered for selection of hybrids. In the present study, for days to 50% flowering none of the hybrids showed negatively significant heterosis in all the three forms *viz.*, Relative heterosis, heterobeltiosis and standard heterosis. Hence, none of the hybrids can be selected for days to 50% flowering with negative heterotic values.

Positively significant heterosis is exhibited in several hybrids for yield attributing characters. Mid Parental Heterosis (RH) was found to be positive and significant for the characters no. of branches/plant, no. of capsules / plant, grain yield (kg/ha) in all the hybrid combinations except VRI (Sv) 2 / Prachi. For plant height, all hybrids except TMV (Sv) 7 / Prachi and TMV (Sv) 7 / DS 5 showed significant relative heterosis. Significant positive relative heterosis was noticed in eleven out of forty five crosses for no. of primary and secondary branches/ plant, no. of capsules/plant, no. of seeds/ capsule and single plant yield in the study conducted by Meenakumari and Ganesamurthy (2017). This indicates that the cross combinations were heterotic for the yield characters and shows good scope for utilization as commercial hybrids as well as in varietal improvement.

Better parental heterosis is considered for selection of hybrids in many cases as this may benefit for yield improvement if the cross combination is for improvement of specific characters choosing one of the parents specifically showing better performance. Heterobeltiosis / better parental heterosis was positive and significant for the character plant height in hybrids TMV (Sv) 7 / E8, VRI (Sv) 2 / DSS 9, VRI (Sv) 2 / DS 5 and VRI (Sv) 2 / Prachi; for no. of branches / plant and grain yield in cross combinations *viz.*, TMV (Sv) 7 / E8, TMV (Sv) 7 / DSS 9, TMV (Sv) 7 / DS 5, VRI (Sv) 2 / E 8 and VRI (Sv) 2 / DS 5 while hybrids TMV (Sv) 7 / E 8, TMV(Sv) 7 / DSS 9, TMV(Sv) 7 / DS 5 and VRI (Sv) 2 / DS 5 showed positive and significant heterobeltiosis for no. of capsules / plant. Monpara and Pawar (2016) also reported significant and positive heterobeltiosis in eight cross combinations for yield attributing traits like primary branches/plant, capsules/plant, seeds/capsule, seed yield and harvest index in sesame. Tripathi *et al.* (2016) identified five superior cross combinations based on heterobeltiosis for seed yield, three cross combinations for capsule number and also reported the significant contribution of capsule number in determining the seed yield in sesame.

Accordingly the hybrids TMV (Sv) 7 / E 8, TMV (Sv) 7 / DSS 9, TMV (Sv) 7 / DS 5, VRI (SV) 2 / DS 5 could be selected since they showed significant heterobeltiosis for important yield contributing characters viz., no. of branches / plant, no. of capsules /plant and grain yield.

Standard heterosis is generally considered for varietal improvement as this is economically useful to the farmers and the commercial success of the new variety depends on the yield advantage of this variety over the ruling variety. Kadambavana sundaram (1980) suggested that heterotic expression over standard variety should be given due importance for exploitation of commercial hybrid. Considering the standard heterosis for plant height, only one hybrid TMV (Sv) 7 / E 8 could be adjudged as the best hybrid showing significant heterosis. For grain yield characters like no. of branches/plant, no. of capsules/plant and grain yield/ha the hybrids viz., TMV (Sv) 7 / E 8, TMV (Sv) 7 / DS 5 and TMV (Sv) 7 / DSS 9 are showing significant heterosis compared to other hybrids and hence can be utilized for exploitation of commercial heterosis. Saravanan and Nadarajan (2002) reported high Standard heterosis for no. of primary branches/plant, no. of capsules/plant, 1000 seed weight and single plant yield in sesame. Shobana Rani *et al.* (2015) assessed heterosis among thirty eight cross combinations and reported two hybrids Swetha thil x VS 07-023 and Swetha thil x JL SEL 05-3 exhibiting significantly superior standard heterosis over three standard checks. High standard heterosis has been recorded in five cross combination over the standard parent by Chaudhari *et al.* 2017.

Considering all the three types of heterosis, the cross combination VRI (Sv) 2 / Prachi is found to exhibit lowest values of relative heterosis, heterobeltiosis and Standard heterosis for economically important characters viz., no. of capsules/plant and grain yield/ha while the cross combination TMV(Sv) 7 / E 8 exhibited highest relative heterosis, heterobeltiosis and Standard heterosis for the characters viz., no. of capsules/plant and grain yield/ha among the eight hybrids studied. Similarly the hybrids viz., TMV (Sv) 7 / DSS 9 and TMV (Sv) 7 / DS 5 also showed significant relative heterosis, heterobeltiosis and standard heterosis for no. of branches/plant, no. of capsules/plant and grain yield/ha. Hence the hybrids TMV(Sv) 7 / E 8, TMV (Sv) 7 / DSS 9 and TMV (Sv) 7 / DS 5 are adjudged as the best heterotic combinations exhibiting all the three types of heterosis for grain yield and its contributing characters. Sushila *et. al.*, 2020 reported significant positive average heterosis in six crosses, heterobeltiosis in five crosses and standard heterosis in seven and also concluded positive and highly significant values for all three kinds of heterosis in two crosses, RT-351×TKG-22 and RT-54 × TKG-22 and could be selected as

promising hybrids for improving seed yield and contributing characters in sesame. Significant and positive standard heterosis and heterobeltiosis for seed yield per plant in sesame have also been reported by earlier workers like Kumar and Ganesan, (2004), Kumar *et al.* (2015), Shoba Rani *et al.*, (2015) and Karande *et al.* (2018). These hybrids also showed high *per se* performance in comparison to all other hybrids and their respective parents. Dela and Sharma (2019) also selected four heterotic combinations on the basis of heterosis and mean performance and also opined that the presence of large genetic diversity among the parents and also unidirectional distribution of allelic constitution contributing towards desirable heterosis.

Heterosis is generally studied for assessing the breeding value of the cross combinations. It is also studied as a measure of varietal superiority performed in selection of superior hybrids in any variety release programme. In general, genetically diverse parents are believed to put forth most heterotic combinations. Hence the superior hybrids *viz.*, TMV(Sv) 7 / E 8, TMV (Sv) 7 / DSS 9 and TMV (Sv) 7 / DS 5 identified based on heterosis may be utilized for further improvement in yield and its contributing characters through heterosis breeding or in identification of superior segregants in further generations.

Conclusions

The highly heterotic combinations are expected to through better transgressive segregants in the future generations and considered to be of high value than the low heterotic combinations. Hence considering all the three types of heterosis, the hybrids TMV (Sv) 7 / E 8, TMV (Sv) 7 / DS 5 and TMV (Sv) 7 / DSS 9 were adjudged to be good for enhancement of grain yield and its attributing characters in Sesame.

Reference

1. Agricultural Statistics at a glance, (2021). [https://eands.dacnet.nic.in/PDF/Agricultural Statistics at a Glance](https://eands.dacnet.nic.in/PDF/Agricultural_Statistics_at_a_Glance_2020-21) 2020-21.
2. Chaudhari, G.B., Naik, M.R., Anarase, S.A. and Ban, Y.G. (2015). Heterosis studies for quantitative traits in sesame (*Sesamum indicum* L.). *Electron. J. Pl. Breed.*, **6(1)**:218-224
3. Coulman, K.D., Liu, Z., Hum, W.Q., Michaelides, J and Thompson, L.U. (2005). Whole sesame seed is as rich a source of mammalian lignan precursors as whole flaxseed. *Nutrition and Cancer* **52(2)**, 156-165.

4. Dela, G.J. and Sharma, L.K. (2019). Heterosis for seed yield and its components in sesame (*Sesamum indicum* L.). *J. of Pharmacognosy and Phytochemistry*, **8(4)**: 1345-1351.
5. Fonseca, S. and Patterson, F.L. (1968). Hybrid vigour in seven parent diallel cross in common wheat (*Triticum aestivum* L.). *Crop Sci.*, 2: 85-88.
6. Gauthaman, K. and Mohamed Saleem, T.S. (2009). Nutraceutical Value of Sesame Oil. *Phcog. Rev.* **3**, 264-269.
7. Jatothu, L. Dangi, K.S. and Kumar, S.S. (2013). Evaluation of sesame crosses for heterosis of yield and yield attributing traits. *J. of Trop. Agric.*, **51 (1-2)** : 84-91.
8. Kadambavana Sundaram, M. (1980). Combining ability as related to gene action in cotton (*G. hirsutum* L.) Ph.D., Thesis, Tamil Nadu Agrl. Univ., Coimbatore.
9. Karande, G.R., Yamgar, S.V., Waghmode, A.A., Wadikar., P. B. (2018). Exploitation of heteosis for yield and yield contributing character in sesame (*Sesamum indicum* L.). *Int. J Curr. Microbiol. App. Sci.*; **7(2)**:299-308.
10. Kempthorne, O. 1957. An introduction to genetic statistics. John Wiley and Sons Ins., New York, pp: 458-471.
11. Kumar, N., Tikka, S.B.S. and Bhagirathram, Dagla, M.C. (2015). Heterosis studies for agronomic trait under different environmental conditions in sesame (*Sesamum indicum* L.). *Electronic J Plant Breed.*, **6(1)**:130-140.
12. Kumar, P.S., Ganesan J. (2004). Heterosis for some important quantitative traits in sesame (*Sesamum indicum* L.). *Agric. Sci. Digest.* **24(4)**:292-294.
13. Meena Kumari, B. and Ganesamurthy. K. (2017). Study on the exploitation of heterosis in sesame (*Sesamum indicum* L.). *Electron. J. of Plant Breed.*, **8(2)**: 712-717.
14. Meredith, W.R., and Bridge, R. R. (1972). Heterosis and gene action in cotton, (*G. hirsutum* L.). *Crop Sci.* **12**:304-310.
15. Monpara, B.A. and Pawar, A.K. (2016). Evaluation of hybrids for heterosis breeding in sesame (*Sesamum indicum* L.). *Electron. J. of Plant Breed.*, **7(4)**: 1183-1187
16. Morris, J.B. (2002). Food, Industrial Nutraceutical Uses of Sesame Genetic Resources. (ASDHS Press, 2002) pp.153-156
17. Nayak, A. J, Patel, S. R. and Shrivastva, A. (2017). Heterosis studies for yield and its components traits in sesame (*Sesamum indicum* L.). *An International e- Journal*, **6(1)**: 38-48.

18. Prasad, R. (2002). Text Book of field Crops Production (Indian Council of Agricultural Research, New Delhi, 2002) p.821
19. Rajput, S.D., P.N.Harer and N.S.Kute. (2017). Heterosis and its relation with combining ability in Sesame. (*Sesamum indicum* L.). for quantitative traits. *Int. J. Curr. Res.*, **9(09)** : 56971-56973.
20. Saravanan, S. and Nadarajan, N. (2002). Studies on heterosis in Sesame (*Sesame indicum* L.). *Indian J.Genet.*,**62(3)**:271-272(2002)
21. Sasikumar B and Sardana, S.(1990). Heterosis for yield and yield components in sesame. *Indian J. Genet.*, **50(1)**: 87–88.
22. Shekhawat, N.R., Macwana, S.S., Choudhary, R. and Patel, B. R. (2014). Line x Tester analysis in sesame (*Sesamum indicum* L.). *Intern. Quarterly J. of Life Sci., The Bioscan*, **9(4)**:1657-1660, 2014 (Suplmnt. on Gen. and Pl. Breed.).
23. Shobana Rani, T., Kiran Babu, T., Madhukar Rao, P., Thippeswamy, S., Kiran Reddy, G. and Soujanya. (2015). Heterosis studies in sesame (*Sesamum indicum* L.). *IJPAES*, **5(3)**: 177-183.
24. Sushila, S., Kumar, S.R. and Babita kumari. (2020). Heterosis and combining ability studies in Sesame (*Sesamum indicum* L.). *J. Oilseeds Res.*, **37(1)** : 16-20.
25. Tripathy, Swapan K., Mishra, D.R., Panda, S., Senapati, N., Pramod K. Nayak, G. B. Dash, Sasmita Dash, Kartik C. Pradhan, Dash, A.P., Mishra, D., Mohanty, S. K., Mohapatra, P. M. and Lenka, D. (2016). Identification of heterotic crosses for sesame breeding using diallel matting design. *Tropical Plant Res.*, **3(2)**: 320–324.
26. Virani, M.B., Vachhani, J.H., Kachhadia, V.H., Chavadhari, R.M. and Mungala, R.A. (2017). Heterosis studies in sesame (*Sesamum indicum* L.). *Electron. J. of Plant Breed.*, **8(3)**: 1006-1012.
27. Wei., P, Zhao, F., Wang, Z., Wang, Q., Chai, X., Hou, G., Meng, Q. (2022). Sesame (*Sesamum indicum* L.): A Comprehensive Review of Nutritional Value, Phytochemical Composition, Health Benefits, Development of Food, and Industrial Applications. *Nutrients*, **14(19)** :4079. doi: 10.3390/nu14194079. <https://aicrp.icar.gov.in/sesame/about-us/>

UNDER PEER REVIEW