

# A Simplified, Efficient and Rapid DNA Extraction Protocol from Rice Grains and Leaf

## Abstracts

DNA extraction is very complicated with different plant species because the presence of secondary metabolites that hinder with DNA isolation processes and application like DNA restriction, Gene amplification, as well as gene cloning. A simple method for preparation of rice genomic DNA was developed. The current study expressed comparatively rapid, cost-effective, less time consuming, methods for DNA extraction from seed, young leaves and old leaves of rice genotypes without using liquid nitrogen, phenol, and  $\beta$ -mercaptoethanol in extraction buffer. Using this method, high quality and quantity of genomic DNA was get from 0.5 g of rice seed, young leaves and 75 days old leaves and extracted total nucleic acids (genomic DNA) were amplified by employing single sequence repeats (SSR) or microsatellite markers, which produced reproducible results. This protocol resolves the problems of DNA degradation, contamination, and low yield due to binding and/or coprecipitation with starches and polysaccharides.

**Keywords:** DNA isolation; PCR; *Oryza sativa*; Molecular Marker, DNA extraction, Molecular Breeding

## Introduction

The word “Dhan” is developed from Sanskrit word “Dhanya” which means “Sustainer” of the human race. Rice (*Oryza sativa*) belongs to the family Poaceae and sub-family *Oryzoideae*. Rice is a staple food that feeds over 50% population of the world; thus, it is essential in global food security, which contains approximately 80% carbohydrates, 7-8% protein, 3 % fats and 3 % fibers [1]. Rice is most important agronomic, monocot model crop, grown under the different eco-geographical environment in various tropical and sub-tropical countries. The genome size of rice is 430 Mb. Breeding techniques viz. Hybrid rice technology, molecular breeding, Quantitative Trait Loci (QTL) analysis etc. used for rice have attracted worldwide concern from both the public and private sectors. Molecular marker-assisted selection (MAS), has been broadly applied in rice in previous studies [2].

Molecular marker-assisted selection (MAS), Quantitative Trait Loci (QTL) analysis, evaluation of genetic diversity, conformation of transgenic and germplasm identification are important techniques in modern breeding programs. Hundreds to thousands of samples are required to be practiced for the MAS, QTL, screening of transformants as well as molecular studies for required rapid DNA isolation methods for such kind of protocols. Several authors [3, 4, 5, 6,7] explain DNA extraction protocols are mostly used in plant biotechnology and/or plant molecular biology, but available different methods are comparatively expensive, used hazardous chemicals, time consuming, involved multiple steps, require expensive enzymes and used liquid nitrogen for grinding the samples [5, 7, 8]. The Cetyl-trimethyl-ammonium bromide (CTAB) is frequently used surfactants in DNA isolation include the lyses of cell wall as well as cell membrane, which release the genetic material into extraction buffer. CTAB protocols are very famous methods for rice DNA extraction, including other plants, organisms such as bacteria [9], Fungi [10], nematods [11], and animals [12]. Several modifications have been present in CTAB protocols [6,13, 14, 15]. Some protocols have been informed to reduce the DNA isolation steps but require more amounts of plant tissue samples and liquid nitrogen [16]. The modified CTAB isolation protocols employing high concentration of salts to use removed polysaccharides because huge quantity of polysaccharide in samples the problematic at the time of PCR reaction.

A number of protocols are available and are being developed for extraction of DNA from plants. While, diverse plants contain different amounts of DNA, Isolation of high-quality nucleic acids is a prerequisite for MAS studies. Many DNA extraction protocols have been developed for rice [17]. However, these protocols typically involve the use of living organs such as green leaves as experimental materials; requiring researchers to germinate the seeds and to select the target plants at the seedling stage based on polymerase chain reaction results. It needs few weeks or few months from plantation to fresh tissue collection and also requires more attention for management practices. To overcome these problems initiate destructive detection methods by isolating nucleic acids from triploid embryos [18], develop nucleic acids isolation protocol using the dry half seeds of rice [5, 14], develop a rapid protocol for DNA extraction from seed of rice.

However, in the protocols provided by [17, 18, 19], the concentrations of nucleic acids from cereal crops (rice, wheat and maize) were relatively low. Most laboratories wish for simple and fast procedure for obtaining plant genomic DNA for PCR, and good-quality DNA for complete enzyme digestion. Therefore, a protocol for extracting genomic DNA from young or old, fresh or dry rice leaves, dry seeds, panicle, and stems that is applicable to a variety of plants,

regardless of the complexity of their genomes is presented. The objective of this study was to develop a simple and rapid method to isolate DNA under normal laboratory conditions (room temperature) from small amount of tissue for large number of samples.

## Material and Methods

### Materials

In present study we were used the Basmati rice and non-Basmati plant material of three stages viz. dry seeds, 15 days old leaf (grown in small Aluminum pots) and 65 to 70 days old leaf (grown in field conditions) for isolation of DNA. All leaves samples were collected from 15 days old plant raised in pots and 70 days old plants grown in field, wrapped in aluminum foil and kept immediately in an ice containing box.

### Primers

Isolated nucleic acids were amplified with Simple Sequence repeat (SSR) markers and specific primers (Table:1).

**Table 1. List of four primers utilized for amplification of rice genomic DNA extracted from twelve basmati and non basmati entries used in the study**

| S.N. | Primers | Chro No. | Forward                | Reverse               | Motif Repets |
|------|---------|----------|------------------------|-----------------------|--------------|
| 1.   | RM 1024 | 5        | GCATATAACCATGGGGATTGG  | GGGATTGGGATAATGGTGTG  | (AC)13       |
| 2.   | RM 1183 | 1        | GGGCACGAATAAAACCAGAG   | GGGATGGTCCAATGACAAAG  | (AG)14       |
| 3.   | RM 1018 | 4        | ATCTTGTCCTCCACTGCACCAC | TGTGACTGCTTTTCTGTGCGC | (AC)13       |
| 4.   | RM 3333 | 4        | AAGCTATCGACACCGTGACC   | GCACCTTACAATTTGGCACC  | (CT)15       |

### Reagents

The solutions used were: 1M Tris-HCl (prepared using 3.94 gTris-HCl dissolved in 20 ml deionized water & pH8.0 adjusted using 1N HCl and/or 5M NaOH solution, makeup the 30ml volume with double distilled water), and 0.5M EDTA (pH 8.0) (prepared using 4.65 g ofEDTA dissolved in 15 ml de-ionized water and 25 ml final volume makeup with de-ionoized water). The 5MNaOH and 1N HCl solution was used to adjust the pH to 8.0. The 2.0M NaCl(11.58 g NaCl added into 40 ml of de-ionized water and adjudgedthe final volume to 100 mL with de-ionized water); 3% CTAB (3 g CTABdissolved into 100 ml de-ionized water); and 70% ethanol (75 ml absolute ethanol mixed with 25 ml de-ionized water.The 3X extraction buffers was prepared by using 3 % (w/v) CTAB solutions 2 % PVP,1MTris-HCl (pH8.0), 0.5M EDTA (pH 8.0), and 2M NaCl (Table-2). The another solution chloroform: isoamyl alcohol (24:1) (CI)(Table 3), 50X Tris-acetate-EDTA (TAE) buffer (Table 4), and 1X TE buffer (Table 5) their composition and preparation are available.

## DNA extraction procedure

The young leaves were collected from 15 days old seedlings raised in the pots, wrapped in aluminium foil and kept immediately in a box containing ice. Same procedure used in 70 days old field growing rice plant and taking seeds are stored in laboratory. 0.5 to 1.0 gram fresh young and older leaf sample and 15 to 20 grains were grind in mortar and pastel in 2000  $\mu$ l 3X extraction buffer.

**Table 2: Composition (only for 100ml) and Preparation of 3X CTAB extraction Buffer solution**

| Reagent                       | Weight in gram | Stock Solution | Final Concentration | 100ml preparation |
|-------------------------------|----------------|----------------|---------------------|-------------------|
| Tris-HCl                      | 3.94           | 1.0 M          | 100 mM              | 10 ml             |
| EDTA                          | 4.65           | 0.5 M          | 20 mM               | 10 ml             |
| NaCl                          | 11.58          | 4.95 M         | 2.0 M               | 40 ml             |
| CTAB                          | 3 (w/v)        | ---            | 3 %                 | 3 g               |
| PVP                           | 2 (w/v)        | ---            | 2 %                 | 2 g               |
| <b>Double Distilled Water</b> |                | ---            | ---                 | 30 ml*            |
| <b>Final Volume =</b>         |                |                |                     | 100               |

\* Water amount more or less as required to maintain 100 ml final volume.

**Table 3: Composition and preparation of fresh Chloroform: Isoamyl alcohol (24 :1) without phenol**

| S.N.         | Reagents        | 100 ml preparation |
|--------------|-----------------|--------------------|
| 1.           | Chloroform      | 96.0 ml            |
| 2.           | Isoamyl alcohol | 4.0 ml             |
| <b>Total</b> |                 | 100.0 ml           |

**Table 4: Composition and preparation of the 50X and 1X TAE Buffer**

| S.N. | Reagent             | 1 Litre 50X Buffer | 1 litre 1 X TAE Buffer           |
|------|---------------------|--------------------|----------------------------------|
| 1.   | Tris Base (pH8.0)   | 242.4 g            | 20.0 ml TAE + 980.0 ml<br>DD H2O |
| 2.   | Glacial Acetic Acid | 57.3ml             |                                  |
| 3.   | 0.5M EDTA (pH 8.0)  | 100 ml             |                                  |

**Table 5: Composition and Preparation of the 1X TE Buffer**

| S. N. | Reagents | Stock Solution (M) | Final Concentration (mM) | 250 ml preparation |
|-------|----------|--------------------|--------------------------|--------------------|
|-------|----------|--------------------|--------------------------|--------------------|

|    |                     |      |       |        |
|----|---------------------|------|-------|--------|
| 1. | Tris-HCl (pH 8.0)   | 1.0  | 10 mM | 2.5 ml |
| 2. | EDTA (pH 8.0)       | 0.5  | 1.0   | 0.5 ml |
| 3. | dd H <sub>2</sub> O | ---- | ---   | 247 ml |

The homogenised mixture transferred to 2.0 ml eppendorf centrifuged tube and the content were mixed gently by swirling and inverting the tube and incubated at 64 °C in water bath for 50 min to 1 hour with occasional mixing at 10 to 15 min intervals. The tube was taken out and an equal volume of Chloroform: isoamylalcohol (24:1) was added. The content was mixed by inversion for 10 min and centrifuged at 10,000 rpm for 10 min. The clear aqueous layer was transferred to new tubes and re-extracted with an equal volume of CI and centrifuged at 10,000 rpm for 10 min. The supernatant was transferred into new tubes. Two third volume of isopropanol was added and mixed gently by inverting the Eppendorf tube and placed at -20 °C for 30 min (4 °C for the whole night). Genomic DNA was pelleted by centrifuging at 10,000 rpm for 10 min and the pellet was washed with 300 µl Ethanol (70%) and centrifuging at 10,000 rpm for 5 min. The DNA pelleted was air dried and dissolved in 30 µl TE buffer by gentle tapping and isolated genomic DNA was stored at -20 °C for further use.

#### **Quality Determination of isolated DNA**

The determination of quality of isolated DNA is essential to assess the suitability of good quality DNA for PCR amplification. The isolated genomic DNA samples were subjected to electrophoresis through agarose gel to assess the quality of DNA isolated from the seed, leaves of young seedlings, and 70 days old rice plant achieved from the genotypes under evaluation in the present study. The DNA sample 3 µl were mixed with 1/8<sup>th</sup> volume of 6 X DNA gel loading dye and was loaded onto 0.8 % agarose gel containing of ethidium bromide, in a submarine horizontal electrophoresis unit filled with 1X TAE buffer. Electrophoresis was carried out at 90V for 45 min. The gel was viewed under UV light to detect the quality of DNA. The DNA fragments were visualized as fluorescent bands because of fluorescence of ethidium bromide intercalated between the nitrogenous bases of DNA under UV light. Appearance of a single, sharp band of high molecular weight without smearing indicated the extraction of a good quality DNA sample with uniform segment size and less damage during extraction process.

#### **Amplification of genomic DNA and separation of amplified DNA through agarose gel electrophoresis**

The primer vials were centrifuged before and after the addition of 1X TE buffer to the vials. After dilution the concentration of each primer stock solution was obtained as 100 µM. The

diluted primers were stored at  $-20^{\circ}\text{C}$ . Amplification of DNA was carried out using a standard protocol of polymerase chain reaction (PCR) adjusted to laboratory conditions. The amplification was carried out in a thermal cycler using 15  $\mu\text{l}$  of reaction mixture (Table 6). The reaction mixture was prepared by varying the components involved in composition of reaction mixture.

**Table6: Composition of PCR reaction mixture (15  $\mu\text{l}$ )**

| Sl. No. | Components  | Volume ( $\mu\text{l}$ ) |
|---------|---|--------------------------|
| 1       | Water (Protase and Nuclease free)                   | 3.0                      |
| 2       | 5X PCR Buffer                                       | 2.8                      |
| 3       | $\text{MgCl}_2$ (10 mM)                             | 1.4                      |
| 4       | dNTPs (200 $\mu\text{M}$ )                          | 2.8                      |
| 5       | Primer F (5 $\mu\text{M}$ )                         | 1.0                      |
| 6       | Primer R (5 $\mu\text{M}$ )                         | 1.0                      |
| 7       | Taq DNA polymerase (1 unit)                         | 1.0                      |
| 8       | Template DNA (Diluted 1 : 9 (DNA:H <sub>2</sub> O)) | 2.0                      |
| Total   |   | 15.0 $\mu\text{l}$       |

The polymerase chain reaction was performed with the help of microsatellite sites based 3 pairs of forward and reverse simple sequence repeat (SSR) primers. The amplification was carried out in a thermo-cycler using the program standardized for this purpose. The amplified product was separated using 2.0 % agarose gel with ethidium bromide as a staining reagent. The amplified product with an expected product size above 100 bp was subjected to 2% gel electrophoresis at 90 V for one hour and then visualized and documented in gel documentation system. The size of the amplified product was estimated with the help of a 100 bp ladder.


### Result and Discussion

The superior quality of DNA sample was obtained from seed, 15 days young leaves, and 70 days old leaves, of twelve rice (*Oryza sativa*) genotypes and amplified using four simple sequence repeat (SSR) or microsatellite markers namely viz. RM 1024, RM 1183, RM 1018, and RM 3333. The amplified product was observed in all samples tested with little variation in intensity of the amplified bands. After amplification RM 1024 produced 110-142 bp, RM 1183 produced 113-142 bp, RM 1018 produced 135-149 bp, and RM 3333 produced 160-189 bp polymorphic product and allelic size difference 32, 29, 14 and 29 bp respectively (Figure 1 and 2). Here is described a simple and reproducible procedure for PCR amplification of rice DNA, that also is applicable for other plant genomes.

DNA samples prepared using the extraction procedures (lane 1 = Marker (M), 2-5 Seed genomic DNA, 6-9 young leaves Genomic DNA and 10-13 mature 70 days old genomic DNA in Figure 1) were subjected to PCR amplification using different primers: RM 1024 on chromosome


5, RM 1183 on chromosome 1, RM 1018 on chromosome 4, and RM 3333 on chromosome 4, and the repeat motif of SSR markers is (AC)<sub>13</sub> (AG)<sub>14</sub>, (AC)<sub>13</sub> and (CT)<sub>15</sub> respectively. All the genomic DNA samples produced a clear, sharp, and reproducible PCR product when the primers were used for PCR amplification (Figure 2). Although two variations of the DNA extraction procedure were used, there was no difference between lanes of two protocols.

Numerous protocols for plant genomic DNA extraction are available however, the better results were obtained with that utilized CTAB [1, 5, 7, 16, 19, 20]. In this experiment is anxious with modification of CTAB method to isolate DNA without using liquid nitrogen, β-mercaptoethanol, and phenol. The concentration of NaCl was high (2M) which extracted high quantity of DNA. The main action of NaCl in extraction buffer is the removal of proteins and carbohydrates which are attached to the DNA. Polyvinyl pyrrolidone forms H-Bonds with phenolic metabolites and prevent the plant genomic DNA contamination by phenolic metabolites [13, 16, 17, 18]. There are several advantages in using genomic DNA extraction protocol suitable for polymerase chain reaction (PCR) genotyping, which can be employed for the recognition of alleles in diverse genetic, breeding and biotechnological approaches, viz. marker-assisted selection (MAS), quantitative trait loci (QTL), genetic mapping, and Transgenic Screening, mutant introgression. All four microsatellite markers show very good amplification and polymorphism with the DNA extracted through this method. Therefore, this method should be recognized as a good rapid and inexpensive method for DNA extraction from rice leaves and seeds.



|               |                       |                     |                   |                      |                        |                     |
|---------------|-----------------------|---------------------|-------------------|----------------------|------------------------|---------------------|
| <b>M-</b>     | 1. Nagina-22          | 2. Punjab Basmati-4 | 3. Kasturi        | 4. Pusa Basmati-1121 | 5. Ranvir Basmati      | 6. CSR-30           |
| <b>Marker</b> | 7. Vallabh Basmati-23 | 8. IR-64            | 9. Pant Basmati-1 | 10. Sarbati          | 11. Vallabh Basmati 24 | 12. Pusa Basmati -1 |

**Figure 1: Crude total genomic DNA isolated from twelve different varieties 1 to 4 isolated from seeds, 5 to 8 isolated young leaves and 9 to 12 isolated 70 days older rice genotypes.**



| M-<br>Marker | 1. Nagina-22             | 2. Punjab<br>Basmati-4 | 3. Kasturi        | 4. Pusa Basmati-1121 | 5. Ranvir<br>Basmati      | 6. CSR-30           |
|--------------|--------------------------|------------------------|-------------------|----------------------|---------------------------|---------------------|
|              | 7. Vallabh<br>Basmati-23 | 8. IR-64               | 9. Pant Basmati-1 | 10. Sarbati          | 11. Vallabh<br>Basmati 24 | 12. Pusa Basmati -1 |

**Figure 2: Amplification of genomic DNA with selected four different primers with all genotypes.**

### Conclusions

In summary, the procedure for rice genomic DNA or nucleic acids extraction described here could be a trusted and reliable protocol to work well for analysis without using liquid nitrogen, 2-mercaptoethanol, and Phenols.

### REFERENCES

1. Kumar, P. (2022). *Expression Analysis of Transcription Factors and Related Genes Induced by Water Stress in Contrasting Genotypes of Rice (Oryza Sativa L)*. S. V. P. U. A. & T., Meerut. 1-7.
2. Jena, K. K., and Mackill, D. J. (2008). Molecular markers and their use in marker- assisted selection in rice. *Crop science*, 48(4): 1266-1276.
3. Murray, M. G., and Thompson, W. F. (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic acids Res.* 8: 4321-4325.
4. Doyle, J. J. and Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19: 11-15.

5. Ahmadikhah, A. (2009). Rapid mini-prep DNA extraction method in rice (*Oryza sativa*). *African journal of Biotechnology*. 8(2).
6. Ferdous, J., Hanafi, M. M., Rafii, M. Y., and Muhammad, K. (2012). A quick DNA extraction protocol: Without liquid nitrogen in ambient temperature. *African Journal of Biotechnology*. 11(27): 6956-6964.
7. Sajib, A. A., Bhuiya, M. A. I., and Huque, R. (2017). A simple, efficient and rapid method for good quality DNA extraction from rice grains. *Rice Science*. 24(2): 119-122.
8. Sharma R., John, S. J., Damgaard, D. M., and McAllister, T. A. (2003). Extraction of PCR-quality plant and microbial DNA from total rumen contents. *BioTechniques*34: 92-97. 96-97.
9. Caccavo, F., Lonergan, D. J., Lovely, D. R., Davis, M., Stolz, J. F., and McInerney, M. J. (1994). *Geobactersulfurreducens* sp. nov., a hydrogen-oxidizing and acetate-oxidizing dissimilatory metal-reducing microorganism. *Appl. Environ. Microbiol.* 60: 3752-3759.
10. Thuan, N. T. N., Bigirimana, J., Roumen, E., Straeten, D. V. D., and Hofte, M. (2006). Molecular and pathotype analysis of the rice blast fungus in North Vietnam. *Eur. J. Plant Pathol.* 114: 381-396.
11. Mondino, E. A., Covacevich, F., Studdert, G. A., Pimentel, J. P., and Berbara, R. L. (2015). Extracting DNA of nematodes communities from Argentine Pampas agricultural soils. *Anais da Academia Brasileira de Ciências*, 87: 691-697.
12. Shahjahan, R. M., Hughes, K. J., Leopold, R. A. and DeVault, J. D. (1995). Lower incubation temperature increases yield of insect genomic DNA isolated by the CTAB method. *Biotechniques*, 19: 332-334.
13. Punya, Sharma, V. K., and Kumar, P. (2017). A modified method for maize genomic DNA extraction without liquid nitrogen and phenol. *Indian Res J Genet & Biotech.* 9: 45-51.
14. Kang, H. W., Cho, Y. G., Yoon, U. H., and Eun, M. Y. (1998). A rapid DNA extraction method for RFLP and PCR analysis from a single dry seed. *Plant Mol. Biol. Rep.* 16: 90.
15. Allen, G. C., Vergara, M. A. F., Krasnyanski, S., Kumar, S., and Thompson, W. F. (2006). A modified protocol for rapid DNA isolation from plant tissue using cetyltrimethylammonium bromide. *Nature protocols* 1: 2320-2325.
16. Tussell, R. T., Ramayo, A. Q., Herrera, R. R., Saavedra, A. L. Brito, D. P. (2005). A fast simple and reliable high yielding method for DNA extraction from different plant species. *Mol. Biol.* 31: 137-139.
17. Clark, K. A., and Krysan, P. J. (2007). Protocol: an improved high-throughput method for generating tissue samples in 96-well format for plant genotyping (Ice-Cap 2.0). *Plant methods*, 3(1), 1-8.
18. Chunwongse, J., Martin, G. B. and Tanksley, S. D. (1993). Pre-germination genotypic screening using PCR amplification of half-seeds. *Theor. Appl. Genet.* 86: 694-698.
19. Kang, T. J., and Yang, M. S. (2004). Rapid and reliable extraction of genomic DNA from various wild-type and transgenic plants. *BMC Biotech.* 4: 20.

20. Suman, S. K., Kumar, M., Sharma, V. K., and Kumar, R. (2020). A modified and efficient CTAB genomic DNA extraction method from maize leaf for PCR analysis. *Journal of Pharmacognosy and Phytochemistry*, 9(4): 3418-3420.

UNDER PEER REVIEW