

# **Molecular Characterization of *Cryptosporidium* Oocysts in Under-five Children with Diarrhoea in Ile-Ife, Nigeria**

## **Abstract**

**Aim:** Cryptosporidiosis is the fifth leading cause of diarrhoea-related death globally among under-five children. This study was conducted to identify and characterize oocysts of *Cryptosporidium* in under-five children with diarrhoea in Ile-Ife.

**Study design:** It was a case-control study.

**Place and duration of the study:** Department of Medical Microbiology and Parasitology, Obafemi Awolowo University between April, and October 2019.

**Methodology:** Ethical approval was obtained for this study. Consent was also obtained from the parents of the children. Stool samples from 53 children with diarrhoea (cases) and 94 children without diarrhoea (controls) under five years of age were collected. A structured proforma was used to obtain relevant information from the parents of the children. Stool samples were collected and examined macroscopically for consistency, appearance, and colour. Modified Ziehl-Neelsen staining was carried out on all the stool samples. The samples positive for *Cryptosporidium* oocysts by staining were genotyped by nested polymerase chain reaction and confirmed by sequencing of their 18S rRNA genes. Data generated were analyzed using descriptive and inferential statistics with SPSS software version 20 and STATA 15.

**Results:** The prevalence of cryptosporidiosis among the study participants was 23.1% with rates of 43.4% (23) and 11.7% (11) for diarrhoeic and non-diarrhoeic children respectively. Of the 34 stool samples analysed, 6 (17.6%) were amplified by nested PCR and 4 (11.8%) were identified by sequencing. The species of *Cryptosporidium* identified were *Cryptosporidium parvum* (75%) and *Cryptosporidium bovis* (25%).

**Conclusion:** In conclusion, Cryptosporidiosis is prevalent among under-five children with *Cryptosporidium parvum* and *Cryptosporidium bovis* as the infecting species.

**Key words-** Cryptosporidiosis, diarrhoea, under-five children, *Cryptosporidium*, Infection.

## **INTRODUCTION**

The protozoal parasite *Cryptosporidium* is an important cause of diarrhoeal disease in humans and animals globally. There are presently over 40 recognized species with most of these species found in humans (Innes *et al.*, 2020). *Cryptosporidium* causes Cryptosporidiosis which is a significant diarrhoeal disease that affects young children especially those under the age of 5 years and individuals that are immunocompromised (Innes *et al.*, 2020; Semani *et al.*, 2023). Like many other important protozoan parasites such as *Giardia*, *Cryptosporidium* is spread through the faecal-oral route. Infection can be acquired through ingestion of oocyst infected water, food and or a direct contact with an infected individual (Shirley *et al.*, 2012; Checkley *et al.*, 2015; Johansen *et al.*, 2015). It is also transmitted zoonotically as *Cryptosporidium* is known to have a wide host range. It was first recognized as a human pathogen in 1976 (Ryan *et al.*, 2021) and since then its status has sailed up from a mere curiosity to a pathogen of worldwide importance. *Cryptosporidium* has also been reported in animals and the environment with various species and genotypes identified from surface water and animals (Mensah *et al.*, 2023; Mugasa *et al.*, 2023; Sharafi *et al.*, 2023). *Cryptosporidium* has recently been established as the second most important infectious agent after rotavirus, and the single most important parasitic cause of gastroenteritis (Kotloff *et al.*, 2013).

Studies have also established a link between *Cryptosporidium* infection and growth impairment, physical fitness, and cognitive function of children. (Kotloff *et al.*, 2013; Korpe *et al.*, 2016; Khalil *et al.*, 2018; Rogawski *et al.*, 2018). Approximately 7.6 million diarrhoea cases annually are attributable to *Cryptosporidium*, with about 2.9 million in Sub-Saharan Africa (Sow *et al.*, 2016 and Mutai *et al.* 2020). Recent studies have also linked cryptosporidiosis to colorectal cancer (Innes *et al.*, 2020). Molecular tools have revolutionized identification of microorganisms and of the 44 recognised *Cryptosporidium* species and greater than 120 genotypes, 19 species, and four genotypes have been reported in humans with *Cryptosporidium hominis*, *Cryptosporidium parvum*, *Cryptosporidium meleagridis*, *Cryptosporidium canis* and *Cryptosporidium felis* being the most prevalent (Ryan *et al.*, 2021). However, there is paucity of information on the species infecting children in Ile-Ife environment. It was against this background this study sought to assess the prevalence and species of *Cryptosporidium* in under-five children.

## MATERIAL AND METHODS

**Study design** - It was a case control study.

**Study area** - The study area was Ile-Ife, Osun State, Nigeria. Ile-Ife lies between Latitudes 7° 31' N and 7° 35' N and Longitudes 4° 30' E and 4° 35' E. The research study was conducted in the Department of Medical Microbiology and Parasitology, Obafemi Awolowo University, Ile-Ife, Osun state, Nigeria.

**Study population** - The participants (under-five children) were recruited from Enuwa Primary Health Center, Oke-ogbo State Hospital and Obafemi Awolowo University Teaching Hospitals Complex (OAUTHC), Ile-Ife, Osun state.

**Sample size and selection of subjects**- A minimum sample size of 147 was calculated for this study comprising 94 controls (children under-five without diarrhoea) and 53 cases (children under-five with acute diarrhoea). The proportion ratio used was 2:1 (controls to cases). The children were randomly selected from these centers.

**Demographic information**- A structured proforma was used to obtain relevant information such as age, sex, weight, height, history of diarrhoea, source of drinking water and contact with pets.

**Sample collection, processing, and storage** - The stool samples were collected into clean grease free universal bottles. Each sample was divided into 2 portions, a portion was preserved with 10% formalin while the other part was preserved with 2.5% Potassium dichromate. Stool samples preserved with 2.5% potassium dichromate were stored at 4°C for molecular analysis, while those preserved in formalin were used for formol-ether concentration and microscopy as recommended (Lemos *et al.*, 2011).

The samples were processed by carrying out macroscopy, direct wet mount examination, formal-ether sedimentation technique, Modified Ziehl-Neelsen staining and molecular analysis. The stained slides were observed using immersion oil objective lens and the size of the oocysts was confirmed using an ocular micrometer. Samples that were positive for *Cryptosporidium* oocysts were further analysed using nested polymerase chain reaction.

## **Molecular analysis**

### **DNA Extraction**

The positive faecal samples that were preserved with 2.5% of potassium dichromate at 4°C were diluted with nuclease-free water at 1:1 proportion. To each of the fecal sample 500µl lysis buffer [50µl Tris-HCl 100 mM (pH=8.0); 50µl ethylenediaminetetraacetic acid (EDTA) 50 mM (pH=8.0); 50 µl 10% sodium dodecyl sulfate (SDS); 3.5µl 70 mM β-mercaptoethanol; 1µl 1% polyvinylpyrrolidone (PVP) and 345.5µl sterile distilled water] were added and incubated at 65°C in a water bath for 90 minutes. Twenty micro-liters of proteinase K (20 mg/ml) was applied after incubation. The samples were incubated for 15 minutes in an ice bath. The tubes were shaken to mix well, and the mixture was centrifuged at 12000 rpm for 5 minutes. After centrifugation, the aqueous phase which is the supernatant was gently removed to a new eppendorf tube, 500µl chloroform was added; homogenized and centrifuged at 12000 rpm for 15 minutes. The supernatant seen after centrifuging was put into another eppendorf tube and precipitated with 500µl ice-cold isopropyl alcohol and held for 10 minutes in an ice bath. After 10 minutes, it was centrifuged at 12000 rpm for 3 minutes. The supernatant was gently discarded, and the precipitated DNA was washed with 500µl of 70% ethanol and centrifuged at 12000 rpm for 3 minutes. The supernatant was discarded, and the DNA was dried.

### **18S rRNA Nested PCR Assay**

The primary mix containing the extracted DNA was amplified using the PCR thermal cycler (GeneAmp® PCR System 9700). The cycling condition were selected as follows: one cycle of denaturation at 94°C for 5 minutes; 45 cycles of 94°C for 30 seconds (denaturation), 48°C for 20 seconds (annealing), 72°C for 30 seconds (extension); then a final extension of 72°C for 7 minutes and held at 4°C until removed from the thermo-cycler.

The secondary mix containing the primary PCR product was amplified using the same thermo-cycler. The cycling condition were selected as followed: one cycle of denaturation at 94°C for 5 minutes; 45 cycles of 94°C for 30 seconds (denaturation), 50°C for 20 seconds (annealing), 72°C for 30 seconds (extension); then a final extension of 72°C for 7 minutes and held at 4°C until removed from the thermo-cycler. Sequencing of 18S rRNA PCR products was carried out using the Nimagen, Brilliant Dye™ Terminator Cycle Sequencing Kit V3.1, BRD3-100/1000 according to the manufacturer's instructions. The labelled products were then cleaned with the ZR-96 DNA Sequencing Clean-up Kit (Catalogue No. D4053). The cleaned products were injected on the Applied Biosystems ABI 3500XL Genetic Analyser with a 50 cm array, using POP7. The sequence chromatogram analysis was performed using

FinchTV analysis software and was changed to FASTA format before blasting using NCBI database. Table 1 shows primers used for the nested PCR while Tables 2 and 3 show the primary and secondary PCR work sheets.

**Table 1. Primers used for nested PCR.**

Primer name	Sequence 5'-3'	Target gene	Amplicon size (bp)
Primer 1-R	CACCAGACTTGCCCTCCAAT	18S rRNA	215
Primer 1-F	CAATGACGGGTAACGGGGAA	18S rRNA	215
Primer 2-R	GCCTGCTGCCTTCCTTAGAT	18S rRNA	78
Primer 2-F	CGGGTAACGGGGAATTAGGG	18S rRNA	78

**Table 2. Primary PCR work sheet**

Primary PCR Mix	n = 1	n = 34
Master mix	12.5 µl	687.5 µl
Primer 1-F	0.5 µl	27.5 µl
Primer 1-R	0.5 µl	27.5 µl
Extracted DNA working solution	4 µl	220 µl
Water	7.5 µl	412.5 µl
Total	25 µl	1375 µl

**Table 3. Secondary PCR work sheet**

Secondary PCR Mix	n = 1	n = 6
Master mix	12.5 µl	75 µl
Primer 2-F	0.5 µl	3 µl
Primer 2-R	0.5 µl	3 µl

---

Primary PCR Product	3 $\mu$ l	18 $\mu$ l
Water	8.5 $\mu$ l	51 $\mu$ l
Total	25 $\mu$ l	150 $\mu$ l

---

**Data Analysis** – Data were analyzed using frequency, proportion, percentages, tables, and Pearson’s chi-square was used to determine the association between Cryptosporidiosis and associated symptoms.

## **Results**

### **Socio-Demographic Characteristics of the Respondents**

Out of the 147 children recruited, 38.8 % were infants while 61.2. % were more than 12 months in age. Also, 51 % were male while 49.0 % were female. Diarrhoea occurred in 36.1% of the study participants while 63.9% did not have diarrhoea. Out of those with diarrhoea, 34.0 % were exposed to all kinds of domestic animals while 66.0% were not

exposed. For the children without diarrhoea, 59.6% were exposed to domestic animals while 40.4% were not. Majority of the respondents were Christians 84.4%; Muslims were 15% and other religions 0.6 %. For the father's occupation, 27.9 % were artisans and 27.2 % traders while for the mother's occupation, 27.2 % were artisans and 36.1 % were traders. Majority of the guardians of the children had tertiary education (46.3%), 43.5% had secondary education, 8.8% had primary education and 1.4 % had no education. Forty-four-point two percent of the children have not started school at all, and majority of the children (89.8%) had their mother as their informant.

Table 4 shows the prevalence of *Cryptosporidium* from the stool samples of the study participants by the Modified Ziehl-Neelsen technique.

**Table 4: Prevalence of *Cryptosporidium* from the stool samples of under-five children analysed by Modified Ziehl- Nelseen staining technique.**

Stool Consistency	<i>Cryptospori dium</i> Positive	<i>Cryptospori dium</i> Negative	$\chi^2$	<i>P-value</i>

---

Diarrhoeic (n= 53)	23 (43.4)	30 (56.6)	19.1479	<b>&lt;0.001</b>
Non-diarrhoeic (n= 94)	11 (11.7)	83 (88.3)		
<b>Total (n= 147)</b>	<b>34 (23.1)</b>	<b>113 (76.9)</b>		

---

**Identification of *Cryptosporidium* from the stool Samples of Under-five Children analysed by nested PCR.**

Tables 5 and 6 shows the analysis done by PCR for further identification of *Cryptosporidium* targeting 18S rRNA gene. Of the 34 stool samples analysed by nested PCR, 6 (17.6%) were amplified and recorded as positive for *Cryptosporidium* infection by possession of 18S rRNA gene. Plates 1 and 2 are amplicons of primary and secondary PCR.


**Table 5: Identification of *Cryptosporidium* from the stool Samples of Under-five Children analysed by PCR.**

Variables	Frequency	Percentage (%)
<b>PCR</b>		
Negative	28	82.4
Positive	6	17.6
<b>Total</b>	<b>34</b>	<b>100</b>

**Table 6: Prevalence of *Cryptosporidium* from the stool Samples of Under-Five Children analysed by PCR.**


Variables	<i>Cryptosporidium</i> positive children		$\chi^2$	<i>P-value</i>
PCR	Diarrhoea n=23	Non-Diarrhoea n=11	0.8191	0.365
Negative	18(78.26)	10(90.91)		
Positive	5(21.74)	1(9.09)		
<b>Total</b>	<b>23(100.00)</b>	<b>11(100.00)</b>		

L 1 2 3 4 5 6 L



**Plate 1: Amplicons of primary PCR of 18S rRNA gene.**  
(Lane L: DNA ladder (50 bp), Lane 2, 4, 5, and 6 = 215 bp)

**L    1    2    3    4    5    6    L**



**Plate 2: Amplicons of secondary PCR of 18S rRNA gene.**  
(Lane L: DNA ladder (50 bp), Lane 2, 3, 4, 5 and 6 = 75 bp)

**Characterization of *Cryptosporidium* species from the stool Samples of Under-five Children analysed by sequencing of 18S rRNA PCR products.**

Table 7 shows the identified *Cryptosporidium* species analysed by sequencing of the 18S rRNA gene. Of the 6 *Cryptosporidium* isolates that were characterised by sequencing analysis, 3 were identified as *Cryptosporidium parvum* and 1 was identified as *Cryptosporidium bovis*. The 4 isolates were from children with diarrhoea while the other 2 were not identified.

UNDER PEER REVIEW

**Table 7: *Cryptosporidium* species identified by sequencing analysis of 18S rRNA PCR products.**

<b>Species Identified</b>	<b>Number of Species Identified (%)</b>	<b>Accession Number</b>	<b>Partial Sequence</b>
<i>Cryptosporidium parvum</i>	3(75%)	FJ039877.1	GGAGCCTGAGAACGGCTACCACA TCTAAGGAAGGCAGCAGGC
<i>Cryptosporidium bovis</i>	1(25%)	KY809006.1	GCAGCCTGAGAAACGGCTACCAC ATCTAAGGAAGGCAGCAGGC

UNDER PEER REVIEW

## Discussion

The prevalence of *Cryptosporidium* among the under-five children investigated in this study by Modified Ziehl-Neelsen (MZN) staining technique is 23.1%. This rate is high, and this finding substantiates *Cryptosporidium* as an important cause of diarrhoea in children. Wellington *et al.* (2008) and Musa *et al.* (2014) reported prevalence rates of 17.1% and 17.8% in Lagos and Zaria respectively which are lower than the observed prevalence in this study. However, other studies such as Nassar *et al.* 2017, Shinkafi and Muhammed, 2017 reported higher prevalence rates of 38.3% in Osun and 62.5% in Sokoto states respectively. The varying prevalence rates could be due to differences in geographical location and possibly levels of hygiene. This study also revealed that diarrhoea was significantly associated with presence of *Cryptosporidium* in the stool samples of the children which is similar to the reports of Inyang-Etoh *et al.* 2007, Dabas *et al.* 2017 and Tamomh *et al.* 2021.

The *Cryptosporidium* oocysts detected by MZN staining were further characterized by nested PCR. Out of the oocysts detected in diarrhoeic and non-diarrhoeic stools, 21.7% and 9.1% were positive for *Cryptosporidium* respectively in 6 samples due to the presence of 18S rRNA gene. Three were further identified as *Cryptosporidium parvum* (75%) and 1 (25%) was identified as *Cryptosporidium bovis* by sequence analysis. *Cryptosporidium* species that are commonly implicated in human cryptosporidiosis are *Cryptosporidium hominis* and *Cryptosporidium parvum* (Lebbad *et al.*, 2021; Krumkamp *et al.*, 2021). *Cryptosporidium felis*, *Cryptosporidium canis* and *Cryptosporidium meleagridis* also cause infections in humans especially those that are immunocompromised (Ryan *et al.* 2021). Of these species of *Cryptosporidium*, the two that commonly cause human infection are *Cryptosporidium parvum* and *Cryptosporidium hominis*, but in addition, *Cryptosporidium parvum* infects domestic or wild ruminants. (Anejo-okopi *et al.*, 2016).

The finding of *Cryptosporidium parvum* in this study is much higher than the reports of Anejo-okopi *et al.*, 2016 and Okojokwu<sup>a</sup> *et al.*, (2016) where prevalence rates of 25% and 54% for were found respectively.

*Cryptosporidium bovis* was the other species detected in this study and it is known to commonly infect cattle in addition to three other species of *Cryptosporidium* (Thomson *et al.* 2017; Abdelaziz *et al.*, 2022). In Colombia, Higuera *et al.*, 2020 reported *Cryptosporidium bovis* as a cause of human infection. The zoonotic potential of *Cryptosporidium* was also

investigated among children less than 10 years and livestock in Egypt and *Cryptosporidium bovis* was found in 4.1% of ruminants; Combination of *Cryptosporidium parvum* plus *Cryptosporidium bovis* (5.3%) and combination of *Cryptosporidium parvum* plus *Cryptosporidium bovis* 1.2% in the children. (Helmy *et al.* 2013). Some other studies have also reported *Cryptosporidium bovis* as a cause of human infection. (Ng *et al.*, 2012 and Khan *et al.*, 2010). Cattle is the major host for *Cryptosporidium bovis* and a known reservoir of zoonotic infections (Ryan *et al.* 2014).

Human infection with *Cryptosporidium bovis* has not been previously reported in Nigeria and its finding in this study suggests that zoonotic transmission of *Cryptosporidium* in Nigeria is not limited to *Cryptosporidium parvum*. However, there have been reports of *Cryptosporidium bovis* infection among cattle in Nigeria (Okojoku<sup>b</sup> *et al.* 2016 and Chukwu *et al.*, 2019)). The presence of the species identified in this study further elucidates the dynamics of transmission of *Cryptosporidium* which could be both anthroponotic and zoonotic.

**Conclusion:** Cryptosporidiosis is highly prevalent in under five children with *Cryptosporidium parvum* and *Cryptosporidium bovis* as the infecting species. This study also highlights that the paediatric population are at risk for zoonotic transmission of *Cryptosporidium*. This study also corroborates anthroponotic and zoonotic modes of transmission in the study environment and further highlights the need for one health approach to the management of Cryptosporidiosis. Routine screening of stool samples from children for *Cryptosporidium* in addition to other intestinal pathogens should be encouraged in laboratories. Further studies are needed on children and domestic animals in Nigeria to better understand the sources of infection, transmission dynamics and genetic diversity of *Cryptosporidium*.

**Ethical approval and Consent** – Approval for the study with ethical clearance certificate was obtained from the Ethics and Research committee of OAUTHC, Ile-Ife with study protocol number (ERC/2019/04/15). The mothers of the children were informed about the research and gave their consent. Confidentiality of all patients and data was strictly maintained.

## REFERENCES

1. Innes EA, Chalmers RM., Wells B, and Pawlowic MC. A One Health Approach to Tackle Cryptosporidiosis. *Trends in Parasitology*. 2020; 36(3): 290–303.
2. Semmani M, Costa D, Achour N, Cherchar M, Ziane H, Mouhajir A, et al. Occurrence and Molecular Characterization of *Cryptosporidium* Infection in HIV/Aids Patients in Algeria. *Viruses*. 2023; 15: 362. <https://doi.org/10.3390/v15020362>.
3. Shirley DA, Moonah, SN, and Kotloff KL. Burden of disease from cryptosporidiosis. *Current opinion in infectious diseases*. 2012; 25(5): 555–563. <https://doi.org/10.1097/QCO.0b013e328357e569>.
4. Checkley W, White AC, Jaganath D, Arrowood MJ, Chalmers RM, et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for *Cryptosporidium*. *The Lancet Infectious Diseases*. 2015; 15(1): 85-94.
5. Johansen OH, Hanevik K, Thrana F, Carlson A and Stachurska-Hagen T. Symptomatic and asymptomatic secondary transmission of *Cryptosporidium parvum* following two related outbreaks in schoolchildren. *Epidemiology and infection*. 2015; 143(8): 1702–1709.
6. Ryan UM, Feng Y, Fayer R and Xiao L. Taxonomy and molecular epidemiology of *Cryptosporidium* and *Giardia* – a 50-year perspective (1971–2021) *International Journal for Parasitology*. 2021; 51: 1099–1119.
7. Mensah GT, Ayeh-Kumi PF, Annang AK, Owusu-Frimpong I, Niampoma S and Brown CA. Molecular epidemiology of *Cryptosporidium species* in Kpong and its environs, Ghana. *PLoS ONE*. 2023; 18(2): e0281216. <https://doi.org/10.1371/journal.pone.0281216>

8. Mugasa CM, Mirembe BB, Ochwo S, Nkamwesiga J, Ndekezi C, Tusabe T, Musoba A, Kankya C. Molecular Detection of *Cryptosporidium* Species in Wildlife and Humans at the Wildlife-Human Interface around Queen Elizabeth National Park, Uganda. *Parasitologia*. 2023; 3(2):181-193. <https://doi.org/10.3390/parasitologia3020019>
9. Sharafi R, Aghapour AA, and Aminpour A. Prevalence and molecular analysis of *Cryptosporidium* spp. collected from surface water. *Environ Monit Assess*. 2023; 195: 499. <https://doi.org/10.1007/s10661-023-11097-w>
10. Kotloff KL, Nataro JP, and Blackwelder WC. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicentre Study, GEMS): a prospective, case-control study. *The Lancet*. 2013;382: 209-222.
11. Korpe PS, Valencia C, Haque R, Mahfuz M, McGrath M, et al. Epidemiology and risk factors for cryptosporidiosis in children from 8 low-income sites: results from the MAL-ED study. *Clinical Infectious Diseases*. 2018; 67(11): 1660-1669.
12. Khalil IA, Troeger C, Rao PC, Blacker BF, Brown A, et al. Morbidity, mortality and long-term consequences associated with diarrhoea from *Cryptosporidium* infection in children younger than 5 years: a meta-analysis study. *Lancet Global Health*. 2018; 6: e758-768.
13. Rogawski SM, Liu ET, Platts-Mills JA, Kabir F, Lertsethtakarn P, et al. Use of quantitative molecular diagnostic methods to investigate the effect of enteropathogen infections on linear growth in children in low-resource settings: longitudinal analysis of results from the MAL-ED cohort study. *The Lancet Global Health*. 2018; 6(12): e1319-e1328.
14. Lemos FO, Almosny NP, Soares AB, and Alencar NX. *Cryptosporidium* species screening using Kinyoun technique in domestic cats with diarrhoea. *Journal of Feline Medicine and Surgery*. 2011;14(2): 113-117

15. Sow SO, Muhsen K, Nasrin D, Blackwelder WC, Wu Y, et al. The Burden of *Cryptosporidium* Diarrheal Disease among Children < 24 Months of Age in Moderate/High Mortality Regions of Sub-Saharan Africa and South Asia, Utilizing Data from the Global Enteric Multicenter Study (GEMS). PLoS neglected tropical diseases. 2016; 10(5): e0004729. <https://doi.org/10.1371/journal.pntd.0004729> .
16. Mutai D, Owili P, and Muga M. Trend of *Cryptosporidium* Infection among Children below 24 Months in an Informal Urban Settlement, Kenya. Open Journal of Medical Microbiology. 2020; 10:153-161.
17. Wellington O, Chika O, and Adetayo F. *Cryptosporidium* and other Intestinal Protozoans in Children with Diarrhoea in Lagos, Nigeria. The Internet Journal of Tropical Medicine. 2008; 5(2):1–5.
18. Musa S, Yakubu AM, and Olayinka AT. Prevalence of cryptosporidiosis in diarrhoeal stools of children under-five years seen in Ahmadu Bello University Teaching Hospital Zaria, Nigeria. Nigerian Journal of Paediatrics. 2014; 41:3
19. Shinakfi SA, and Muhammed Z. Prevalence of *Cryptosporidium* oocysts among primary school children in Wamakko local government of Sokoto State, Nigeria. Nigerian Journal of Paediatrics. 2017; 25: 1.
20. Inyang-Etoh PC, Etim NG, Useh MF, Udiong CEJ, and Essien AW. Cryptosporidiosis and Infantile Diarrhoea in Calabar, Nigeria. Journal of Medical Sciences. 2007; 7:1325 – 1329.
21. Dabas A, Shah D, Bhatnagar S, and Lodha R. Epidemiology of *Cryptosporidium* in Pediatric Diarrheal Illnesses. Indian Pediatr. 2017;54: 299-309.
22. Tamomh AG, Agena AM, Elamin E, Suliman MA, Elmadani M, et al. Prevalence of cryptosporidiosis among children with diarrhoea under five years admitted to Kosti teaching hospital, Kosti City, Sudan. BMC infectious diseases.2021; 21(1): 349. <https://doi.org/10.1186/s12879-021-06047-1>.
23. Lebbad, M., Winiacka-Krusnell, J., Stensvold, C. R., and Beser, J. (2021). High Diversity of *Cryptosporidium* Species and Subtypes Identified in Cryptosporidiosis Acquired in Sweden and Abroad. Pathogens (Basel, Switzerland). 2021; 10(5): 523. <https://doi.org/10.3390/pathogens10050523>
24. Krumkamp R, Aldrich C, Maiga-Ascofare O, Mbwana J, Rakotozandrindrainy N, Borrmann S, et al. Transmission of *Cryptosporidium* Species Among Human and Animal Local Contact Networks in Sub-Saharan Africa: A Multicountry Study. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America. 2021;72(8):1358–1366. <https://doi.org/10.1093/cid/ciaa223>

25. Anejo-Okopi, JA, Okojokwu JO, Ebonyi AO, Ejeliogu EU, Isa SE, Audu O, et al. Molecular characterization of *Cryptosporidium* in children aged 0-5 years with diarrhea in Jos, Nigeria. The Pan African Medical Journal. 2016; 25:253. [doi: [10.11604/pamj.2016.25.253.10018](https://doi.org/10.11604/pamj.2016.25.253.10018)].
26. Okojokwu OJ, Ileigo IH, Ezemuel YS, Oluyinka OO, Akpakpan EE, Kolawole T, et al. Molecular characterisation of *Cryptosporidium* species among patients presenting with diarrhoea in some parts of Kaduna State, Nigeria. American Journal of research communication. 2016; 4:87-106.
27. Thomson S, Hamilton CA, Hope JC, Katzer F, Mabott NA, Morrison LJ, et al. Bovine cryptosporidiosis: impact, host-parasite interaction, and control strategies. Vet Res. 2017; 48:42 <https://doi.org/10.1186/s13567-017-0447-0>.
28. Abdelaziz AR, Tahoun A, El-Sharkawy H, Abd El-Salam MM, Alorabi M, El-Shehawi AM et al. Overview on *Cryptosporidium bovis* and Its Effect on Calves in Some Governorates in Egypt. Journal of tropical medicine. 2022; 4271063. <https://doi.org/10.1155/2022/4271063>
29. Higuera A, Villamizar X, Herrera G, Giraldo JC, Vasquez-A LR, Urbano P, et al.. Molecular detection and genotyping of intestinal protozoa from different biogeographical regions of Colombia. PeerJ. 2020; 8: e8554 <https://doi.org/10.7717/peerj.8554>
30. Helmy YA, Krücken J, Nöckler K, Samson-Himmelstjerna G, Zessin K. Molecular epidemiology of *Cryptosporidium* in livestock animals and humans in the Ismailia province of Egypt. Veterinary Parasitology. 2013; 193: 15–24.
31. Ng J, Eastwood K, Walker B, Durreihm DN, Massey PD, Porigneaux P, et al. Evidence of *Cryptosporidium* transmission between cattle and humans in Northern New South Wales. Experimental Parasitology. 2012;130(4): 437-441.
32. Khan SM, Debnath C, Pramanik AK, Xiao L, Nozaki T and Ganguly S. Molecular characterization and assessment of zoonotic transmission of *Cryptosporidium* from dairy cattle in West Bengal, India. Veterinary Parasitology. 2010; 171: 41–47.
33. Ryan, U., Fayer, R., and Xiao, L. *Cryptosporidium* species in humans and animals: Current understanding and research needs. Parasitology. 2014;141(13):1667-1685. doi:10.1017/S0031182014001085.
34. Okojokwu OJ, Ileigo IH, Ezemuel YS, Oluyinka OO, Akpakpan EE, Kolawole T, et al. Molecular Characterisation of *Cryptosporidium* Species from Extensively Managed Cattle

Slaughtered in Abattoirs in Kaduna State, Nigeria. *Advances in Applied Science Research*. 2016; 7(1):17-22.

35. Chukwu EV, Olayemi OD, Mohammed BR, Opara NM and Agbede RIS. Research trends and prevalence of *Cryptosporidium* infections in animals in Nigeria: A review. *Nigerian Journal of Parasitology*. 2019; 40(1):103-109.

UNDER PEER REVIEW

UNDER PEER REVIEW