

Original Research Article

EFFECT OF ORGANIC AND INORGANIC NUTRIENT SOURCES ON GROWTH AND YIELD OF MAIZE (*Zea mays* L.)

Abstract: The present investigation titled “Effect of Organic and Inorganic Nutrient Sources on Growth and Yield of Maize” was conducted during *khariif* season of 2022 at Chamelti Agriculture Farm, MS Swaminathan School of Agriculture, Shoolini University, Solan, and Himachal Pradesh. The field experiment was laid out in Randomized Block Design comprising of 10 treatments and 3 replications which are (T₁) Control, (T₂) 100% RDF, (T₃) 120% RDF, (T₄) 50% RDF + FYM (10 t), (T₅) 75% RDF + FYM (5 t), (T₆) 100% RDF + FYM (5 t) + Mulch, (T₇) 50% RDF + Vermicompost (2.5 t ha⁻¹), (T₈) 75% RDF + Vermicompost (2.5 t ha⁻¹), (T₉) 100% RDF + Vermicompost (2.5 t ha⁻¹) + Mulch and (T₁₀) 75% RDF + FYM (5 t) + Vermicompost (2.5 t ha⁻¹) + Mulch. Mustard stover @ 2 t ha⁻¹ was used as mulching. RDF used is (100:40:40 kg ha⁻¹) was applied through urea (46% N), SSP (16% P₂O₅) and MOP (60% K₂O). One third N and full dose of P and K was applied at the time of sowing as basal application. Remaining nitrogen was applied in 2 equal splits at 30 and 50 DAS as top dressing. PSC-3322 variety of maize was used for sowing. Other crop management practices were followed as per the recommendation of the area. This study concluded that application of 120 % RDF recorded significantly higher plant height, yield and was economically better than other treatments.

Comment [HT1]: Keywords ?

INTRODUCTION

The rising population and consumption, reduction in available land and other productive units are placing unprecedented pressure on the current agriculture and natural resources to meet the increasing food demand. Achieving food security under sustainable system possesses a significant challenge in the developing world and is highly critical for alleviating poverty. To circumvent this challenge, crop producers tended to over use certain inputs such as chemical fertilizers and pesticides which in turn have already started deteriorating environment and soil as well. To meet the world's future food security and sustainability needs, food production must grow substantially, while the negative impact of agriculture on environment must shrink dramatically at the same time (Foley *et al.*, 2011). In India, maize is the third most important cereal crop after rice and wheat. It has got immense potential

therefore, called as “Miracle crop” and also as “Queen of Cereals”. Maize, with its high content of carbohydrates, fats, proteins, some of the important vitamins and minerals has acquired a well-deserved reputation as a poor man’s ‘nutricerea’ and contributes more than 9% to national food basket. Maize grain has elevated nutritive value as it contains about 72% starch, 10% protein, 4.8% oil, 5.8% fiber and 3% sugar (Rafiq *et al.*, 2010). The consumption pattern for maize produced in India at present includes poultry feed (52%), human food (24%), animal feed (11%), starch (11%), brewery (1%) and seed (1%) (Dass *et al.*, 2007).

It occupies 9.86 million ha area and production of 31.51 million tonnes of production with average productivity of 3195 kg ha⁻¹ (Agricultural Statistics at a Glance, 2021). The area, production and productivity of maize in Himachal Pradesh is 26.74 thousand ha, 725014 metric ton and 2730 kg ha⁻¹ respectively. In Solan district of

Comment [HT2]: Update the latest 2022 or 2023.

Himachal Pradesh total area under maize is 22435 ha with the production of 73276 metric tonnes and average productivity of about 3270 kg ha⁻¹ (Statistical Abstract of Himachal Pradesh, 2021-22).

It is evident that the productivity of maize in tropical nations is constrained due to inherently poor soil fertility, low soil organic matter and further more low water holding capacity. Maize crop has a higher nutrient uptake character and leaves soil exhaustive. The method of nutrition and its management plays a crucial role in production of maize. The cereal production versus fertilizer consumption of India indicates low fertilizer use efficiency (Prasad, 2009).

Farmyard manure is a traditional, well known, readily available and widely used input since time immemorial. It is a conspicuous organic component of an integrated nutrient supply system, which improves soil health, increases the productivity and releases macro and micronutrients. It is costlier than chemical fertilizers on nutrient basis but other beneficial effect on soil aggregates, cation exchange capacity, water holding capacity, fertilizers use efficiency, microbial activity and nutrient availability in soil (Sharma *et al.*, 2004).

Vermicompost plays a significant role in improving the fertility of top soil and in boosting the productivity of the crop. Vermicompost has also been advocated as good organic manure for use in integrated nutrient management practices in field crops (Singh and Nepalia, 2009). It is proven fact that productivity of any crop cannot be further increased by use of high doses of fertilizer alone. Balanced nutrition through right proportion of organic manures and chemical fertilizers is essential for boosting QPM production and sustaining soil productivity (Singh *et al.*, 2017).

MATERIALS AND METHODS

The present research work titled “Effect of Organic and Inorganic Nutrient Sources on Growth and Yield of Maize (*Zea mays* L.)” was carried out during kharif season of 2022

at Chamelti Agriculture Farm, MS Swaminathan School of Agriculture, Shoolini University of Biotechnology and Management Sciences, Solan. The experimental plot was assigned well-drained soil which had homogenous fertility and textural arrangement. Geographically, Chamelti Agriculture Farm is situated 30 km away from Solan city at an elevation of 1,270 meters above mean sea level lying between latitude 30° 85'67.30 N and longitude 77° 13'20.38 E. It falls under the mid-hill zone of Himachal Pradesh. The field of the experimental site represented ideal spatial unit in respect of texture, make up and fertility status. The soil of the experimental site was sandy loam in texture, slightly alkaline in reaction with EC in safer range, medium in organic carbon, available nitrogen, potassium and high in available phosphorus. The field experiment was laid out in Randomized Block Design comprising ten treatments and replicate thrice. The experiment consists (T₁) Control, (T₂) 100% RDF, (T₃) 120% RDF, (T₄) 50% RDF + FYM (10 t), (T₅) 75% RDF + FYM (5 t), (T₆) 100% RDF + FYM (5 t) + Mulch, (T₇) 50% RDF + Vermicompost (2.5 t ha⁻¹), (T₈) 75% RDF + Vermicompost (2.5 t ha⁻¹), (T₉) 100% RDF + Vermicompost (2.5 t ha⁻¹) + Mulch and (T₁₀) 75% RDF + FYM (5 t) + Vermicompost (2.5 t ha⁻¹) + Mulch. Mustard stover @ 2 t ha⁻¹ was used as mulching. Recommended dose of nitrogen, phosphorus and potassium (100:40:40 kg ha⁻¹) was applied through urea (46% N), SSP (16% P₂O₅) and MOP (60% K₂O). One third nitrogen and full dose of phosphorus and potassium was applied at the time of sowing as basal application. Remaining nitrogen was applied in two equal splits at 30 and 50 DAS as top dressing. However, FYM and Vermicompost were applied one month before sowing. PSC-3322 variety of maize was used for sowing. Other crop management practices were followed as per the recommendation of the area.

STATISTICAL ANALYSIS

The data presented in the thesis are the mean values. All the observations are statistically analyzed by using the analysis

of variance. The results were tested for the treatments mean by applying F- test of significance on the basis of null hypothesis (Cochran and Cox, 1957). Wherever necessary, standard errors along with critical difference at 5 % of significance were computed for discriminating the treatment effects for chance effects.

RESULTS AND DISCUSSION

A. Growth parameters

Different growth parameters such as plant height and dry matter accumulation were showed in (Table 1). Among treatments, (T₃) 120% RDF recorded significantly higher plant height (40.50 cm) which was statistically at par with (T₉) 100% RDF + Vermicompost (2.5 t ha⁻¹) + Mulch, (T₆) 100% RDF + FYM (5 t ha⁻¹) + Mulch and (T₂) 100% RDF, respectively. However, least plant height was noted under (T₁) control treatment. The higher level of nitrogen increased the availability and absorption of nitrogen which resulted in more vegetative growth due to increase in plant height on account of enlargement of cells and increased photosynthesis Mohamoud *et al.* (2002); (Tiwana *et al.*, 2003); Sobhana *et al.* (2012); Gul *et al.* (2015) and Wadile *et al.* (2016).

Whereas application of (T₃) 120% RDF recorded significantly higher dry matter accumulation (18.33 g plant⁻¹) which was statistically at par with (T₉) 100% RDF + Vermicompost (2.5 t ha⁻¹) + Mulch, (T₆) 100% RDF + FYM (5 t ha⁻¹) + Mulch and (T₂) 100% RDF, respectively. However, least dry matter accumulation (11.98 g plant⁻¹) was noted under (T₁) control treatment.

B. Yield attributes

Data on the effect of organic and inorganic nutrient sources on various yield attributes parameters of maize have been presented in Table 2. Among the treatments, application of (T₃) 120% RDF recorded significantly higher number of cobs plant⁻¹ (2.20), cob length (21.45 cm), number of grains cobs⁻¹ (221.45) and seed index (26.51 g) however, least number of cobs plant⁻¹ (1.20), cob length (15.94 cm), number of grains cobs⁻¹

(142.35) and seed index (26.05 g) was noted under (T₁) control treatment. Maximum number of rows cob⁻¹ was recorded under the application of 120 % RDF through inorganic source mainly due to more availability and steady nutrients release. Use of fertilizer did bring about significant improvement in overall growth of the crop by providing needed nutrients from initial stage and increase in supply of N, P and K in more synchronize way. These findings are corroborating the results of Iqbal *et al.* (2014) and Nagavani and Subbian (2014) in maize.

C. Yield

Data on the effect of organic and inorganic nutrient sources on yield of maize have been presented in Table 3. Among the treatments, application of (T₃) 120% RDF recorded significantly higher grain yield (3753 kg ha⁻¹), stover yield (13511 kg ha⁻¹), biological yield (17264 kg ha⁻¹) and harvest index (21.74%) however least grain yield (1805 kg ha⁻¹), stover yield (6697 kg ha⁻¹), biological yield (8502 kg ha⁻¹) and harvest index (21.23%) was noted under (T₁) control treatment. The probable reason for these results might be attributed to better nitrogen availability. Nitrogen being a major constituent of chlorophyll molecule, might have played a positive role in increasing the photosynthetic activity and ultimately reflected in the acceleration of meristematic activity and increase in the yield. The significant positive correlation between yield and various other parameters have indicated the positive response of higher dose fertilizers on various other parameters. The results of present investigation are in close agreement with the findings of Girija Devi (2002); Dudhat *et al.* (2004); Hani *et al.* (2006); Kumar *et al.* (2008); Rizwan *et al.* (2008); Ramanjaneyulu *et al.* (2010) and Reddy and Bhanumurthy (2010).

D. Economics

Data on the effect of organic and inorganic nutrient sources on Economics is showed in Table 4. Among the treatments, application of (T₃) 120% RDF recorded significantly higher gross returns (₹ 123099 ha⁻¹), net

returns (₹ 90513 ha⁻¹) and B: C ratio (2.78) The data clearly revealed that higher gross returns, net returns and B:C ratio were observed under application of (T₃) 120% RDF over rest of the treatments. This might be due to higher yield and least cost of cultivation. Integration of vermicompost and

FYM with inorganic source in different proportion recorded low net realization and B:C ratio mainly due to high cost of manure over 120 % RDN through inorganic source. These results are in accordance with the findings Meena *et al.* (2013) and Nagavani and Subbian (2014).

Table 1. Effect of organic and inorganic nutrient sources on growth parameters

| Treatments | Plant Height at harvest (cm) | Dry Matter at harvest (g plant ⁻¹) |
|--|------------------------------|--|
| T ₁ : Control | 177.63 | 82.43 |
| T ₂ : 100% RDF | 212.90 | 98.79 |
| T ₃ : 120% RDF | 222.00 | 103.02 |
| T ₄ : 50% RDF + FYM (10 t ha ⁻¹) | 192.45 | 89.30 |
| T ₅ : 75% RDF + FYM (5 t ha ⁻¹) | 197.90 | 91.83 |
| T ₆ : 100% RDF + FYM (5 t ha ⁻¹) + Mulch | 216.25 | 100.35 |
| T ₇ : 50% RDF + Vermicompost (2.5 t ha ⁻¹) | 196.69 | 91.27 |
| T ₈ : 75% RDF + Vermicompost (2.5 t ha ⁻¹) | 198.39 | 92.06 |
| T ₉ : 100% RDF + Vermicompost (2.5 t ha ⁻¹) + Mulch | 219.20 | 101.72 |
| T ₁₀ : 75% RDF + FYM (5 t ha ⁻¹) + Vermicompost (2.5 t ha ⁻¹) + Mulch | 199.90 | 92.76 |
| <i>SEm±</i> | 6.60 | 3.11 |
| <i>LSD (p=0.05)</i> | 19.91 | 9.00 |

Table 2. Effect of organic and inorganic nutrient sources on yield attributes

| Treatments | Cobs plant ⁻¹ | Cob length (cm) | Grains cob ⁻¹ | Seed index (g) |
|--|--------------------------|-----------------|--------------------------|----------------|
| T ₁ : Control | 1.20 | 15.94 | 142.35 | 26.05 |
| T ₂ : 100% RDF | 2.00 | 19.12 | 194.31 | 26.32 |
| T ₃ : 120% RDF | 2.20 | 21.45 | 221.45 | 26.51 |
| T ₄ : 50% RDF + FYM (10 t ha ⁻¹) | 1.40 | 17.24 | 174.63 | 26.14 |
| T ₅ : 75% RDF + FYM (5 t ha ⁻¹) | 1.60 | 18.31 | 184.65 | 26.21 |
| T ₆ : 100% RDF + FYM (5 t ha ⁻¹) + Mulch | 2.00 | 19.42 | 197.37 | 26.39 |
| T ₇ : 50% RDF + Vermicompost (2.5 t ha ⁻¹) | 1.40 | 17.61 | 180.61 | 26.18 |
| T ₈ : 75% RDF + Vermicompost (2.5 t ha ⁻¹) | 1.60 | 18.49 | 188.85 | 26.26 |
| T ₉ : 100% RDF + Vermicompost (2.5 t ha ⁻¹) + Mulch | 2.00 | 19.78 | 200.65 | 26.44 |
| T ₁₀ : 75% RDF + FYM (5 t ha ⁻¹) + Vermicompost (2.5 t ha ⁻¹) + Mulch | 1.60 | 18.71 | 191.98 | 26.29 |

| | | | | |
|------------------------------|------|------|-------|------|
| <i>SEm</i> ± | 0.06 | 0.39 | 74.58 | 0.18 |
| <i>LSD</i> (<i>p</i> =0.05) | 0.21 | 1.24 | 21.54 | NS |

Table 3. Effect of organic and inorganic nutrient sources on economic

| Treatments | Yield (kg ha ⁻¹) | | | Harvest index (%) |
|--|------------------------------|--------------|------------------|-------------------|
| | Grain yield | Stover yield | Biological yield | |
| T ₁ : Control | 1805 | 6697 | 8502 | 21.23 |
| T ₂ : 100% RDF | 3315 | 12133 | 15448 | 21.46 |
| T ₃ : 120% RDF | 3753 | 13511 | 17264 | 21.74 |
| T ₄ : 50% RDF + FYM (10 t ha ⁻¹) | 2231 | 8255 | 10486 | 21.28 |
| T ₅ : 75% RDF + FYM (5 t ha ⁻¹) | 2815 | 10387 | 13202 | 21.32 |
| T ₆ : 100% RDF + FYM (5 t ha ⁻¹) + Mulch | 3451 | 12562 | 16013 | 21.55 |
| T ₇ : 50% RDF + Vermicompost (2.5 t ha ⁻¹) | 2416 | 8939 | 11355 | 21.28 |
| T ₈ : 75% RDF + Vermicompost (2.5 t ha ⁻¹) | 2942 | 10856 | 13798 | 21.32 |
| T ₉ : 100% RDF + Vermicompost (2.5 t ha ⁻¹) + Mulch | 3542 | 12857 | 16399 | 21.6 |
| T ₁₀ : 75% RDF + FYM (5 t ha ⁻¹) + Vermicompost (2.5 t ha ⁻¹) + Mulch | 3025 | 11132 | 14157 | 21.37 |
| <i>SEm</i> ± | 103 | 421 | 536 | 0.43 |
| <i>LSD</i> (<i>p</i> =0.05) | 312 | 1274 | 1619 | NS |

Table 4. Effect of organic and inorganic nutrient sources on yield

| Treatments | Economics (₹ ha ⁻¹) | | | B:C ratio |
|--|---------------------------------|---------------|-------------|-----------|
| | Cost of cultivation | Gross returns | Net returns | |
| T ₁ : Control | 24000 | 59801 | 35801 | 1.49 |
| T ₂ : 100% RDF | 31155 | 109329 | 78174 | 2.51 |
| T ₃ : 120% RDF | 32586 | 123099 | 90513 | 2.78 |
| T ₄ : 50% RDF + FYM (10 t ha ⁻¹) | 34578 | 73847 | 39269 | 1.14 |
| T ₅ : 75% RDF + FYM (5 t ha ⁻¹) | 32866 | 93091 | 60225 | 1.83 |
| T ₆ : 100% RDF + FYM (5 t ha ⁻¹) + Mulch | 34655 | 113608 | 78953 | 2.28 |
| T ₇ : 50% RDF + Vermicompost (2.5 t ha ⁻¹) | 40078 | 79969 | 39891 | 1 |
| T ₈ : 75% RDF + Vermicompost (2.5 t ha ⁻¹) | 41866 | 97292 | 55426 | 1.32 |
| T ₉ : 100% RDF + Vermicompost (2.5 t ha ⁻¹) + | 43655 | 116495 | 72840 | 1.67 |

| | | | | |
|--|-------|-------|-------|-----|
| Mulch | | | | |
| T ₁₀ : 75% RDF + FYM (5 t ha ⁻¹) + Vermicompost (2.5 t ha ⁻¹) + Mulch | 45366 | 99946 | 54580 | 1.2 |

UNDER PEER REVIEW

CONCLUSION

On the basis of experimental finding summarized above, the following conclusions are drawn: Marked improvement in growth, yield attributes and yield of maize were observed with application of (T₃) 120% RDF which was statistically at par with (T₉) 100% RDF + Vermicompost (2.5 t ha⁻¹) + Mulch, (T₆) 100% RDF + FYM (5 t ha⁻¹) + Mulch and (T₂) 100% RDF over rest of the treatments.

On the basis of B: C ratio, application of (T₃) 120% RDF was found to be remunerative for maize under Mid hills of Himachal Pradesh.

REFERENCE

- Agricultural Statistics at a Glance. Government of India. Ministry of Agricultural & Farmers Welfare. Department of Agriculture, Cooperation & Farmers Welfare. Directorate of Economics and Statistics. 2021.
- Dass, S., Singh, K. P. and Yadav, V. K. Present status and potential of maize hybrids in enhancing the productivity. National Conference on-Doubling Maize Production Organized by IFFCO Foundation, ICAR, DMR, DAC and IFFCL at New Delhi. 2007; 5: 13-19.
- Dudhat, M.S., Savalia, M.G. and Ramdevputra, M.V. Response of forage maize to nitrogen and phosphorus levels. Forage Research. 2004; 30: 34-35.
- Foley, J.A., Ramankutty, N., Brauman, K.A., Cassidy, E.S., Gerber, J.S. and Johnston, M. Solutions for a cultivated planet Nature. 2011; 478(7369): 337-342.
- Girija Devi, L. Forage yield of maize (*Zea mays* L.) as influenced by nitrogen levels and biofertilizers. Forage Research. 2002; 27: 263-266.
- Gul, S., Khan, M.H, Khanday, B.A. and Nabi S. Effect of sowing methods and NPK levels on growth and yield of rainfed maize (*Zea mays* L.). Scientifica. 2015; 1-6.
- Hani, E.A., Hamed, M.A. and Ali. E.E. The effect of nitrogen and phosphorus fertilization on growth, yield and quality of forage maize. Journal of Agronomy. 2006; 5: 515-518.
- Iqbal, A., Iqbal, M. A., Raza, A., Akbar, N., Abbas, R. N. and Khan, H.Z. Integrated Nitrogen Management Studies in Forage Maize. American-Eurasian Journal Agricultural & Environment Science. 2014; 14(8): 744-747.
- Kumar, A., Singh, R., Rao, K. L. and Singh, U. K. Effect of integrated nitrogen management on growth and yield of maize (*Zea mays* L.) cv. PAC- 711. Madras Agricultural Journal. 2008; 95(7-12): 467-472.
- Meena, S. K. Mundra S. L. and Singh, P. Response of maize (*Zea mays*) to nitrogen and zinc fertilization. Indian Journal of Agronomy. 2013; 58(1): 127- 128.
- Mohamoud, A.K., Sharanappa, I.S. and Reddy, P.J. Effect of compost and fertilizer levels on the structure of growth and yield in maize. Madras Agriculture Journal. 2002; 89(10-12): 720-723.
- Nagavani, A. V. and Subbian, P. Productivity and economics of hybrid maize as influenced by integrated nutrient management. Current Biotica. 2014; 7(4): 283-293.
- Prasad R. Efficient fertilizer use: the key to food security and better environment. Journal of Tropical Agriculture. 2009; 47 (1): 1-17.
- Rafiq, M.A., Ali A., Malik, M.A. and Hussain, M. Effect of fertilizer level and plant densities on yield and protein contents of autumn planted maize. Pakistan Journal of Agricultural Science. 2010; 47(3): 201-208.
- Ramanjaneyulu, A.V., Giri, G. and Kumar, S.R. Biofertilizer, nitrogen and phosphorus on yield and nutrient economy in forage sorghum affected by nutrient management in preceding mustard. International Journal of Bio-resource Management. 2010; 1: 66-68.

Comment [HT4]: Spelling mistake and check all the references and add more following references as well as for quality of papers.

- Reddy, D.M. and Bhanumurthy, V.B. Fodder, grain yield, nitrogen uptake and crude protein of forage maize as influenced by different nitrogen management practices. *International Journal of Bio-resource Management*. 2010; 1: 69-71.
- Rizwan, A., Muhammed, A., Azeem, K. and Zahir, A. Effectiveness of organic biofertilizer supplemented with chemical fertilizers for improving soil water retention aggregates stability, growth and nutrient uptake of maize. *Journal of Sustainable Agriculture*. 2008; 31: 57-77.
- Sandon, F. (1958). *Experimental Designs*. By WG Cochran and GM Cox. Pp. xiv, 611. 82s. 1957.(John Wiley and Sons, New York; Chapman and Hall, London). The Mathematical Gazette, 42(342),334-334.
- Sharma, V., Kanwar, K and Dev, S.P. Efficient recycling of obnoxious weed plants (*Lantana camara* L.) and congress grass (*Parthenium hysterophorus* L.) as organic manure through vermicomposting. *Journal of The Indian Society of Soil Science*. 2004; 52(1): 112-114.
- Singh, D. and Nepalia, V. Influence of integrated nutrient management on quality protein maize (*Zea mays*) productivity and soils of southern Rajasthan. *Indian Journal of Agricultural Sciences*. 2009; 79(12): 1020-1022.
- Singh, L., Kumar, S., Singh, K. and Singh, D. Effect of integrated nutrient management on growth and yield attributes of maize under winter season (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*. 2017; 6 (5): 1625- 1628.
- Sobhana, V., Kumar, A., Idnani, L.K., Singh, I. and Shivadhar. Plant population and nutrient requirement for baby corn hybrids (*Zea mays* L.). *Indian Journal of Agronomy*. 2012; 57(3): 294-296
- Statistical Abstract of Himachal Pradesh. Department of Economic and Statistics. Government of Himachal Pradesh, Shimla. 2021-22.
- Tiwana, U.S., Puri. K.P. and Singh, S. Fodder yield and quality of multicut pearl millet as influenced by nitrogen and phosphorus under Punjab conditions. *Forage Research*. 2003; 28: 190-193.
- Wadile, S.C., Pawar, P.P., Iihe, S.S. and Rathod, V.M. Nutrient management on growth, yield and quality of sweet corn, baby corn and maize. *Bioinfolet*. 2016; 13(1A): 67-69.

Add the following references are compulsory:

- Singh, D., Yadav, A., Tiwari, H., Singh, A. K., Singh, S., Yadav, R. K., ... & Sachan, D. S. (2023). Nitrogen Management through Nano Urea and Conventional Urea and its Effect on Wheat (*Triticum aestivum* L.) Growth and Yield. *International Journal of Plant & Soil Science*, 35(18), 1466-1473.
- Kumar, S., Agrawal, S., Jilani, N., Kole, P., Kaur, G., Mishra, A., ... & Tiwari, H. (2023). Effect of integrated nutrient management practices on growth and productivity of rice: A review.
- Tiwari, H., Naresh, R. K., Singh, P. K., & Kumar, Y. (2023). Effect of Integrated Nutrient Management on Growth and Productivity of Wheat (*Triticum aestivum* L.) in Typic Ustochrepts Soils of Western UP, India. *International Journal of Plant & Soil Science*, 35(11), 129-142.

UNDER PEER REVIEW