

## Original Research Article

### The Affects of Smoking on Cataract Formation in Turkish Patients

#### ABSTRACT

**Aims:** To investigate whether smoking influence on cataract formation age and cataract type in patients who underwent senile cataract surgery.

**StudyDesign:**Retrospectivecross-sectional.

**Place and Duration of Study:** The study conducted in Department Of Ophthalmology, Ahi Evran Faculty Of Medicine between November 2014 and March 2018.

**Methodology:**Five hundred eighty-one patients were included, who were operated for senile cataracts. Age, gender, cataract type and smoking status of patients were recorded.Congenital, traumatic, drug induced and uveitic cataracts were excluded from the study. The smoking status of patients were divided into three groups:currently smoking, never smoked, quitted smoking. The smoking load of patients were calculated as packages/year.

**Results:**Of totally 581 patients, 47.8% (278) were female and 52.2% (303) were male and mean ages were  $66.68 \pm 10.68$  years and  $66.17 \pm 9.37$  years, respectively. There was an association between gender and smoking status ( $p < 0.05$ ).Smoking rate (both currently smoking and quit smoking) was higher among males. The mean age of cataract patients who were smokers was statistically lower than the mean ages of non-smokers and quitters ( $p < 0.001$ ).There was a statistical association between cataract type of patients and smoking status ( $p = 0.0152$ ). This was resulted from lower incidence of cortical cataracts and higher incidence of posterior subcapsular and nuclear cataracts in smokers.Among these three groups, cortical cataract was the most frequent in non-smoker group. It was found that smoking increases the posterior subcapsular cataract, but it has less effect on cortical cataract formation.

**Conclusion:** Smoking causes cataract formation in younger ages and higher posterior subcapsular cataracts rate. We suggest that this finding may prevent beginning to smoking and increase the motivation of individuals to quit smoking.

**Key Words:** Cataract, cataract type, cigarette cessation, smoking.

## 1. INTRODUCTION

At the present time, the most prevalent cause of treatable blindness is cataract. The data of WHO (World Health Organization) in 2014 indicated that 35% of blindness in the world is resulted from the cataracts. According to WHO estimates, 54 million people over the 60 years of age will have low vision because of cataracts by 2020 [1]. Although surgical removal is the only available treatment for the cataract, identifying the risk factors to take protective measures is also important. Aging is the most important known risk factor for cataracts. The other risk factors are genetic disposition, diabetes, ultraviolet radiation, corticosteroid use, hypoparathyroidism, trauma and persistent intraocular inflammation[2,3].

One of the factors accused for cataracts formations is smoking. After prolonged smoking, ocular side effects in addition to systemic diseases and various malignancies caused by chemicals ingredients are known so far[4,5]. These ocular side effects shown previously include irritation to ocular surface, cataract formation and progression, increased intraocular pressure, vasoconstriction in ophthalmic artery, decreased ocular perfusion pressure, higher risk, rate and severity of Grave's ophthalmopathy, formation and progression of age-related macular degeneration[6-9]. As it is a preventable risk factor, the association between smoking and cataracts is important. Avoiding smoking may delay cataract formation and progression, as well as decrease financial and clinical load needed for cataract surgery considerably.

To our knowledge, there is no study investigating the relationship between smoking and cataract formation in Turkish patients. In this study, it is therefore aimed to investigate whether smoking influence cataract formation age and cataract type in patients who underwent senile cataract surgery.

## 2. MATERIALS AND METHODS

Five hundred eighty-one patients who were operated for senile cataracts between November 2014

and March 2018 in Ahi Evran University, Faculty of Medicine, Department of Ophthalmology were evaluated retrospectively. Information about the study was provided to patients and written consent was obtained from all subjects. The age, gender, cataract type and smoking status of patients were recorded. Use of smoking and alcohol was questioned again by phone-calls and missing information was completed. Patients who cannot be reached by phone-call or who use alcohol were excluded from the study. Information about remaining 581 patients were recorded.

All the patients underwent full ophthalmological examination. Patients who has 0.3 or less visual acuity according to decimal Snellen chart were operated. Cataract type was recorded by slit lamp examination after pupillary dilatation. The cataract classification was made according to its anatomical location by Brown and Hill [4] and Lens Opacities Classification System III [5] which included nuclear, cortical and posterior subcapsular cataracts (PSC). Mixed type and mature cataracts were not included in the study. Congenital, traumatic, drug induced and uveitic cataracts were excluded from the study.

The smoking status of patients were divided in three groups: currently smoking, never smoked, quitted smoking. The smoking load of patients were calculated as packages/year (grouped as  $\leq 5$  packages/year, 6-10 packages/year, 11-20 packages/year, 21-30 packages/year, 31-40 packages/year,  $\geq 41$  packages/year).

## **2.1 Statistical Analysis**

The associations between cataract type and smoking status, age, gender, time since quitting, smoking load (packages/year) were investigated. Categorical variables were given as number and percentage, numerical variables were given as mean - standard deviation and as median and minimum-maximum values. Normal distribution of numerical variables were tested by Shapiro– Wilks test and when assumption was validated analysis of one way variance (ANOVA) and independent T tests were used for comparisons between groups and when normal distribution was not validated Mann-Whitney U test were used. When a difference was found among the groups, Bonferroni pair comparison test was used to find out the group that causing the difference. Chi-square test was used to compare categorical variables. All the analysis was conducted by IBM SPSS 22.0 package program (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.), and  $p < 0.05$  was accepted as statistical significance for all analysis.

### 3.RESULTS

The demographical data, cataract types and smoking status of the patients were summarized in Table 1 and Table 2.

There was an association between the gender and the smoking status ( $p < 0.05$ ). The smoking rate (both currently smoking and quitted smoking) was higher in males and female ration was higher in the non-smoker group (Table 3).

The mean age of smokers was lower than the mean age of non-smokers and quitters and this was statistically significant ( $p < 0.001$ ) (Table 4).

The associations between smoking status and Diabetes Mellitus (DM), and Hypertension (HT) were also investigated. There was no statistical difference between the smoking status and cataract type in patients with DM ( $p = 0.807$ ). Similarly, there was no statistical difference between the smoking status and cataract type in patients with and without HT [respectively  $p = 0.192$  (smoking- cataract type comparison in patients with HT) / and  $p = 0.067$  (smoking- cataract type comparison in patients without HT)].

There was statistically significant difference between the cataract type and the smoking status of patients ( $p = 0.0152$ ). This difference was due to less frequent cortical cataract and more frequent PSC and nuclear cataracts in currently smoking group. Additionally the frequency of cortical and PSC cataract type in the quitter group was between the frequency of currently smokers and nonsmokers (Table 5). Almost half of the cataracts in currently smoker group was PSC cataract and cortical cataract was the least frequent type. In the quitter group, nuclear and PSC frequency was 41%. The frequency of PSC and nuclear cataract types were similar in the nonsmoker group. Among the three groups cortical cataract was the most prevalent in the nonsmoker group. With smoking, the PSC frequency was increased and it is found that cortical cataract was less affected by smoking (Table 5).

When the association between the gender and cataract type was evaluated within smoking status, the frequencies of nuclear and PSC type was more prevalent in smoking males compared to smoking females ( $p < 0.05$ ).

The association between smoking load of patients (packages year) and cataract type was also analyzed. According to the results, there was no statistically significant association between the smoking load and cataract type ( $p=0.157$ ).

#### **4.DISCUSSION**

Cigarette smoke is a mixture of thousands of chemicals including the particles such as nicotine, volatile chemicals such as formaldehyde and gases such as carbon-monoxide[4]. In addition to reported side effects of smoking on almost all layers of the eye[6-9], many molecular mechanisms were suggested for enhancing cataract formations[10-19]. One of those mechanisms is impairment of alpha crystallin protein structure which is a major protein in lens. In Alpha crystalline proteins there are two sub-units as Alpha-A and Alpha-B[10]. Alpha crystalline has a chaperon function and it has a significant role in lens transparency. Alpha crystalline prevent protein aggregates which may cause light scattering by binding denaturated proteins with is chaperon feature[11]. However, chaperon like activity of alpha crystalline may be disturbed or lost by various genetical and environmental factors, aging and some clinical conditions. Ultimately, the lens loses its transparency[11]. In a recent experimental study, p-parabenzoquinone was detected in porcine lenses and lenses obtained from smoking humans. It was shown that p-parabenzoquinone caused cataracts by disturbing the structure of alpha crystalline[12].

The epithelial cells are the first line cells in the lens which are vulnerable to many damages causing cataracts because of their superficial localization. These cells are well-equipped to protect the lens from harmful factors and metabolically they are quite active. Therefore, the epithelial cells were investigated by many researchers for cataract pathogenesis related to age and disease[13]. It is suggested that oxidative stress may trigger cataract formation by causing changes in lens epithelium[14]. Smoking may cause additional oxidative stress by indirectly decreasing antioxidant molecules like ascorbic acid and nicotinamide[15,16]. In addition, heavy metals in the cigarette like lead and cadmium may cause oxidative stress by depletion of glutathione and thiol pool and may disturb antioxidant protections system. A previous study reported that cadmium caused apoptosis in human lens epithelial cells[14-19]. In another study, hyperplasia, hypertrophy and stratification of lens epithelial cells were observed in histopathological examination of rat lenses which exposed to cigarette smoking 2 hours a day for 60 days[20]. Additionally, zinc level was decreased, and iron and copper levels were increased in lenses. It was thought that oxygen radicals produced by Fenton's reaction of

that increased metal load and increased oxidative stress may cause lens damage[20].In our study, currently smokers and quitters had cataracts in younger ages compared to nonsmoker group.We suggest that it might be caused by earlier aging of lens because of the reasons discussed above.

Many studies were conducted previously about the effect of smoking on cataract formation and cataract types. Among those studies, some reported no effect on cataract type while the others reported the opposite. Many researchers reported that smoking had a positive association with nuclear and PSC cataract formation while it had no effect on cortical cataract formation[21-24].In Blue Mountains Study including 3645 people, the effect of smoking and alcohol use on cataract were investigated and nuclear cataract prevalence was found to be higher than PSC prevalence in smokers and alcohol users. It is reported that the cataract formation risk was much higher in people who both smoke and use alcohol[21].Prospective studies including large number of cases like Physicians Health Study (17824 cases) [22]and Nurses Health Study and Health Professionals Follow up Study (124690 cases) [23] reported that smoking is a risk factor for PSC cataract.Similarly, in our study PSC rate was significantly higher in smokers compared to nonsmokers.

Many studies showed that cataract risk decreased after quitting smoking[25-27].One of the most extended information about the cataract formation in people who quit smoking was in 2 studies of Lindblad et al. [25,26].In this prospective study including 34595 females, when compared to nonsmokers, 10 years after quitting smoking in people who smoked 6-10 cigarettes a day and 20 years after quitting smoking in people who smoked 10-15 cigarettes a day, there was no statistical difference for cataract surgery need.For smokers who smoke 15 cigarettes a day or more, after quitting smoking, cataract formation risk was continued even the risk decreased[26].Likewise, decreased PSC rate in quitters in our study showed that cataract risk may decrease after quitting smoking. In a prospective study with 5 years follow up, of 17824 doctors between 45 and 84 years of age, 33% was smokers, 39% was quitters and 51% was nonsmokers. It is reported that nuclear and PSC rate was increased in people who smoke 20 or more cigarettes a day. They reported that while nuclear cataract formation risk decreases in people who quit smoking, PSC rate did not change[22].In our study, while a significant association between smoking and PSC formation was found, there was no effect on cortical cataract formation. In the people who quit smoking, cataract formation age was higher and PSC rate was decreased.

## **5.CONCLUSION**

The pathogenesis of cataract formation is influenced by many factors including genetic and environmental factors. We are aware of this is the first study investigating smoking affects on cataract formation in Turkish patients. In this study, it was shown that smoking caused higher PSC rate and cataract surgery in earlier age.As it is reported that10 years delay in cataract formation decreased the cataract surgery rate and related costs by 50% [28], we suggest that more critical public health care policies should be raised to prevent andquit smoking

## **ETHICALAPPROVAL**

The study protocol was approved by the Ahi Evran University Faculty of Medicine Clinical Research Ethics Committee (Ethical approval number: 2018-07/69).

## **CONSENT**

It is not applicable

## **REFERENCES**

1.Casparis H, Lindsley K, Kuo IC, Sikder S, Bressler NB. Surgery for cataractsin people with age-related macular degeneration. Cochrane Database Syst Rev. 2012;6:CD006757.

2. Skalka HW, Prchal JT. Effect of corticosteroids on cataract formation. *Arch Ophthalmol* 1980;98:1773-7.
3. Glynn RJ, Christen WG, Manson JE, Bernheimer J, Hennekens CH. Body mass index. An independent predictor of cataract. *Arch Ophthalmol*. 1995;113:1131-7.
4. Bartecchi CE, MacKenzie TD, Schrier RW. The human costs of tobacco use. *N Engl J Med* 1994;330:907-12.
5. Toll BA, Rojewski AM, Duncan LR, et al. "Quitting smoking will benefit your health": the evolution of clinician messaging to encourage tobacco cessation. *Clin Cancer Res*. 2014;20:301-9.
6. Avunduk AM, Avunduk MC, Evirgen O, et al. Histopathological and ultrastructural examination of the rat conjunctiva after exposure to tobacco smoke. *Ophthalmologica*. 1997;211:296-300.
7. Lietz-Partzsch A, Griesser SM, Flammer J, Haefliger IO. Decreased Heidelberg Retina Flowmeter (HRF) parameter "flow" at the papilla shortly after smoking a cigarette. *Klin Monbl Augenheilkd*. 2001;218:332-4.
8. Klein R, Klein BE, Linton KL, DeMets DL. The Beaver Dam Eye Study: the relation of age-related maculopathy to smoking. *Am J Epidemiol*. 1993;137:190-200.
9. Çankaya AB, Elgin U, Şimşek T, Batman A. The effect of smoking on intraocular pressure in nonglaucomatous cases. *Glo-kat* 2007;2:27-30.
10. Sharma KK, Santhoshkumar P. Lens aging: effects of crystallins. *Biochim Biophys Acta*. 2009;1790:1095-108.
11. Moreau KL, King JA. Protein misfolding and aggregation in cataract disease and prospects for prevention. *Trends Mol Med*. 2012;18:273-82.
12. Chowdhury A, Choudhury A, Chakraborty S, et al. p-Benzoquinone-induced aggregation and perturbation of structure and chaperone function of  $\alpha$ -crystallin is a causative factor of cigarette smoke-related cataractogenesis. *Toxicology*. 2018;394:11-18.

- 13.**Berthoud VM, Minogue PJ, Osmolak P, Snabb JI, Beyer EC. Roles and regulation of lens epithelial cell connexins. *FEBS Lett.* 2014;588:1297-303.
- 14.**Wang W, Schaumberg DA, Park SK. Cadmium and lead exposure and risk of cataract surgery in U.S. adults. *Int J Hyg Environ Health.* 2016;219:850-6.
- 15.**Flaye DE, Sullivan KN, Cullinan TR, Silver JH, Whitelocke RA. Cataracts and cigarette smoking. The City Eye Study. *Eye.* 1989;3:379-84
- 16.**Taylor A, Jacques PF, Epstein EM. Relations among aging, antioxidant status, and cataract. *Am J Clin Nutr.* 1995; 62:1439-47.
- 17.**Shalini VK, Luthra M, Srinivas L, et al. Oxidative damage to the eye lens caused by cigarette smoke and fuel smoke condensates. *Indian J Biochem Biophys.* 1994;31:261-6.
- 18.** Ramakrishnan S, Sulochana KN, Selvaraj T, et al. Smoking of beedies and cataract: cadmium and vitamin C in the lens and blood. *Br J Ophthalmol.* 1995;79:202-6.
- 19.**Harding JJ. Cigarettes and cataract: cadmium or lack of vitamin C? *Br J Ophthalmol.* 1995;79:199-200.
- 20.**Rao CM, Robison WG Jr, Zigler JS. Effect of smoke condensate on the physiological integrity and morphology of organ cultured rat lenses. *Curr Eye Res.* 1995;14:295-301.
- 21.**Cumming RG, Mitchell P. Alcohol, smoking, and cataracts; the Blue Mountains Eye Study. *Arch Ophthalmol.* 1997;115:1296-1303.
- 22.**Christen WG, Manson JE, Seddon JM, et al. A prospective study of cigarette smoking and risk of cataract in men. *JAMA.* 1992;268:989-93.
- 23.**Weintraub JM, Willett WC, Rosner B, et al. Smoking cessation and risk of cataract extraction among US women and men. *Am J Epidemiol.* 2002;155:72-9.
- 24.**Klein BE, Klein R, Linton KL, Franke T. Cigarette smoking and lens opacities: the Beaver Dam Eye Study. *Am J Prev Med.* 1993;9:27-30.

25.Lindblad BE, Hakansson N, Wolk A. Smoking cessation and the risk of cataract: a prospective cohort study of cataract extraction among men. JAMA Ophthalmol. 2014;132:253-7.

26.Lindblad BE, Hakansson N, Svensson H, Philipson B, Wolk A. Intensity of smoking and smoking cessation in relation to risk of cataract extraction: a prospective study of women. Am J Epidemiol. 2005;162:73-9.

27.Christen WG, Glynn RJ, Ajani UA, et al. Smoking cessation and risk of age-related cataract in men. JAMA. 2000;284:713-6.

28.Brian G, Taylor H. Cataract blindness challenges for the 21st century. Bull World Health Organ. 2001;79:249-56.

**Table 1: Demographical Features**

<b>Numerical Variables</b>	<b>n</b>	<b>Mean ± SD</b>
<b>Age</b>	581	66.49 ± 9.67
<b>Female - Age</b>	278	66.68 ± 10.68

---

<b>Male - Age</b>	303	66.17 ± 9.37
-------------------	-----	--------------

<b>Categorical Variables</b>	<b>n</b>	<b>%</b>
------------------------------	----------	----------

<b>Gender</b>		
---------------	--	--

<b>Male</b>	303	52.2
-------------	-----	------

<b>Female</b>	278	47.8
---------------	-----	------

<b>HT</b>	239	41.1
-----------	-----	------

<b>DM</b>	173	29.8
-----------	-----	------

---

n; number, SD; Standard Deviation, DM; Diabetes Mellitus, HT; Hypertension

**Table 2:Smoking status and cataract types**

---

	n	%
--	---	---

---

Smoking Load (packages/years)	≤5	13	5
	6-10	24	9.2
	11-20	63	24
	21-30	50	19.1
	31-40	37	14.1
	≥41	75	28.6
Time since quitting smoking (years)	≤1	4	2.5
	1-5	24	15.2
	6-10	46	29.1
	11-15	20	12.7
	≥16	64	40.5
Smoking Status	Never smoked	318	54.7
	Currently Smoking	106	18.2
	Quitted Smoking	157	27.1
Cataract Type	Nuclear	228	39.2
	Cortical	114	19.6
	PSC	239	41.1

n; number, SD; Standard Deviation, PSC; Posterior Subcapsular Cataract.

Table 3: Association between smoking and gender

Gender	Never Smoked	Currently Smoking	Quitted Smoking	p
Female	237(85.3%)	20 (7.2%)	21 (7.6%)	
Male	81 (26.7%)	86(28.4%)	136 (44.9%)	<0.001

**Table 4: The mean ages according to smoking status, and p value.**

Smoking Status	n	Mean ± SD (years)	p
Never Smoked	318	67.82± 9.81	
Currently Smoking	106	61.39 ± 9.36	<0.001
Quitted smoking	157	67.29 ± 8.47	

n; number, SD; Standard Deviation.

**Table 5: The association between cataract type and smoking (Pearson Chi-Square:12.302; p-value:0.015)**

Cataract Type	Never Smoked (n,%)	Currently Smoking (n,%)	Quitted smoking(n,%)	Total (n,%)
Nuclear	119 (37.4)	44 (41.5)	65 (41.4)	228(39.3)
Cortical	77 (24.2)	10 (9.4)	27 (17.2)	114 (19.6)
PSC	122 (38.4)	52 (49.1)	65 (41.4)	239 (41.1)
<b>Total</b>	318 (100)	106 (100)	157 (100)	581 (100)

n; number, PSC; Posterior Subcapsular Cataract.