

Case report

TUBERCULOSIS: MASQUERADE IN OTORHINOLARYNGOLOGY

Abstract

Tuberculosis is a completely curable disease affecting any part of the body. But rare site TB can lead to diagnostic dilemma as it mimics other conditions. This study aims at identifying cases at an early stage preventing morbidity and mortality associated with the same.

KEYWORDS: cholesteatoma, diagnostic error, parotid, radiology of the paranasal sinuses

INTRODUCTION

Tuberculosis (TB) is a bacterial infection caused by *Mycobacterium tuberculosis*. Commonest site of involvement is lung, but it can also affect other parts of the body known as extrapulmonary tuberculosis. However, extrapulmonary tuberculosis poses a diagnostic challenge. TB has surpassed AIDS as the leading cause of death due to infectious disease.¹ Even though the mortality rate is high, it's a completely curable condition with prompt treatment.

Most common site of extrapulmonary TB is lymph node. Our study covers one of its rare forms, the otorhinolaryngeal TB. Our study aims at understanding this diagnostic puzzle and its timely management.

CASE REPORT

We had four cases of rare site TB in the head and neck region.

A 45-year-old female presented with a non-healing ulceroinflammatory lesion over the right medial canthus which was not improving with antibiotics. On anterior rhinoscopy, a fleshy pink bleeding nasal mass was visualised below the right inferior turbinate. Imaging studies revealed a heterogeneously enhancing mass lesion with erosion of adjacent walls. All these investigations narrowed down the differential diagnosis to the possibility of either a lacrimal sac neoplasm or a granulomatous lesion involving the lacrimal sac. Hence to obtain a confirmatory diagnosis, biopsy from representative area was taken. To our surprise, the histopathological assessment was reported as caseating granuloma suggestive of TB. Anti-tubercular therapy

(ATT) was then started and resulted in a healed lacrimal tract fistula (Figure 1) which was later taken for secondary suturing.

Next was a 38-year-old male patient who had slow growing painless swelling in the left infraauricular region. An ultrasonography (USG) was advised initially which reported as suspected ruptured dental abscess. Fine needle aspiration cytology (FNAC) showed inflammatory pathology, acid fast bacilli stain was negative. To assess the spread a CT scan was done which had features suggestive of parapharyngeal neoplasm with submandibular, parotid glands, mandible and left petrous temporal bone involvement with necrotic lymph node (Figure 2). Here our differentials were parapharyngeal neoplasm and osteomyelitis of mandible. In view of those necrotic lymph nodes, a biopsy was advised. USG guided biopsy was performed and sent for HPR which came as necrotising granulomatous lymphadenitis possibly of tubercular aetiology, but special stains for TB were negative. Still, we went on with TB management and the patient started showing recovery in 3 weeks of commencement of ATT.

Third case was that of a 14-year-old female who presented with gradual onset diplopia on right lateral gaze. Clinical examination showed right lateral rectus palsy, rest everything was normal. Radiological imaging of paranasal sinus revealed isolated sphenoid sinus disease eroding the roof of sphenoid (Figure 3) with contrast enhancement in parasellar cavernous sinus. Differential diagnosis of sellar neoplasm, osteomyelitis of sphenoid or allergic fungal rhinosinusitis were taken into consideration. She was posted for surgical clearance of the sinus. Intraoperatively we found granulation tissue (Figure 4) over the sinus mucosa and sample was taken for histopathology reporting. The report was conclusive of multidrug resistant TB on gene xpert. Visual complaints reversed within 1 month of starting ATT.

Another such case was of a 50-year-old female presented with left sided lower motor neuron (LMN) facial palsy and left sided gradual hearing loss associated with vertigo for 8 months. High resolution CT of temporal bone and MRI revealed contrast enhanced erosive lesion involving the left petrous apex (Figure 5). So, we made a differential list of temporal bone neoplasm, osteomyelitis involving the petrous apex and petrous apex cholesteatoma. Since her hearing in left side was non salvageable, we went on with trans cochlear approach and tissue sample were sent for HPR and the result came as TB.

DISCUSSION

Tuberculosis is an endemic disease in developing countries and now there is resurgence of TB in developed countries due to the increased global incidence of AIDS.² Despite lungs being the primary site of involvement, extrapulmonary TB is the major clinical problem as it poses a diagnostic challenge. The presentations are often

misleading camouflaging other conditions.³ It accounts for around 20% of overall active tuberculosis.⁴ Rare forms of extrapulmonary TB include cutaneous, breast, otorhinolaryngeal and disseminated TB.

In our study, first case was an ulceroproliferative lesion of the lacrimal tract failed with conventional treatment modalities, second such case was a parapharyngeal swelling involving mandibular bone, then we had isolated sphenoid sinusitis with cranial neuropathy not recovering with antibiotics and the last patient had petrous apex inflammation with cranial neuropathies. In all these cases, the initial differential diagnosis was neoplastic lesion considering its spread and erosive bony margins. To confirm the diagnosis, biopsy sample was taken in all cases. Keeping in mind, the rise in incidence of extrapulmonary TB with its multidrug resistant form in hike, the tissue samples were also sent for testing TB and reported positive for the same. Still, there was a significant delay in our diagnosis due to its features on imaging favouring neoplasm with subsequent negative stain (acid fast bacilli, AFB) of the FNAC sample.

TB in these rare sites is less mentioned in literature. Failure of initial management protocol should always be an alerting sign to consider TB as a probable diagnosis, especially in a developing country. Routine testing often shows false negative results, so it is necessary to do a cytological/ biopsy evaluation and GeneXpert study to confirm.⁵ With a clinical suspicion of tuberculosis and a positive histopathological report, anti-tubercular therapy should be commenced even on negative AFB staining.¹ Early initiation of treatment will prevent hematogenous spread and thereby mortality.

CONCLUSION

Although the incidence of tuberculosis is high, unusual site tuberculous infections are rare. Rare site tuberculosis mimics malignancy resulting in a diagnostic dilemma. A high index of suspicion of TB should be there when the conventional treatment fails, especially in a developing country where it is endemic. Biopsy sample should always be sent for gene Xpert in doubt. Caseating necrosis with epithelioid granuloma even in the absence of AFB should be treated as TB. If diagnosed early and is still sensitive to first line drugs, then tuberculosis is a completely curable disease.

REFERENCES

1. Clinical study of tuberculosis in the head and neck region—11 years' experience and a review of the literature. *Emerging Microbes & Infections*. Published 2018. Accessed June 10, 2022. <https://doi.org/10.1038%2Fs41426-017-0008-7>
2. SCASSO F, FERRARI G, DE VINCENTIIS GC, et al. Emerging and re-emerging infectious disease in otorhinolaryngology. *Acta Otorhinolaryngologica Italica*. 2018;38(2, SUPPLEMENT 1):S1-S106. doi:10.14639/0392-100x-suppl.1-38-2018
3. Das S. Head and Neck Tuberculosis: Scenario in a Tertiary Care Hospital of North Eastern India. *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. Published online 2016. doi:10.7860/jcdr/2016/17171.7076
4. Sharma S, Rana AK. ENT manifestations of tuberculosis: an important aspect of ENT practice. *Pan African Medical Journal*. 2020;36. doi:10.11604/pamj.2020.36.295.24823
5. Qian X, Albers AE, Nguyen DTM, et al. Head and neck tuberculosis: Literature review and meta-analysis. *Tuberculosis*. 2019;116:S78-S88. doi:10.1016/j.tube.2019.04.014

Figure 1



Figure 1: Right medial canthus ulceroinflammatory lesion before and after anti-tubercular therapy

Figure 2



Figure 2: Left parapharyngeal soft tissue enhancement with erosion of mandible.

Figure 3



Figure 3: Soft tissue enhancement in sphenoid sinus with erosion of its walls

Figure 4



Figure 4: Pale granulation tissue obtained during surgery.

Figure 5




Figure 5: CT showing soft tissue lesion involving left petrous apex with bony wall erosion.