

“EFFECT OF MULCHING AND DIFFERENT-TILLAGE PRACTICES ON YIELD & YIELD ATTRIBUTING CHARACTERS OF SORGHUM CROP”

Abstract: A field experiment was conducted to study the effect of mulching and tillage practices on yield & yield attributes of sorghum at Regional Agricultural Research Station, Andhra Pradesh during 2020-21 & 2021-22 for three Kharif seasons (2020-21 & 2021-22) at Regional Agricultural Research Station, Nandyala, Andhra Pradesh to study the effect of mulch and different tillage practices on yield & yield attributing characters of sorghum crop. Mulching, Tillage practices, mulch practices and their interaction had showed no significant influence on growth parameters viz., plant height and dry matter production at harvest, days to 50 % population. Whereas, panicle weight, panicle length, grain yield, straw yield and harvest index were significantly higher with zero tillage and with residue mulch. The interaction between tillage and mulch practices was non significant. Maximum gross and net returns were observed with zero tillage with residue mulch practice in grain sorghum. The economic evaluation of mulching and tillage practices in sorghum crop revealed that maximum gross returns (Rs 1,18,659), net returns (Rs 78,579) and B:C ratio (2.96) were obtained with zero tillage practice with mulching practice. The study indicated that tillage and mulching had significant effect on soil fertility and grain yield of sorghum crop.

Key Words: Sorghum-crop, Tillage Systems, Mulching, Soil fertility, & Economics

1. Introduction : The World population is growing population has increased the competition pressing demand for land, water, food and other resources, whilst also raising the demand for food. With rapidly increasing population the demand for food grains is continuously increasing in India. Arid and semi-arid regions, where sorghum production practiced, account for about half of the world's total land area, and could play a vital role in solving the world's food security problems. Sorghum is fifth economically important cereal crop in world after wheat, rice, barley and maize. It is an important feed, food and forage crop. The crop is also used in, which also provides raw material for the production of alcohol, biofuels, starch, fiber, dextrose syrup and many other products (Shahid Mehmood *et al.*, 2014). Sorghum is a staple food for millions of people in semiarid regions of the World as a grain crop can and supports wheat, rice, maize and other grain crops, as for food it is a staple food for millions of people in semiarid regions of the World (Shakoor, 1999). Agricultural production in drylands is limited due to the scarcity of rainfall. Increasing the yield per unit of grain is an effective measure to ensure food security. As agricultural production from drylands is limited due to the scarcity of rainfall, therefore, it is essential to increase crop productivity and yields by optimizing agricultural field management practices. Sorghum is fifth economically important cereal crop in world after wheat,

Comment [U1]: Why? Brief the importance of the problem

Comment [U2]: The season discussed different to season stated in Abstract

Comment [U3]: Briefly state treatments and design of the experiment

Comment [U4]:

Comment [U5]: Not the treatment but present as the factor significantly affect the variable

Comment [U6]: Rearrange the sequence as commented

Comment [U7]: Variable showed no significant difference not known

Comment [U8]: Is that the interaction effect? If so, rewrite as combined with otherwise

Comment [U9]: Not clear, may be rewrite

Comment [U10]: No result regarding soil fertility, how measured???

Comment [U11]:

Comment [U12]:

Comment [U13]: Rewrite with improved expressions.

Comment [U14]: ???

Comment [U15]: Show the problem potential and constraint

Show the production potential of sorghum

~~rice, barley and maize. It is an important feed, food and forage crop, which also provides raw material for the production of alcohol, biofuels, starch, fiber, dextrose syrup and many other products (ShahidMehmood *et al.*, 2014). With rapidly increasing population the demand for food grains is continuously increasing in India. Sorghum as a grain crop can support wheat, rice, maize and other grain crops, as it is a staple food for millions of people in semiarid regions of the World (Shakoor, 1999).~~

~~Among different agricultural measures, practices such as mulching, tillage, and their combination are considered as most sustainable agricultural practices in arid and semi-arid regions as they in retain ing soil water in the upper soil layers and reducing the need for irrigation. Compared with traditional soil tillage measures, the implementation of mulching and reduced tillage techniques can significantly reduce soil surface water evaporation, surface ~~runoff~~ runoff, and soil ~~erosion~~ and erosion and increase soil water storage. To an extent, it increases crop yield and water use efficiency and guarantees the sustainable development of ~~agriculture~~ agriculture. Straw mulching has been widely used for the cultivation of maize (*Zea mays* L.), spring wheat (*Triticum aestivum* L.), rice (*Oryza sativa* L.), potatoes (*Solanum tuberosum* L.), vegetables and fruit trees. ~~This farming method not only guarantees the production of crops but is also extremely friendly to the environment.~~~~

~~To attain this probability, soil erosion will have to be condensed. Mulches are loose coverings or sheet of organic material that is placed on the soil surface. It helps to preserve moisture, repression of weeds, improving soil consistency, insect pest assault and guard roots from severe temperature. Organic mulches improve soil, pleasant soil temperature, hinder weed growth, lessen soil moisture evaporation and improve the visual qualities of landscapes. A good layer of mulch will help to preserve moisture and suppress weed germination. Mulch enhanced root and increased maize grain yield by increasing plant N-uptake efficiency, falling N discharge~~

Comment [U16]:

Comment [U17]: Support with literatures

Comment [U18]: Replace with literatures

losses and improving nutrient preservation over unmulched plots (Aulakhet *al.*, 2000). Straw mulch is practiced successfully in many advanced countries like America and Australia where it improved many soil aspect as support soil moisture retention ability, prevent wind erosion, control of weeds, nutrient return and soil structure improvement.

Comment [U19]: Search literatures from India

2. MATERIALS AND METHODS

Comment [U20]: Where is the objectives? Why this study designed?

The experiment was conducted during the Rabi season of 2020-2021 and 2021-2022 at Regional Agricultural Research Station Nandyal. This Theentrecenter is located Searee Rainfall zone at 15.47° N latitude and 78.48° E longitude. The experiment was conducted in an area of 90 m² plot size with 45 cm × 15 cm spacing and fertilizer application of 80-60-40 NPK kg/ha. The total 24 treatments were plotted by using strip plot design consisted of three horizontal practices, T₁ or CT : Conventional tillage (primary, secondary and tertiary tillage as per existing farmers/ researchers practice and sowing by seed cum ferti-drill), T₂ or ZT : Zero tillage (Sowing by seed cum-ferti-drill / cultivator; glyphosate spray to kill existing vegetation after sowing), T₃ or MT : Minimum (Reduced) tillage (Basal fertilizer dose application followed by Disking and sowing by seed drill after running cultivator that incorporates basal fertilizers) and two vertical practices (residue mulch-M1 and without residue mulch-M2). All the six treatments were replicated by 4 times. Strip plot design was statistically analysed by using OPISTAT Software and NTJ 5 variety was selected for the experiment. The parameters like plant height, dry mass production, 100 seed weight, grain yield, straw yield & harvest index were recorded by using standard standard procedures. Initial soil samples were collected from each plot every year from 0 to 30 cm deep. The samples were passed through a 2 mm sievesubjected for analysis of available N, P, and K. For organic carbon analysis, samples were passed through a 0.2-mm sieve. The organic carbon was measured by wet oxidation (Walkley and Black 1934); available N by

Comment [U21]: Not well organized: Need rewriting: State where the experiment conducted, show the clear geographical location. Show the factors to be tested, how these factors were laid out. Show the plot size and area considered for observation. Show the data to be collected, soil sampled collected how analyzed. Finally, the anova procedure used and how Isd tested.

Comment [U22]: Is that Rabi of Kharifseason ??

Comment [U23]: Stated the location before the years

Comment [U24]: Present next to treatments description and layout and area allotted for each treatment

Comment [U25]: Present the treatments, design ahead.

Comment [U26]: Not clear, Re write: how many mulching practices are there, how many tillage practices are they? How these treatments were combined.

Comment [U27]: How?

Comment [U28]: State the data observed not the design subjected to analysis

Comment [U29]: What is the test crop name??

alkaline– potassium permanganate ($KMnO_4$) (Subbaiah and Asija 1956); available P by 0.5 M sodium bicarbonate ($NaHCO_3$) (Olsen *et al.* 1954); and available K using neutral normal ammonium acetate method (Jackson 1973).

3. RESULTS & DISCUSSION

3.1 Growth, yield attributes and yields under sorghum crop

Data present in Table 1 shows that plant height, dry mass production and test weight not significantly affected due to.... there is no significant difference between any treatment for yield attributes like plant height, dry mass production and test weight. Highest plant height (317cm) was observed in CTM2 treatment. Lowest drymass production was observed in CTM2 treatment (8278 kg) may be due to lower nitrogen fertilization than recommended dose of fertilization. These results were in accordance with the findings of Prasad Rao *et al.* (2013).

Mulching and Tillage practices, mulch practices and their interaction had significant influence on yield parameters of sorghum like panicle weight, panicle length, grain yield, straw yield and harvest index. The interaction between tillage and mulch practices was non-significant. Yield attributes and yields of *rabi* sorghum as influenced by tillage and mulch as shown in Table 1. The highest pooled grain yield and straw yield was obtained at zero tillage and lowest pooled grain yield and straw yield at conventional tillage because of zero tillage was attributed to improvement in yield components namely, number of panicle weight and 1000 grain weight. In addition to mulching practices, the highest pooled grain yield and straw yield (3447 kg/ha) (8605 kg/ha) was obtained at zero tillage system compare to other tillage, because of mulching practices reduces the evaporation of water and increases the water holding capacity which results in higher yields. Although, harvest index was not affected by mulch rate, tillage system and their

Comment [U30]: Rewrite: write the soil parameters to be tested and procedure adopted in brief with connecting words

Comment [U31]: ANOVA of the variables not clearly presented and discussed. Language of expression need to be edited and the flow of statement should be improved to standard

Comment [U32]: Correct the tense

Comment [U33]: The factors tested

Comment [U34]: Bring the variable tested to subject line

Comment [U35]: Unknown abbreviations

Comment [U36]: For no significant difference among the treatments we are not supposed to discuss the result. Otherwise, check the data

Comment [U37]:

Comment [U38]: ???

Comment [U39]: Write directly to the variable affected and write the result in brief

Comment [U40]: Show the result Table

Comment [U41]: The statement is against what said in first line in this paragraph

Comment [U42]: ???

Comment [U43]:

Comment [U44]: Present before the result discussion

Comment [U45]: Is this number or weight?

Comment [U46]: The result not presented where Show as zero tillage affected statistically those variable ahead

Comment [U47]: For which does it belong???

Comment [U48]: How does it known?? Writ in probability terms of expression... this suggests or the result indicated that...

interaction (X. Wang and Dai, 2012) also obtained highest harvest index value on the zero-tillage system. This was due to mechanical modifications of soil profiles in Conventional Tillage system which could alleviate high subsoil strength, facilitating deeper rooting and, thus, the plant-availability of subsoil resources (Patil *et al.*, 2016, Schneider *et al.* 2017 & Mishra *et al.*, 2019). Higher soil water availability in Conventional Tillage system compared to Reduced Tillage and No Tillage from sowing to harvest might be responsible for higher leaf area index, better crop growth, and dry matter accumulation and translocation to head at reproductive stage of crop growth producing higher sorghum grain yield (Patil 2013). Deep tillage has been reported to enhance earthworm activities—~~and activities and~~ increase the abundance of plant growth-promoting rhizobacteria and mycorrhizae in the ~~subsoil-subsoil~~. Similar ~~were the findings were~~ reported by Patil (2013), according to him the grain yield of sorghum was lowest in first year of experiment where minimum tillage operation was carried out as against the conventional tillage. In contradictory to this Agbede and Ojeniyi (2009) recorded higher sorghum yields in zero tillage against the conventional tillage. Similarly, Sen *et al.* (2002) also advocated the yield advantage of zero tillage against the conventional tillage. On the other hand, results depicted that mulch had affected the grain yield significantly are in line with Agbede and Ojeniyi (2009) and Shahid Mehmood *et al.*, (2014), that sorghum grain yield was significantly higher in mulched sorghum than that in non mulched (control).

3.2 Soil available nutrient status

Initial and final soil available nitrogen, phosphorous and potassium nutrients ~~status~~ ~~was were~~ estimated in ~~ra~~ sorghum crop and presented in ~~table~~ Table 2. There ~~is~~ no significant difference between ~~any treatment~~ for soil pH, E.C and available potassium. Soil available nitrogen content was significantly highest (168 kg ha⁻¹), with Zero tillage with mulching

Comment [U49]: Present the result of anova before conclusion
Rewrite in new line

Comment [U50]: ??? use last name only for citing

Comment [U51]: Not clear and unknown discussion

Comment [U52]:

Comment [U53]: There is not association of the variables ANOVA result and the supported literatures.

Show each variables tests of significance and associate the discussion and literature at the same paragraph

Comment [U54]: Remove either word

Comment [U55]: The same

Show the result at the start of the result section

Comment [U56]: The result and discussion didn't follow the scientific report writing standard. Language of expression must be improved

Comment [U57]: Is it for each replicated treatments? If so indicate in material and methods section too and

Comment [U58]:

Comment [U59]: Improve the tense

Comment [U60]: Show the treatments

treatment followed by Zero tillage without residue mulching treatment after two years of experimentation. However after two years of experimentation soil available nitrogen content was maintained without depletion. However soil available nitrogen content was significantly decreased due to application of 50% STCR fertilizer dose. Soil available phosphorous content values were significantly differing with ~~tillage—mulching~~ and ~~mulching—tillage~~ practices. Significantly highest available phosphorus recorded for zero tillage (43.6 kg ha^{-1}) and minimum tillage (40.8 kg ha^{-1}) was statistically similar. ~~Mulching had showed~~ significant effect on the soil available phosphorus. Relatively, greater soil available phosphorus was recorded for zero tillage as compared to conventional tillage and minimum tillage. Similar trend was followed in soil available potassium content as in available phosphorous content. But significantly higher values of available potassium were recorded with minimum tillage treatment followed by zero tillage treatment. Relatively, greater soil available potassium was recorded for zero tillage (438 kg ha^{-1}) as compared to conventional tillage and minimum tillage. Maximum increase of as compared to among mulching practices was recorded for poultry manure in all tillage treatments. These results are in agreement with Agbede and Ojeniyi (2009), ~~Shahid Mehmood~~ *et al.* (2014) and ~~Patilet~~ *al.* (2016) who stated that in zero tillage and mulched plots, available potassium was higher compared to minimum and conventional tillage and unmulched plots. Similar results were noticed in secondary nutrients like calcium, magnesium and sulphur availability during two years of sorghum crop growth in soil.

3.3 Economics

The economic evaluation of mulching and tillage practices in sorghum crop revealed that maximum gross returns (Rs 1,18,659), net returns (Rs 78,579) and B:C ratio (2.96) were obtained with zero tillage and mulching practices. The cost incurred on mulching application and

Comment [U61]: Not clear

Comment [U62]: What does it mean?

Comment [U63]: ANOVA result should be presented ahead of discussion

Comment [U64]: Why? discuss

Comment [U65]: May b combined/ merged

Comment [U66]: What??? Define the variable

Comment [U67]:

Comment [U68]: Use last name

Comment [U69]:

Comment [U70]: Not indicated in material methods section

conventional tillage practices reduced the net returns and B: C ratio (1.75) in CTM1 treatment. But the treatment CTM2 recorded lowest B: C ratio (1.64). Suitable tillage practices reduce cost of cultivation; increase income, reduce hazardous effects and increase water use efficiency of sorghum that stabilizes and/or improves the food production in the region. Similar economic benefits ~~have been~~ were reported by Thavaprakash and Malligawad (2002) in sunflower and Schlegelet *et al.*, 2017 in sorghum crop.

Comment [U71]: ???

Comment [U72]: Different crop

4. CONCLUSION

The effect of ~~different mulching and~~ tillage practices ~~and with mulch & without mulch~~ on sorghum crop was conducted at Regional Agricultural Research Station, Nandyal. ~~From this study it is concluded—observed~~ that ~~mulching and~~ tillage ~~practices and mulching had~~ ~~showed~~ significant effect on soil ~~fertility—nutrient content~~ and grain yield of sorghum. ~~The results were observed with~~ ~~It is concluded that~~ highest grain and straw yield was obtained at zero tillage ~~system—practice~~ compare to conventional and minimum tillage ~~system practice~~. The highest ~~harvest index~~ and benefit cost ratio was observed in zero tillage system compared to other treatments.

Comment [U73]: Make similar manner of writing

Comment [U74]: Where is the result and discussion??

REFERENCES:

1. Agbede, Taiwo Michael & Ojeniyi, S.O. (2009). Tillage and poultry manure effects on soil fertility and sorghum yield in southwestern Nigeria. *Soil and Tillage Research*. 104: 74-81. 10.1016/j.still.2008.12.014.
2. Aulakh S.M., T.S. Khera, J.W. Doran, K. Singh, B. Singh, 2000 - Yields and nitrogen dynamics in a rice-wheat system using green manure and inorganic fertilizer. *Soil Science. Society of American Journal*. 64:1867-1876.

Comment [U75]: The style of referencing violates the rule or not according to the authors' guide - The punctuation not properly applied

Comment [U76]: Author name (s), year, title of the article, publisher, volume and page number not properly presented for the reference lists

Rewrite

Comment [U77]:

Comment [U78]: Missed/ not used

Comment [U79]:

Comment [U80]:

3. GebreyesusBrhaneTeshfahunegn,2012,EffectofTillageandFertilizerPracticesonSorghumProductioninAbergelleArea,NorthernEthiopia.4(2):52-69.
4. Jackson, M.L. 1973. Soil Chemical Analysis. Prentice Hall of Inco. New York, USA. 498.
5. Mishra, S SRao And I K Das. 2019.Effect of tillage and nutrient management on sorghum (Sorghum bicolor) productivity in Alfisols of semi-arid tropical India.*Indian Journal of Agricultural Sciences*. 89 (7): 1133–9
6. Olsen SR, Cole CV, Watnabe FS, Dean LA.Estimation of available phosphorus in soil by extracting with sodium bicarbonate. USDA Circular No.989, United state department of agriculture, Washington DC, 1954. 15.
7. PatilS L. 2013. Winter sorghum (Sorghum bicolor) productivity as influenced by tillage practices and nitrogen management in Vertisols of SAT, India. *Soil & Tillage Research*126: 183–92.
8. PatilS L, Mishra P K and Ramesha M N. 2016. Tillage and integrated nitrogen management: Does sustain sorghum productivity in Vertisols of Semi-Arid Tropics under varying rainfall situations? *Indian Journal of Soil Conservation*44 (2): 198–205.
9. Prasada Rao, Subbaiah VG, Chandra Sekhar K. Department of Agronomy, Agricultural College, Bapatla, Acharya N.G Ranga Agricultural University, International Journal of Applied Biology and Pharmaceutical Technology, 2013, 4(4). 18.
10. Schlegel A J, Assefa F, Haag L A, Thompson C R and Stone L R. 2017. Long-term tillage on yield and water use of grain sorghum and winter wheat. *Agronomy Journal* 110: 1–12.
11. Schneider F, Don A, Hennings I, Schmittmann O and Seidel S J. 2017. The effect of deep tillage on crop yield – What do we really know? *Soil & Tillage Research* 174: 193–204.
12. Sen A, Shama N, Singh RK, Pandy M. 2002. Effect of different tillage systems on the performance of wheat.Paper for international workshop on herbicide resistance management and zero tillage in rice- wheat cropping system.March 4-6, Department of Agronomy, CCS Haryana Agriculture University, Hisar-125 004, India.
13. ShahidMehmood ,ShahidIbniZamir , Tassadduq Rasool and Waseem Akbar. 2014. Effect of Tillage and Mulching on Soil Fertility and Grain Yield of Sorghum.*ScientiaAgriculturae*.4(1), 2014: 31-36. DOI: 10.15192/PSCP.SA.2014.4.1.3136.
14. Shakoor A. 1999. Evaluation of exotic and local cultivars of sorghum for yield, maturity and nonscesence associated characters under rain-fed conditions. *Pakistan Journal of Biology Sciences*.13: 167-174.
15. SubbiahBV, Asija GL.A rapid procedure for estimation of available nitrogen in soil. *Current Science*. 1956; 25:259-260. 14.

Comment [U81]:

Comment [U82]:

Comment [U83]: Year of publication

Comment [U84]:

Comment [U85]:

Comment [U86]:

Comment [U87]:

Comment [U88]:

Comment [U89]:

Comment [U90]: Does the first name need space, or separated with dot,? Not uniform

Comment [U91]: ???yr

Comment [U92]:

Comment [U93]: missed

16. Thavaprakash N, and Malligawad LH. 2002. Effect of nitrogen and phosphorus levels and ratios on yield and economics of sunflower. *Research on Crops*; 3:40-43.
17. Walkley, A. and Black, C.A. 1934. Estimation of organic carbon by chromic acid titration method. *Soil Science*. 37: 29-38.
18. X. Wang, H. Wu, K. Dai., 2012, Tillage and crop residue effects on rainfed wheat and maize production in northern China. *Field Crops Research*, 132: 106–116.

Comment [U94]: Which one is the last name

Table 1. Yield & Yield attributing characters of Sorghum crop as influenced by tillage and mulch (Mean data of 4 replications)

	Plant Height (cm)	Dry Mass production (Kg)	Days to 50% population	Test weight (g)	Grain yield (Kg/ha)	Stover yield (Kg/ha)	Harvest Index
CTM ₁	310	8650	75.75	32.55	3171	8379	27.46
CTM ₂	317	8278	76.00	31.78	2838	8004	26.18
ZTM ₁	312	8835	74.75	35.63	3447	8605	28.61
ZTM ₂	311	8692	76.25	31.30	3267	8353	28.09
MTM ₁	303	8800	75.50	33.88	3365	8568	28.19
MTM ₂	306	8554	75.75	32.18	3205	8333	27.77
SEm _±	28	223	7.3	3.04	201	257	
C.D(P=0.05)	NS	685	NS	NS	547	783	
C.V (%)	8.4	10.1	8.7	6.3	12.3	12.7	
Interaction(Mulch xTillage)							
SEm _±	23.67	210.50	7.25	3.12	204	236	
CD(P=0.05)	NS	NS	NS	NS	NS	NS	
Interaction(Tillage xMulch)							
SEm _±	31.44	215.60	8.32	4.32	191	257	

CD(P=0.05)	NS	NS	NS	NS	NS	NS	NS	
------------	----	----	----	----	----	----	----	--

Table 2: Influence of Tillage & mulching on soil properties of sorghum crop

Treatment name	pH	E.C (d Sm ⁻¹)	O.C (%)	Available N (Kg/ha)	Available P ₂ O ₅ (Kg/ha)	Available K ₂ O (Kg/ha)	Ca (C mol (p+) kg-1)	Mg (C mol (p+) kg-1)	Sulphur (ppm)
CTM ₁	8.31	0.10	0.30	146	41.2	398	9.0	3.2	13.1
CTM ₂	8.33	0.14	0.30	142	40.2	404	8.8	3.2	12.8
ZTM ₁	8.22	0.08	0.33	168	43.6	422	9.7	4.6	14.1
ZTM ₂	8.30	0.14	0.33	165	42.9	434	9.5	4.3	14.4
MTM ₁	8.28	0.12	0.30	154	39.3	438	9.2	4.5	15.2
MTM ₂	8.30	0.17	0.30	158	40.8	447	9.4	3.9	15.1
S. Em ±	0.45	0.01	0.02	9.04	2.80	24.47	0.51	0.26	0.81
C. D (P=0.05)	NS	NS	0.057	26.23	7.10	NS	NS	0.79	NS
C V (%)	6.8	4.5	8.87	9.32	7.88	12.1	9.82	8.4	7.4

Interaction(Mulch xTillage)									
SEm±	0.42	0.01	0.01	7.88	2.04	23.61	0.34	0.17	0.74
CD(P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interaction(Tillage xMulch)									
SEm±	0.38	0.01	8.32	8.04	2.01	22.83	8.57	0.16	0.71
CD(P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS
Initial	8.32	0.06	0.33	135	43.05	383	8.2	2.4	12.2

Table 3. Economics of sorghum crops influenced by tillage and mulch practices

Treatment name	Grain yield (Kg/ha)	Stover yield (Kg/ha)	Gross returns (Rs/ha)	Net returns (Rs/ha)	Benefit :cost ratio (B:C ratio)
CTM ₁	3171	8379	97411	37623	1.63
CTM ₂	2838	8004	104663	44875	1.75
ZTM ₁	3447	8605	118659	78579	2.96
ZTM ₂	3267	8353	107714	67634	2.69
MTM ₁	3365	8568	104071	62036	2.48
MTM ₂	3205	8333	113451	71416	2.70