

Original Research Article

Influence of Irrigation Scheduling on Weeds among Different Sowing Dates in Wheat under Vertisol

ABSTRACT

Wheat is an important cereal grain crop and is staple to millions. Weeds are the major constraint that lower the wheat yield. The knowledge of weeds under different sowing dates and Irrigation can assist in controlling weeds. A field experiment was conducted during the *rabi* season of 2020-21 at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, M.P. to study the influence of irrigation scheduling on weeds at different sowing dates in wheat. The experiment was laid out in split-plot design with three replications. The main plot consisted of three sowing dates, i.e., 3rd December, 18th December and 2nd January and four Irrigation Water/Cumulative Pan Evaporation (IW/CPE) based irrigation scheduling, 1.0, 0.9, 0.8 and 0.7 in the sub-plots. The results revealed that lowest weed density and dry weight were observed in 2nd January sown date, as compared to 3rd December and 18th December sown date. Among the irrigation scheduling, 0.7 IW/CPE observed lowest weed density and dry weight than 1.0, 0.9, 0.8 IW/CPE ratio. 3rd December sown date exhibited highest grain yield (4637 kg ha⁻¹) and straw yield (6788 kg ha⁻¹) than 18th December and 2nd January sown date. Among the irrigation schedule, IW/CPE ratio 1.0 exhibited maximum grain yield (4510 kg ha⁻¹) than 0.9, 0.8 and 0.7 IW/CPE ratio. The results of the study concludes that sowing dates and irrigation schedules had crucial role for controlling weeds.

Keywords: IW/CPE ratio, sowing date, Weed density and dry weight

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the major cereal grain cultivated over a large area in the world and consumed as major source of carbohydrate in the human nutrition. It is sown from mid-November to the end of December in the Central India. In recent years, the sowing gets delayed due to the addition of garden pea in the cropping system, as a catch crop after harvesting of *kharif* crops. This delay affects the sowing as it is vulnerable to temperate conditions. Late sowing may lead to reduce crop growth duration as phenological stages were not compatible with late weather conditions. Delayed sowing results in decrease in number of days to attain different growth stages in wheat, [Khande et al.](#) [1] and this delay in sowing affects the grain yield of wheat [2]. It also affects the weed density thereby competitiveness with the main crops. Weeds rivalry for reducing the yield of crop as they contend with crop for moisture, light and nutrient. Water management and weed management are the crucial factors affecting the yield of wheat [3]. Uncontrolled weed growth may reduce wheat yield ranging from 15-40 % depending upon magnitude, nature, and duration of weed infestation [4]. [Jain and Agarwal](#) [5] observed presence of weeds causes reduction in yield. Moreover, sowing date has pronounced effect on weed density

and biomass [6]. The role of irrigation is crucial for sowing window as it may cope up the effect of warm weather conditions for late maturing crop. Chouhan et al. [7] observed more yield of wheat at 1.0 IW/CPE ratio. Therefore, different irrigation scheduling based on Irrigation Water: Cumulative Pan Evaporation (IW/CPE) ratio for wheat under different thermal environments is studied. Irrigation scheduling based on climatological approach, IW/CPE ratio method, in which a known amount of water is applied when cumulative pan evaporation reaches a predetermined level. It enables to apply exact amount of water thereby increased the irrigation efficiency. It is anticipated that the impact of late sowing can be alleviated through irrigation regimes upto some extent. However, the role of weeds under thermal environments and irrigation scheduling needs to be assessed out. In view of the above facts, a study was carried out with an objective to assess weed density and weed dry weight under different sowing dates and IW/CPE levels of irrigation scheduling in wheat grown at Central Indian conditions.

2. MATERIAL AND METHODS

The research study was conducted during the *rabi* of 2020-21 at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur (M.P.) situated at 23°09' N latitude and 79°58' E longitude and an altitude of 411 above mean sea level. According to U.S.D.A. classification, the soil of this region is broadly classified as "Vertisol". The experimental soil consists of 30 % sand, 20 % silt and 46 % clay with organic matter content of 0.94 % and 7.1 pH. The total rainfall received was 22.0 mm distributed in two rainy days with total sunshine hours of 132.2 hours during whole crop season. The field experiment was laid out in split-plot design with three date of sowing (December 03, December 18 and January 02) as main treatment and IW/CPE ratio (1.0, 0.9, 0.8 and 0.7) as the sub-plot treatment in three replications. Wheat variety, MP 3336 tolerant to heat and suitable for late sowing under irrigated conditions for Madhya Pradesh was selected for the experiment. Irrigation was given prior to land preparation and all the management practices were done as per the prescribed package of practices recommended for this region. For irrigation scheduling, a uniform irrigation was applied immediately after sowing to all the treatments for better germination of the crop. After then, water was applied as per the irrigation schedule based on climatological approach to the experimental plots. The depth of irrigation was fixed as 50 mm for four IW/CPE ratio i.e., 1.0, 0.9, 0.8 and 0.7 as per the formula by Parihar et al. [8]

$$IW/CPE = \frac{\text{Depth of irrigation (mm)}}{\text{Cumulative pan evaporation (mm)}}$$

There is a difference in the number of irrigations between different IW/CPE treatment. Total weed density and weed dry weight were observed at 30 days after sowing (DAS) but before applying of herbicide; 30 and 60 days after herbicide application (DAHA); and at harvest, using a 0.5 m² quadrat placed two times per plot. Before analysis of data on weed density and dry weight, it was subjected to square root transformation ($\sqrt{x+0.5}$). The total weed density and dry weight were expressed in numbers (no.) m⁻² and g m⁻², respectively. The herbicide, Clodinafop-propargyl 15 % + Metsulfuron methyl 1%, was applied at 60+4 g/ha at 30 days after sowing wheat. The efficiency to control weeds were analyzed by differentiating between weeds observed in 1.0 IW/CPE (control) and weeds observed in the other 0.9, 0.8 and 0.7 IW/CPE irrigation approaches. The field data obtained was statistically analyzed using OPSTAT software available online at CCS Haryana Agriculture University [9]. The data were tabulated and analyzed statistically by using ANOVA as suggested by Gomez and Gomez [10] and significant difference between treatment means were compared with critical difference at 5 % levels of probability for F- test.

3. RESULTS AND DISCUSSION

3.1 Weed Flora

The wheat crop was majorly infested with divergent types of weed flora as *Medicago denticulata* L., *Chenopodium album* L., *Anagallis arvensis* L., *Alternanthera sessilis* L. The other weed species were *Cyperus rotundus* and *Echinochloa colona* in very low density among all the treatments.

Medicago denticulata L., *Chenopodium album* L., *Anagallis arvensis* L., *Alternanthera sessilis* L. were observed among the broad leaf weeds, while *Cyperus rotundus* among the sedges and *Echinochloa colona* among the grassy weeds were also observed.

3.2 Weed Density (No. m⁻²)

Data pertaining to weed density were presented in **Table 1**. Among the sowing dates, maximum total weed density (4.36) was observed in 3rd December sowing date, however, it was at par with 18th December (4.32) sowing date before herbicide application. At 30 DAHA, total weed density was significantly higher in 3rd December (3.68) followed by 2nd January (3.52) and 18th December (3.50) sowing date. At 60 DAHA, significantly maximum weed density was observed at 3rd December (2.40) sown date, however, it was at par with 18th December (2.36) sown date. At harvest, it was maximum at 3rd December (2.37) sown date as compared to 18th December and 2nd January. This might be because of varying temperature received at different timings of sown date. High temperature received with the delay in sowing might have affected the germination of weed seeds. If germinate, then it is difficult to establish due to non-favorable weather environment for weeds. Furthermore, reduction in weed density might be because of shift in weed flora in delayed sowing. Similar finding has been reported by **Thorat et al.**[5] and **Jain**[11] that delay in sowing date have profound effect on weed density and biomass.

Among the irrigation schedule, maximum weed density was observed at 1.0 IW/CPE ratio (4.42) followed by other irrigation schedules before herbicide application (**Table 1**). At 30 DAHA, maximum weed density was observed in 1.0 IW/CPE ratio (3.72) that was at par with 0.9 schedule while minimum weed density exhibited in 0.7 (0.35). At 60 DAHA, similar results were observed except, 0.9 and 0.8 schedules are non-significant in the population of total weeds. At harvest, maximum weed density observed at 1.0 while in 0.8 and 0.7 IW/CPE ratio. The maximum weed density in 1.0 IW/CPE ratio might be due to sufficient moisture availability of frequent irrigations than 0.8 and 0.7 IW/CPE approach and later on due to congenial weather conditions for good germination and growth of weeds that helped to establish easily. **Nadeem**[12] reported increase in weed population with the increase in irrigation frequency from 0.6 to 1.0 IW/CPE ratio.

3.3 Weed dry weight (g m⁻²)

The data regarding weed dry weight at before herbicide application, 30 DAHA, 60 DAHA and at harvest is presented in **Table 1**. It was observed that weed dry weight was significantly higher at 3rd December and 18th December sown date than 2nd January before herbicide application. At 30 DAHA, it was significantly maximum at 3rd December and 18th December sown date than 2nd January. The similar results were also observed at 60 DAHA and harvest. Among the irrigation schedules, 1.0 IW/CPE recorded significantly higher weed dry weight followed by 0.9 and 0.8, and minimum in 0.7 irrigation schedules. Similar results were also observed at 30 and 60 DAHA. At harvest, 1.0 IW/CPE ratio observed significantly higher weed dry weight at 1.0 than the other IW/CPE ratio. The interaction was found to be non-significant between sowing date and irrigation schedules. This suggested that the total weed dry weight exhibited more at higher irrigation applied than with less irrigation applied in wheat. The results are in confirmatory with **Verma et al.**[13] that dry weight of weeds significantly affected by irrigation regimes and weed management practices. **Thorat et al.** [6] and **Jain** [11] also observed increase in weed biomass with the increase in irrigation frequency.

3.4 Efficiency to control weeds

The efficiency to control weeds were analyzed between 1.0 IW/CPE as control and other treatments at different sown dates, and is presented in the form of interaction in **Table 2**. The result exhibited that sown dates were not significantly different. However, irrigation schedule observed significant different at 30 and 60 DAHA, and harvest. It exhibited that less irrigated schedule (0.7) controls more weeds than the other 0.8 and 0.9 IW/CPE, respectively. This suggested that irrigation schedule is proportional to weed occurrences that

were also suggested by Thorat et al.[6] exhibited lowest weed density and dry weight at low frequency of irrigation. Jain[11] also observed effect of irrigation schedule on weed biomass. The interaction between sown date and irrigation schedule observed at the initial stage of crop growth at 30 DAHA, while it was found to be non-significant at later stages of crop growth.

UNDER PEER REVIEW

Table 1. Influence of IW/CPE based irrigation scheduling on weed density and weed dry weight among different sowing dates in wheat under Vertisol

Treatments	Density (No. m ⁻²)	Dry weight (g)	Density (No. m ⁻²)	Dry weight (g)	Density (No. m ⁻²)	Dry weight (g)	Density (No. m ⁻²)	Dry weight (g)
	(Before herbicide application)		(30 days after herbicide application)		(60 days after herbicide application)		(Harvest)	
Sowing dates								
December 03	4.36 (59.33)	1.81 (4.79)	3.68 (29.08)	2.19 (13.57)	2.40 (22.50)	2.08 (10.56)	2.37 (21.00)	1.94 (7.23)
December 18	4.32 (57.83)	1.78 (4.26)	3.50 (22.75)	2.16 (12.98)	2.36 (20.50)	2.05 (9.83)	2.29 (17.75)	1.89 (6.13)
January 02	3.99 (41.08)	1.72 (3.37)	3.52 (21.33)	2.09 (11.09)	2.30 (18.08)	2.00 (8.54)	2.21 (14.58)	1.82 (4.98)
SE m±	0.02	0.01	0.02	0.01	0.02	0.01	0.03	0.004
CD (5%)	0.09	0.04	0.08	0.04	0.07	0.05	0.10	0.02
Irrigation levels- IW/CPE ratio								
1.0	4.42 (63.33)	1.83 (5.03)	3.72 (29.89)	2.22 (14.82)	2.47 (25.77)	2.11 (11.36)	2.38 (21.66)	1.94 (7.23)
0.9	4.19 (52.56)	1.78 (4.19)	3.67 (26.22)	2.16 (13.04)	2.38 (21.33)	2.06 (10.09)	2.31 (18.33)	1.91 (6.61)
0.8	4.22 (50.22)	1.76 (3.88)	3.53 (22.33)	2.14 (12.22)	2.32 (18.77)	2.03 (9.30)	2.25 (16.33)	1.86 (5.60)
0.7	4.06 (44.89)	1.72 (3.47)	3.35 (19.11)	2.06 (10.1)	2.23 (15.55)	1.97 (7.82)	2.21 (14.77)	1.83 (5.00)
SE m±	0.04	0.007	0.04	0.01	0.02	0.006	0.02	0.008
CD (5%)	0.13	0.02	0.11	0.03	0.07	0.02	0.05	0.02

*Values are $\sqrt{x+0.5}$ square root transformed. Values in parenthesis are original

Table 2. Influence of IW/CPE based irrigation scheduling on Efficiency to control weeds among different sowing dates in wheat under Vertisol

Treatments	30 DAHA				60 DAHA				Harvest			
	Dec 3	Dec 18	Jan 2	Mean	Dec 3	Dec 18	Jan 2	Mean	Dec 3	Dec 18	Jan 2	Mean
1.0	-	-	-	-	-	-	-	-	-	-	-	-
0.9	22.3	23.3	34.0	26.5	30.8	25.3	20.9	25.6	26.2	21.0	25.3	24.2
0.8	30.1	32.3	29.3	30.6	34.2	31.5	27.9	31.2	37.1	34.7	30.1	34.0
0.7	45.4	39.5	37.8	40.9	40.7	42.0	38.5	40.4	41.8	39.0	40.7	40.5
Mean	32.6	31.7	33.7		35.2	32.9	29.1		35.0	31.6	32.0	
Comparing means	SE m±	CD (5%)		SE m±	CD (5%)		SE m±	CD (5%)		SE m±	CD (5%)	
D- Sowing Date	2.63	NS		1.33	NS		3.02	NS				
I- Irrigation levels- IW/CPE ratio	1.53	4.77		0.90	2.81		1.28	4.00				
I at same level of D	4.56	9.71		2.30	NS		5.23	NS				
D at same level of I	3.41	12.48		1.84	NS		3.52	NS				


Efficiency was a difference between IW/CPE of 1.0 (control) with the other irrigated treatments of 0.9, 0.8, and 0.7 at different sowing dates

UNDER PEER REVIEW

3.5 Yield

Sowing date 3rd December recorded higher grain yield (4637 kg/ ha) and straw yield (6788 kg/ ha) were recorded at 3rd December sown date than other sown dates. Among the irrigation schedule, IW/CPE ratio 1.0 recorded significantly higher grain yield (4510 kg/ ha) which was statistically at par with 0.9 IW/CPE ratio and straw yield (6677 kg/ ha) as compared to 0.9, 0.8 and 0.7. This might be because of favourable environment and adequate moisture availability that favoured in gaining more yield and hence more profitable. The results of present investigation were in line with Verma et al.[14], Thorat et al.[6], Buttar et al.[15], Jain[11], Prasad et al. [16], Bhaskar et al. [17] and Lanjhewar et al. [18]. Alam et al. [19].

Fig 1. Influence of IW/CPE based irrigation scheduling on grain and straw yield among different sowing dates in wheat under Vertisol



4. CONCLUSION

From the present research, it can be concluded that sowing dates and IW/CPE based irrigation scheduling had significant role in weed control and grain yield. Thus, adjusting sowing dates and scheduling irrigation could be used to reduce weed and enhance grain yield of wheat under late sowing condition.

REFERENCES

1. Khande D, Agrawal KK and Bhan M. Phenology and heat unit requirement of wheat under various thermal environments and different methods of sowing in Tawa Command area of Madhya Pradesh. 2021; Anveshak. 51 (2).
2. Agrawal KK, Shankar U, Upadhyay AP and Gupta VK. Accumulated heat unit requirements for different Phenophases of wheat cultivar as influenced by sowing dates at Jabalpur. Journal of Agrometeorology. 1999. 1(2): 173-176.

3. Nadeem MA, Tanveer A, Ali A, Ayub M and Tahir M. Effect of weed-control practices and irrigation levels on weeds and yield of wheat. *Indian Journal of Agronomy*. 2007; 52(1):60-63.
4. Jat RS, Nepalia V, Chaudhary PD. Influence of herbicides and method of sowing on weed dynamics in wheat (*Triticum aestivum* L.). *Indian Journal of Weed Science*. 2003; 35: 18-20.
5. Jain NK and Agrawal KK. Effect of irrigation levels, (based on IW/CPE ratio) fertility levels and weed control methods in linseed (*Linum usitatissimum* L.). *World Weeds*. 1998; 5: 41-45.
6. Thorat TN, Bhan M and Agrawal KK. Effect of sowing dates on weed infestation and yield of wheat (*Triticum aestivum* L.) varieties under different irrigation schedules. *JNKVV ResJ*. 2015; 49(1):41-45.
7. Chauhan BS, Kaushik MK, Nepalia V, Solanki NS, Singh B, Davra NS, Kumavat P and Kumar. Effect of sowing methods, scheduling of irrigation based on IW/CPE ratio and chemical weed control on plant height, dry matter accumulation and yield of wheat. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6(30): 169-172.
8. Parihar SS, Khera KL, Sandhu KS and Sandhu BS. Comparison of irrigation schedules based on pan evaporation and growth stages in winter wheat. *Indian Journal of Agronomy*. 1976; 68: 650-653.
9. www.hau.ernet.in
10. Gomez KA and Gomez AA. Statistical procedure for Agriculture Research, 2nd Edition, pp 241-71. John Wiley & Sons, New York; 1984.
11. Jain P. Fertilizer and water interaction for predicting wheat yield under varying thermal regimes in vertisol using CERES Wheat model. Ph.D. Thesis Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur; 2019.
12. Nadeem MA, Tanveer A, Ali L, Ayub M and Tahir M. Effect of weed control practices and irrigation levels on weeds and yield of wheat. *Indian Journal of Agronomy*. 2010; 52(1): 60-63.
13. Verma SK, Singh SB, Prasad SK, Meena RN and Meena RS. Influence of irrigation regimes and Weed management practices on water use and weed management in wheat (*Triticum aestivum* L. and Emend. Fiori and Paol). *Bangladesh Journal of Botany*. 2015; 44(33):437-442.
14. Verma BR, Ram S, Sharma B. Effect of soil moisture regimes and fertility levels on growth, yield and water use efficiency of wheat (*Triticum aestivum* L.). *Progressive Agriculture*. 2011; 11(1): 73-78.
15. Buttar PM, Kingra PK, Singh SP. Effect of sowing dates, irrigation and mulching on growth and yield of wheat. *Agric Res J*. 2018; 55(2): 243-250.
16. Prasad S, Agrawal K K, Kumar R. Productivity, Profitability, Quality and nutrient uptake of heat tolerant wheat (*Triticum aestivum*) cultivars as influenced by staggered sowing and nitrogen levels. *Indian Journal of Agriculture Science*. 2019; 89(4): 670-7.
17. Bhaskar GP, Shrivastava GK and Sumit. Effect of irrigation levels and weed management on yield attributes and Yield of wheat (*Triticum aestivum* L.) in North-Hill zone of Chhatisgarh. *The Pharma Innovation*. 2022; 11(11):225-231.
18. Lanjhewar A, Agrawal KK, Bhan M and Patel K. Influence of Thermal Regimes and Irrigation Schedules on Growth, Yield of Wheat (*Triticum aestivum* L.) Varieties. *International Journal of Environment and Climate Change*. 2022; 12(12): 484-490.

19. Alam MS, Naresh RK, Vivek, Kumar S and Singh HL. Effect of sowing methods and irrigation scheduling on production and productivity of wheat crop. 2022. Biological Forum – An International Journal. 14 (2a): 445-452.

UNDER PEER REVIEW