

Effect of cultivar and nutrient management on productivity, and profitability of summer greengram (*Vigna radiata* (L.) Wilczek

ABSTRACT

The present study was carried out at Research Farm of Regional Research Station, Bawal, CCS HAU, Hisar, Haryana during *summer*, 2020. In this study, main plot consisted of three different varieties, namely MH-421 (V₁), MH-318 (V₂) and SML-668 (V₃) and subplot included four nutrient treatments, namely Control (F₁), 100% RDF (F₂), 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB (F₃) and 50% RDF + 50% N/ha (FYM) + ST with *Rhizobium* + PSB (F₄). The result revealed that, greengram variety and nutrient treatment significantly influenced the growth, yield and economics of summer greengram. Among the greengram variety, MH-318 variety yielded significantly higher seed yield, net return and B:C ratio compared to MH-421 and SML-668. Further, application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB resulted in significantly higher yield, and monetary benefits. Therefore, the study suggests that fertilizing MH-318 variety with 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB to obtain higher yield and economic returns of summer greengram.

KEYWORDS: Economics, Greengram, *Rhizobium*, Seed yield, Varieties.

INTRODUCTION

Pulses play a crucial role in our foodgrain system along with cereals to stabilize the demand and fulfil nutritional requirements. India holds the top position as both the primary producer and consumer of pulses worldwide. Nevertheless, the per capita availability of pulses has been gradually declining over the years (John *et al.*, 2021). Greengram (*Vigna radiata* (L.) Wilczek) is the third most important legume crop cultivated in India, followed by chickpea and pigeon pea. It belongs to the family Fabaceae and sub family Papilionaceae and is believed to be native of India and central Asia (Vavilov, 1926). This crop is extensively grown across various regions in Asia, encompassing countries such as India, Pakistan, Sri Lanka, Indonesia, Bangladesh, Cambodia, Malaysia, Thailand, Vietnam, Laos and southern China. Additionally, it is also cultivated in certain areas of Africa.

In 2021-22, area and production of greengram in India is 55.07 lakh ha and 31.66 lakh tonnes with productivity of 570 kg/ha, whereas in Haryana it is 0.2 lakh ha and 0.12 lakh tonnes with productivity of 601 kg/ha (Anonymous, 2022).

Greengram grains typically have a protein content of 23.88%, which is significantly higher compared to wheat (10-11%) and rice (6-7%). Furthermore, it is a valuable source of essential amino acids like tryptophan, lysine and arginine. In addition to its importance in human nutrition, green gram also contributes to enhancing soil fertility through the process of atmospheric nitrogen fixation. It can fix 43-85 kg/ha atmospheric nitrogen annually (Rosales *et al.*, 1995). Due to its short duration, wider adaptability, photo and thermo insensitive, it can adjust in various cropping systems.

Despite significant importance, the yield of greengram is very low in India (Nair *et al.*, 2012). The factors contributing to low productivity include challenges such as susceptibility to pests and diseases, losses caused by shattering, limited availability of high-yielding varieties, reliance on rainfed conditions and cultivation in infertile soils with minimal or no financial inputs (Lal *et al.*, 2015). Among the various factors responsible for maximization of yield, selection of high yielding variety along with balanced nutrition is essential for high yield (Meena *et al.*, 2016). A strategy that combines diversification, enhanced productivity and improved soil characteristics is needed to address the significant gap between the demand and supply of pulses. This three-pronged approach aims to meet the demand and overcome the gap by incorporating measures to broaden the variety of crops grown, increase the yield of existing crops and enhance the quality of soil (Bhardwaj *et al.*, 2022). The yield of any crop is product of production potential of the variety, climatic conditions, soil fertility and management practices. Therefore, breeding methods such as selection, hybridization and mutation have been applied and some additional improvement has also been done during last two decades (Pareek *et al.*, 2022). Similarly, ensuring an adequate supply of nutrients, particularly nitrogen and phosphorus, is essential to prevent any constraints on the growth of greengram. The positive impact of organic manures, including farmyard manure (FYM), on enhancing soil fertility and crop productivity has been extensively documented. Nevertheless, there is limited information available regarding the optimal combination of nutrients from different sources, such as bio-fertilizers, inorganic fertilizers and organic fertilizers, specifically for summer greengram cultivation. Thus, this present study is conducted to evaluate the effect of varieties and nutrient treatments on growth, yield attributes, yield and economics of summer greengram.

MATERIALS AND METHODS

A field experiment was conducted at Research Farm of Regional Research Station, Bawal, CCS Haryana Agricultural University, Rewari, Haryana located at 28.07 °N latitude 76.59 °E longitude and elevation of 266 m above mean sea level during *summer* of 2020. The experimental site had a loamy sand soil texture with a pH of 8.17 and an electrical conductivity (ECe) of 0.16 dS/m. The soil was characterized by low organic carbon content (0.16%) and medium availability of nutrients, with 137.3 kg/ha of available nitrogen, 11.18 kg/ha of available phosphorus and 149.1 kg/ha of available potassium at a soil depth of 0-15 cm. The study consisted twelve treatment, main plot occupied by three varieties, namely 'MH-421', 'MH-318' and 'SML-668', and subplots were allotted to nutrient management practices, namely Control (F₁), 100% RDF (F₂), 75% RDF + 25% FYM N/ha + ST with *Rhizobium* + PSB (F₃) and 50% RDF + 50% FYM N/ha + ST with *Rhizobium* + PSB (F₄). The treatment were arranged in split-plot design with four replications. In this study, each plot had a size of 3 m x 5 m. The crop was sown on April 25, 2020 with a row-to-row spacing of 30 cm with a seed rate of 25 kg ha⁻¹. Before sowing, the seeds were treated with Bavistin at a rate of 3 g/kg to safeguard against seedborne diseases. The application of fertilizers, organic matter in the form of farmyard manure (FYM) and biofertilizers followed as per the treatment. One hoeing-cum-weeding was done manually at 20 DAS. After harvesting the crops from each plot, they were carefully bundled and appropriately labelled. Once the second harvest was completed, the produce was sun-dried on the threshing floor. Statistical analysis of data was carried out by "Analysis of Variance" method (Panse and Sukhatme, 1967), the data were analyzed statistically using ANOVA and treatment comparison was made at 5% level of significance.

RESULTS AND DISCUSSION

Growth Attributes

Table 1 displays plant height (cm) of greengram crop at different growth stages. The data revealed that plant height was significantly influenced by both varieties and nutrient treatment, except at 15 DAS. MH-318 variety exhibited significantly taller plants than SML-668 and MH-421. Application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB resulted in significantly higher plant height at 30, 45 DAS and at harvest, compared to other treatment, except for 100% RDF, where it was at par. The increase in plant height resulting from the application of 75% RDF + 25% FYM + ST with *Rhizobium* + PSB, in comparison

to the control, was observed to be 27.45%, 17.83%, 8.95% and 12.17% at 15, 30, 45 DAS and at harvest, respectively.

Table 1 also presents data on dry matter accumulation, indicating a continuous increase as the crop matures until harvest. Varieties and nutrient treatment showed significant impact on dry matter accumulation at all stages, except at 15 DAS. MH-318 and SML-668 demonstrated significantly higher dry matter accumulation than MH-421. The maximum dry matter accumulation was observed with 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB at all growth stages, showing significant differences at all growth stages except 100% RDF, where it was at par. The application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB outperformed other treatments at 30, 45 DAS and at harvest. The combined utilization of inorganic fertilizers, organic manures and seed treatment may have facilitated significant dry matter accumulation by promoting the production of growth-enhancing hormones such as auxins, gibberellins and vitamins through the supplied phosphorus, thus fostering plant growth. Prior studies by Singh *et al.*, (2011), Ranpariya *et al.*, (2017) and Singh *et al.*, (2019) have also reported similar observations.

Yield attributes

The data on yield attributes of greengram as influenced by varieties and nutrient management is presented in Table 2. Greengram variety MH-318 (17.9) and SML-668 (16.7) recorded significantly higher pods/plant compared to MH-421 (14.3). Similarly, pod length was found to be significantly higher with MH-318 (8.5 cm) and SML-668 (7.8 cm) over MH-421 (7.5 cm). The 100 seed weight under variety MH-318 and SML-668 was at par and significantly higher over MH-421 at 5.09, 4.81 and 4.48, respectively.

Among nutrient management treatments, plots applied with 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB demonstrated notable superiority in terms of pods/plant (18.1), resulting in a 29.3% increase compared to the control. The higher number of seeds/pod (10.7) was recorded under the treatment of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB. Additionally, for pod length, the most effective treatment was 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB (8.6 cm). Finally, the application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB (5.13 g) resulted in significantly higher 100 seed weight in comparison to the other treatments. The outcomes align with the findings reported by Tyagi and Upadhyay (2015), Prajapati *et al.*, (2016), Mondal and Sengupta (2019) and Singh *et al.*, (2018).

Seed Yield and Harvest Index

The data presented in Table 2 indicated that both varieties and nutrient treatment significantly impacted on the seed yield of greengram. Among varieties, MH-318 exhibited the highest seed yield (1110 kg/ha), significantly surpassing MH-421 (952 kg/ha) but comparable to SML-668 (1013 kg/ha). Further, application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB resulted in the maximum grain yield (1269 kg/ha), surpassing all other treatments. This treatment showcased an 84%, 62% and 48% increase in yield compared to the control, 50% RDF + 50% N/ha (FYM) + ST with *Rhizobium* + PSB and 100% RDF, respectively. The increase in yield may be attributed to increased growth parameters. Similar results have been reported by Pareek *et al.*, (2022), Pandey *et al.*, (2019) and Hussain *et al.*, (2017).

Greengram variety MH-318 (27.1) showed 1.6% and 2.1% higher harvest index, respectively over SML-668 and MH-421. Among the nutrient management treatment, application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB (27.3%) and 100% RDF (26.8%) were statistically at par and had a significantly higher harvest index than 50% RDF + 50% N/ha (FYM) + ST with *Rhizobium* + PSB (26.0%) and control (23.75%). High seed yield in greengram can be attributed to the high yield of yield-contributing attributes, including the number of pods/plant, number of seeds/pod and 100-seed weight. Pandey *et al.*, (2019) and Hussain *et al.*, (2017) have also documented comparable findings.

Economics and Nutrient Uptake

Monitory analysis of greengram as influenced by varieties and nutrient management is presented in Table 2. Greengram MH-318 variety imparted significantly higher net returns (include data) compared to SML-668 (46,230 ₹/ha) and MH-421 (52,972 ₹/ha). Similarly, application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB resulted in the highest net returns (62,743 ₹/ha), followed by 100% RDF (54,922 ₹/ha) and 50% RDF + 50% N/ha (FYM) + ST with *Rhizobium* + PSB (42,772 ₹/ha). The benefit-cost ratio was also higher for MH-318 (3.12) variety compared to the other two varieties and the highest benefit-cost ratio was observed with the application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB (3.40), followed by 100% RDF (3.35).

The uptake of nutrients in greengram, as demonstrated in Table 3, is a function of biological yield and nutrient content. In particular, variety MH-318 exhibited significantly higher NPK uptake compared to SML-668 and MH-421. Application of 75% RDF + 25% N/ha (FYM) + ST with *Rhizobium* + PSB led to significantly higher NPK uptake (128.8 kg/ha), surpassing all other treatments. These findings are consistent with the results reported by Gorade *et al.*, (2014), Tambe *et al.*, (2019) and Dubey *et al.*, (2018).

CONCLUSION

In conclusion, the study suggests that cultivation of greengram variety MH-318 during the summer season, with an application of 75% RDF, 25% N/ha (FYM) and seed treatment with *Rhizobium* and PSB would provide higher yields and economic benefits while maintaining soil fertility.

References

- Anonymous (2022). Ministry of Agriculture and Farmers Welfare, Govt. of India. *11th Advance Estimates, 2021-22*.
- Bhardwaj, S., Sutaliya, J.M., Parkash, R., Bhardwaj, K.K., Kumar, N., Ahlawat, I. and Pareek, A. (2022). Zero Tillage in Combination with Seed Treatment by Different Biofertilizers Increased Soil Organic Carbon, Macro and Micronutrients Status and Nodulation under Faba Bean (*Vicia faba* L.). *Legume Research*. DOI: 10.18805/LR-4971.
- Dubey, S.N., Rajpal, S., Rohit, K. and Sauhard, D. (2018). Effect of phosphorus and PSB on growth, nodulation and fertility status in different mungbean (*Vigna radiata* L.) varieties and its residual effect on fodder yield of sorghum in indo-gangetic plain zone of India. *International Journal of Agricultural Sciences* **14**(1): 196-201.
- Gorade, V.N., Chavan, L.S., Jagtap, D.N. and Kolekar, A.B. (2014). Response of green gram (*Vigna radiata* L.) varieties to integrated nutrient management in summer season. *Agricultural Science Digest-A Research Journal* **34**(1): 36-40.
- Hussain, Z., Khan, N., Ullah, S., Liaqat, A., Nawaz, F., Rehman, F. and Ali, M. (2017). Response of mungbean to various levels of biochar, farmyard manure and nitrogen. *World Journal of Agricultural Sciences* **13**(1): 26-33.
- John, A.T., Makkar, S., Swaminathan, S., Minocha, S., Webb, P., Kurpad, A.V. and Thomas, T. (2021). Factors influencing household pulse consumption in India: A multilevel model analysis. *Global food security*, **29**, 100534.
- Lal, B., Rana, K.S., Gautam, P., Rana, D.S., Shivay, Y.S., Meena, B.P. and Singh, P. (2015). Ethiopian mustard–chickpea intercropping system is a viable option for yield advantage in dryland condition of north India-Part II. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences* **86**(3): 757-766.
- Meena, B.P., Kumar, A., Dhar, S., Paul, S. and Kumar, A. (2016). Productivity, nutrient uptake and quality of popcorn and potato in relation to organic nutrient management practices. *Annals of Agricultural Research New Series* **37**(1): 72-79.
- Mondal, R. and Sengupta, K. (2019). Study on the performance of mungbean varieties in the new alluvial zone of West Bengal. *Journal of Crop and Weed* **15**(1): 186-191.
- Nair, R.M., Schafleitner, R., Kenyon, L., Srinivasan, R., Easdown, W., Ebert, A.W. and Hanson, P. (2012). Genetic improvement of mungbean. *SABRAO Journal of Breeding and Genetics* **44**(2): 177-190.
- Pandey, O.M., Shahi, S.K., Dubey, A.N. and Maurya, S. (2019). Effect of integrated nutrient management of growth and yield attributes of green gram (*Vigna radiata* L.). *Journal of Pharmacognosy and Phytochemistry* **8**(3): 2347-2352.

- Panase, V.G. and Sukhatme, P.V. (1967). *Statistical methods for Agricultural workers*. ICAR Publication, New Delhi: 199-210.
- Pareek, A., Satyajeet, Satpal, Bhardwaj, S. and Nagora. M. (2022). Effect of varieties and fertility levels on fodder productivity, NPK content, uptake and protein content of summer green gram (*Vigna radiata* L.). *Forage Research* **47**(4):507-512.
- Prajapati, S.K., Tyagi, P.K., Chourasia, S.K. and Upadhyay, A.K. (2016). Effect of integrated nutrient management practices on growth and yield of summer mungbean (*Vigna radiata* L.). *Technofame-A Journal of Multidisciplinary Advance Research* **5**(1): 102-107.
- Ranpariya, V.S., Polara, K.B., Hirpara, D.V. and Bodar, K.H. (2017). Effect of potassium, zinc and FYM on content and uptake of nutrients in seed of summer green gram (*Vigna radiata* L.) and post-harvest soil fertility under medium black calcareous soil. *International Journal of Chemical Studies* **5**(5): 1055-1058.
- Rosales, C.M., Rivera, F., Hautia, R.A. and Del Rosario, E. (1995). Nitrogen fixation by mung bean (*Vigna radiata* L.) under field conditions in the Philippines as quantified by ¹⁵N isotope dilution (No. PNRI-C (AG)--95001). Philippine Nuclear Research Inst.
- Singh, G., Sekhon, H.S., Singh, G., Brar, J.S., Bains, T.S. and Shanmugasundaram, S. (2011). Effect of plant density on the growth and yield of mungbean [*Vigna radiata* (L.) Wilczek] genotypes under different environments in India and Taiwan. *International Journal of Agricultural Research* **6**(7): 573-583.
- Singh, R.P., Dhillon, B.S. and Sidhu, A.S. (2019). Productivity of summer moong (*Vigna radiata* L.) as influenced by different sowing dates and varieties. *Journal of Pharmacognosy and Phytochemistry* **8**(3): 781-784.
- Singh, Y.P., Tomar, S.S., Singh, A.K. and Yadav, R.P. (2018). Nutrient management and irrigation scheduling effect on blackgram (*Vigna mungo*)-frenchbean (*Phaseolus vulgaris*) yield, economics, water productivity and soil properties. *Journal of Soil and Water Conservation* **17**(1): 58-64.
- Tambe, A.D., Surva, U.S. and Kumbhkar, J.S. (2019). Integrated Nutrient Management in soybean-wheat cropping system. *Journal of Agroecology and Natural Resource Management* **6**(4): 160-162.
- Tyagi, P.K. and Upadhyay, A.K. (2015). Effect of integrated nutrient management on yield, quality, nutrients uptake and economics of summer greengram. *Annals of Plant and Soil Research* **17**(3): 242-247.
- Vavilov, N.I. (1926). Studies on the origin of cultivated plants. *Bulletin of Applied Botany of Genetics and Plant Breeding* **16**(2): 139-148.

Table 1: Effect of varieties and nutrient management practices on plant height (cm) and dry matter accumulation (g/plant) of summer greengram.

| Treatment | Plant height (cm) | | | | Dry matter accumulation (g/plant) | | | |
|--------------------------------------------------------------------|-------------------|------------|------------|------------|-----------------------------------|-------------|-------------|-------------|
| | 15 DAS | 30 DAS | 45 DAS | At Harvest | 15 DAS | 30 DAS | 45 DAS | At Harvest |
| Variety | | | | | | | | |
| V ₁ - MH-421 | 10.7 | 21.1 | 33.5 | 37.5 | 0.33 | 1.63 | 8.19 | 14.3 |
| V ₂ - MH-318 | 11.3 | 23.9 | 36.2 | 40.5 | 0.35 | 1.72 | 9.01 | 15.5 |
| V ₃ - SML-668 | 11.0 | 22.6 | 35.3 | 38.1 | 0.34 | 1.67 | 8.53 | 15.1 |
| CD (p≤0.05) | NS | 1.3 | 2.0 | 2.2 | NS | 0.06 | 0.36 | 0.91 |
| Nutrient Treatment | | | | | | | | |
| F ₁ - Control | 9.4 | 20.1 | 33.3 | 36.3 | 0.32 | 1.61 | 7.84 | 13.6 |
| F ₂ - 100% RDF | 11.6 | 23.5 | 35.5 | 39.2 | 0.34 | 1.69 | 8.78 | 15.4 |
| F ₃ - 75% RDF + 25% N/ha (FYM) + <i>Rhizobium</i> + PSB | 12.0 | 23.7 | 36.3 | 40.7 | 0.35 | 1.70 | 9.16 | 16.2 |
| F ₄ - 50% RDF + 50% N/ha (FYM) + <i>Rhizobium</i> + PSB | 11.1 | 22.7 | 34.7 | 38.7 | 0.33 | 1.67 | 8.51 | 14.7 |
| CD (p≤0.05) | NS | 1.0 | 1.8 | 2.0 | NS | 0.06 | 0.39 | 1.19 |

*Significant at p≤0.05, RDF: Recommended dose of fertilizer, FYM: Farm Yard Manure, PSB: Phosphorus Solubilizing Bacteria, DAS: Days after sowing, NS: Non-Significant

Table 2: Effect of varieties and nutrient management practices on yield attributes, yield and economics of summer greengram.

| Treatment | No. of pods per plant | No. of seeds per pod | Pod length (cm) | 100 seed weight (g) | Seed yield (kg/ha) | Harvest index (%) | Net returns (₹/ha) | B:C ratio |
|--------------------------------------------------------------------|-----------------------|----------------------|-----------------|---------------------|--------------------|-------------------|--------------------|-----------|
| Variety | | | | | | | | |
| V ₁ - MH-421 | 14.3 | 9.9 | 7.48 | 4.48 | 952 | 25.0 | 41898 | 2.68 |
| V ₂ - MH-318 | 17.9 | 10.6 | 8.54 | 5.09 | 1110 | 27.1 | 52972 | 3.12 |
| V ₃ - SML-668 | 16.7 | 10.1 | 7.80 | 4.81 | 1013 | 25.8 | 46230 | 2.86 |
| CD (p≤0.05) | 1.44 | 0.54 | 0.65 | 0.33 | 108.2 | 1.10 | - | - |
| Nutrient management practice | | | | | | | | |
| F ₁ - Control | 14.0 | 9.6 | 7.22 | 4.34 | 689 | 23.8 | 27697 | 2.34 |
| F ₂ - 100% RDF | 17.3 | 10.4 | 8.13 | 4.91 | 1117 | 26.8 | 54922 | 3.35 |
| F ₃ - 75% RDF + 25% N/ha (FYM) + <i>Rhizobium</i> + PSB | 18.1 | 10.7 | 8.55 | 5.13 | 1269 | 27.3 | 62743 | 3.40 |
| F ₄ - 50% RDF + 50% N/ha (FYM) + <i>Rhizobium</i> + PSB | 15.8 | 10.1 | 7.87 | 4.79 | 1024 | 26.0 | 42772 | 2.47 |

| | | | | | | | | |
|--------------------|------|------|------|------|--------------|-------------|---|---|
| CD (p≤0.05) | 1.31 | 0.48 | 0.55 | 0.28 | 129.4 | 1.04 | - | - |
|--------------------|------|------|------|------|--------------|-------------|---|---|

*Significant at p≤0.05, RDF: Recommended dose of fertilizer, FYM: Farm Yard Manure, PSB: Phosphorus Solubilizing Bacteria, NS: Non-Significant

Table 3: Effect of varieties and nutrient management practice on N, P and K uptake (kg/ha) by plant of summer greengram

| Treatment | Total nitrogen uptake (kg/ha) | Total phosphorus uptake (kg/ha) | Total potassium uptake (kg/ha) |
|--------------------------------------------------------------------|--------------------------------------|----------------------------------------|---------------------------------------|
| Variety | | | |
| V ₁ - MH-421 | 93.7 | 9.6 | 31.3 |
| V ₂ - MH-318 | 111.6 | 13.4 | 37.8 |
| V ₃ - SML-668 | 102.1 | 11.2 | 34.6 |
| CD (p≤0.05) | 6.42 | 0.92 | 2.98 |
| Nutrient management practice | | | |
| F ₁ - Control | 68.4 | 7.06 | 23.4 |
| F ₂ - 100% RDF | 111.3 | 12.6 | 37.4 |
| F ₃ - 75% RDF + 25% N/ha (FYM) + <i>Rhizobium</i> + PSB | 128.8 | 14.9 | 43.2 |
| F ₄ - 50% RDF + 50% N/ha (FYM) + <i>Rhizobium</i> + PSB | 101.2 | 10.9 | 34.1 |
| CD (p≤0.05) | 7.05 | 1.17 | 3.32 |

*Significant at p≤0.05, RDF: Recommended dose of fertilizer, FYM: Farm Yard Manure, PSB: Phosphorus Solubilizing Bacteria