

## Original Research Article

### Assessing the Different Bio-priming Methods and Knowledge Dissemination through On Farm Trial in Salem district

#### ABSTRACT

One of the most important pre sowing seed improvement techniques for managing biotic and abiotic challenges is seed bio-priming, which ensures uniform stand establishment under adverse conditions. "Bio-priming" seed treatment combines biological (inoculating seeds with protective organisms) and physiological (hydrating seeds) components of vigour improvement & disease management. Recently, it has been employed as an alternate technique for not only eradicating numerous soil- and seed-borne diseases and also improves the establishment and vigour. In the current study, an On-Farm Trial bhendi seeds were soaked with two biopriming treatments Viz., (i) 10% *Pseudomonas fluorescense*, (ii) 10% *Trichoderma viridi*, for 6 hours was conducted in farmers' fields during kharif, 2020. Untreated seeds served as the control. The findings showed that 10% *Pseudomonas fluorescense* outperformed than trichoderma and untreated seed in terms of yield (172.5 q/ha), net return (Rs. 2,58,750/ha), and benefit-cost ratio (2.59) than the untreated control. The 10% trichoderma primed seeds, which was the next-best treatment, had high yields, net returns, and BCR values of 169.8q/ha, Rs. 2,54,700/ha, and 2.56 respectively. So, it was concluded that biopriming of seeds with 10% *Pseudomonas fluorescense* for six hours increased the production of bhendi. Hence, through The dissemination of extension activities like training and method demonstration revealed that increased the understanding of farmers about biopriming is from 5.4 to 52.5 percent.

Key words: Bhendi, seed biopriming, Knowledge dissemination and seed yield

#### INTRODUCTION

Bio-priming is an originative skill and cost effective seed treatment that assimilates biological (inoculation of seed with beneficial organism to protect seed) and physiological facets (seed hydration) of disease control [1]. By excreting chemicals and solubilizing minerals, bio-priming directly contributes to the enrichment of plant development [2]. Advance technologies are being used in modern agriculture to increase crop productivity and break through yield barriers. Creating diverse seed enhancement technologies play a significant role in ensuring uniform field emergence, better crop stands and higher yield for various crops. Bio-priming, which combines a variety of plant extracts, microbial products and biotic agents to manage seed crops and target them against biotic and abiotic stresses, has been hailed as a

Comment [MB1]: and

Comment [MB2]: to eradicate

Comment [MB3]: but also, to improve

Comment [MB4]: fluorescens

Comment [MB5]: Capital T=Trichoderma

Comment [MB6]: Not understood

novel management strategy because it uses fewer chemicals, increases the efficacy of the seeds, lowers management costs, eliminates pollution risks and interferes with biological equilibrium to the least amount possible. Therefore, using "On Farm Testing" in farmers' fields, an effort was undertaken to gauge the effectiveness of seed biopriming with novel biocontrol agents (*Tirchodermaviridae* and *Pseudomonas flurescence*) in bhendi.

## MATERIALS AND METHODS

In order to shorten the time lag between the development of a technology and its adoption by farmers, Krishi Vigyan Kendra, Salem demonstrate the newest agricultural technologies to farmers and extension workers of the State Agriculture Department through "On Farm Testing (OFTs) and Frontline demonstrations (FLDs). During the kharif season in 2020, the "On Farm Testing" was done at five sites in Puthuragraharam Village, Veerapandi block of Salem district, Tamilnadu. Farmers received information about seed biopriming treatments and other improved crop management techniques through training programmes prior to the OFT. The CO(Bh)H4 Bhendi seeds were bioprimed for 6 hours with 10% *Trichoderma viridi*, 10% *Pseudomonas flurescence* and were compared with untreated control seeds. The beneficiaries received the crucial inputs, including bhendi seeds (CO(Bh)H 4), Arka vegetable special, and IPDM components. The Tamil Nadu Agricultural University's approved bhendi farming package and methods were taught to farmers. The KVK scientists routinely visited the OFT fields during various crop stages to ensure that nutrients and plant protection measures were applied in a timely manner. They also offered the farmers other suggested measures and collected feedback data on each stage to further improve the research and extension programme. The yield data was assessed using the cumulative yield method, with 10% *Tirchoderma viridi*, 10% *Pseudomonas flurescence* and iii). A field day was held in conjunction with extension agents from the Department of Agriculture to share the improved farming practices used by OFT farmers with other farmers. A well planned interview schedule with a predetermined list of questions was used to gather the pertinent information regarding the cost of cultivation, market preferences and other restrictions. Each farmer provided statistics on crop output and profitability for the OFT and control plots, which were then averaged across all locations. Using the appropriate statistical methods, the gathered data were pooled and tabular analysis was completed to determine the technical gap. When the cropping phase is completed, furthermore, using the knowledge level of OFT farmers using improved bhendi production technology was compared before and after KVK interventions and knowledge test was done as recommended by [3]. The respondents also received the various improved production technologies such as the choice of improved varieties,

**Comment [U7]:** kharif

**Comment [U8]:** OFT done for what aspect should be mentioned here clearly.

**Comment [U9]:** If possible, What nutrients are present should be mentioned

**Comment [U10]:** No connection was observed between this and previous sentence.

**Comment [U11]:** Please provide the name also

Integrated Nutrient Management (INM), foliar spray of crop boosters (Arka vegetable special), and Integrated Pest and Disease Management (IPDM).

[4] States that the knowledge level was scored, with each correct response receiving two points and each erroneous response receiving one point. The pre- and post-evaluation scores were evaluated, and the respondents' knowledge Index was determined as shown below.

**Comment [U12]:** this can be provided as reference and Method used by the authors should be given here

The formula used for the calculation of knowledge index of each respondent was

$$\text{Knowledge Index} = \frac{K}{P} \times 100$$

Where,

K - Knowledge scores obtained by an individual respondent

P - Maximum possible scores for all items

The respondents were classified into three categories such as low, medium and high using mean and standard deviation.

## RESULTS AND DISCUSSION

The results of the current study, which evaluated the efficiency of seed biopriming with *Tirchoderma viridae* and *Pseudomonas flurescence* of bhendi from seed to harvest in farmers' fields, are summarized below.

**Table 1. Performance of biopriming in bhendi under OFT programme**

Technology option	Field Emergence (%)	Plant population at 60 DAP (%)	Fruit Nos.	No. Harvest	Fruit Yield (q/ha)
Untreated	82	74	23	19	159.7
Biopriming with 10% <i>Pseudomonas flurescence</i>	94	90	27	22	172.5
Biopriming with 10 % <i>Tirchoderma viridi</i>	90	84	25	21	169.8

The results showed that the farmers, using 10% *Pseudomonas flurescence* and *Tirchoderma viridi* 10% primed seeds, respectively, produced an average fruit production of 172.5 and 169.8 q/ha, compared to the control's 159.7 q/ha (Table 1). The difference in fruit output between *Pseudomonas flurescence* primed seed and control. *Pseudomonas flurescence* primed seeds also performed exceptionally well in terms of plant population field emergence number of fruits per plant and number of harvest.

**Comment [U13]:** incomplete sentence

**Comment [U14]:** no comma

The same outcome is also supported by [5], who discovered that seeds treated with *Pseudomonas flurescence* had improved bhendi yield and growth. An increase in okra plant height, fruit production, and

yield per ~~plant had received~~ [6]. Bioprime seeds with *Trichoderma harzianum*, ~~which increases~~ plant tolerance to abiotic stressors and controls disease-causing organisms through the release of antimicrobial compounds [7,8]. ~~As a result of seed treatment~~ with *Pseudomonas fluorescens*, and *Trichoderma viridi* have been shown to protect a variety of crop plants by hyperparasitizing pathogenic fungi [9,10]. ~~When used~~, bioagents have a remarkable ability for multiplication and to grow exponentially, they can even resist stress conditions by spores with strong walls form [11]. The results of the present experiment are supported by data from [12,13] on improvements in the vegetable output of bitter gourd and brinjal respectively. *Pseudomonas fluorescens* were used to increase yield in white cabbage and cauliflower, according to [14].

**Comment [U15]:** incomplete sentence. i.e., it should be mentioned why increased yield obtained?

**Comment [U16]:** enhanced

**Comment [U17]:** The treatment.....

Seed hydration ~~is~~ followed by the introduction of helpful microorganisms to the seed surface in a process known as seed bioprime, ~~which~~ is regarded as an advanced method of seed treatment [1,15,16]. According to initial inoculum levels, it was found that during bioprime, bacterial populations increased (10 to over 10,000 folds) [17]. Different priming techniques are used in the seed bioprime process, such as wet finely crushed lignite or coal (solid matrix priming) [18] or moist conditions in a plastic bag [19]. Before priming, it's crucial to disinfest seeds in ~~an~~ order to minimise or get rid of the undesirable microorganisms, if the seeds are diseased or contaminated with pathogens, this will be amplified during the priming process, having unfavourable consequences on emerging plants [1, 20, 21]. Additionally, the survival of helpful bacteria used during the seed bioprime process may be harmed by the growth of undesirable indigenous microorganisms [21].

The economic analysis of bioprime was studied based on gross income which was calculated with average yield multiplied by prevailing market price of Rs.1000 /quintal during that particular year. It could be observed that the average net income of the ~~₹~~ *Pseudomonas fluorescens* bioprime seeds was Rs.1,01,250/- ha, which was Rs.98,550 per ha, and Rs.88,950 in *Trichoderma viride* and control plots, respectively. Further, the benefit cost ratio was 2.42, 2.38 and 2.26 respectively (**Table 2**).

**Table 2. Economic analysis of bioprime seeds in bhendi**

Treatments	Gross cost (Rs./ha)	Gross income(Rs./ha)	Net income(Rs./ha)	BCR
Untreated	70,750	1,59,700	88,950	1:2.26
Bioprime with 10 % <i>Pseudomonas fluorescens</i>	71,250	1,72,500	1,01,250	1:2.42
Bioprime with 10% <i>Trichoderma viride</i>	71,250	1,69,800	98,550	1:2.38

Knowledge is a requirement for innovation adoption and would give farmers the ability to fully understand a technology and its relative advantages. When evaluating the effective diffusion of

technology, the level of farmer knowledge regarding the effects of biopriming and other improved techniques is essential. So, an effort was undertaken to compare the level of knowledge before and after KVK interventions during the conduct of OFT programme. The degree of expertise of medium category bhendi growing farmers in upgraded technologies increased from 52.5 to 62.5% (Table 3). Due to extension activities, the percentage of farmers with modest expertise was reduced from 35 to 12.5%.

Table 3. Distribution of respondents based on their knowledge level (n = 40)

S.No.	Category	Before OFT		After OFT	
		Number	Per cent	Number	Per cent
1	Low	14	35.0	5	12.5
2	Medium	21	52.5	25	62.5
3	High	5	12.5	10	25.0
	Total	40	100	40	100

When it comes to farmers' understanding of new technologies, it can be shown that before the OFT programme was implemented, only about 12.5 percent of farmers knew about presowing seed treatment of biopriming, and that number rose to 87.5 percent after the programme was implemented (Table 4).

Similar to this, the percentage improvement in farmers' knowledge on the usage of hybrid bhendi, optimum seed rate, seed biopriming, foliar nutrition, INM, IWM and IPDM was 55.0, 15.0, 75.0, 32.5, 42.5, 40.0, and 65.0 percent, respectively. Improved level of knowledge on bhendi producing technologies could be due to the extension activities like training sessions, method demonstrations, and field days during the conduct of the OFT programme by KVK. Additionally, KVK Salem contributed significantly to the spread of technologies by providing important material throughout the extension operations.

Comment [U18]: production

Comment [U19]: conducted during

Table 4. Knowledge level of the bhendi farmers in improved production technologies

(n=40)

S. No.	Technologies	Before OFT programme		After OFT programme		% Increase in Knowledge level of farmers
		Number	%	Number	%	
1	Application of FYM	40	100	40	100	0
2	Improved Bhendi varieties from public sector	10	25.0	32	80.0	55.0

3	Optimum seed rate	32	80.0	38	95.0	15.0
4	<b>Seed biopriming</b>	5	12.5	35	87.5	75.0
5	Foliar nutrition	20	50.0	33	82.5	32.5
6	INM	19	47.5	36	90.0	42.5
7	IWM	10	25.0	26	65.0	40.0
8	IPDM	10	25.0	36	90.0	65.0

## CONCLUSION

It may be established that seeds biopriming with 10% *Pseudomonas fluorescens* out-performed other bioprimed seeds and untreated control in terms of yield (165.1 q/ha), net return (Rs. 93,340/ha), and cost benefit ratio (2.36) followed by the 10% *Trichoderma viride* primed seeds, which had high yield, net return and BCR values of 162.3 q/ha, Rs, 90,340/ha and 2.28, respectively. ~~So it was discovered that biopriming of seeds with 10% *Pseudomonas fluorescens* for six hours increased the yield of bhendi~~ and the **knowledge on seed bio-priming** rose from 12.5 to 87.5 percent.

**Comment [U20]:** Repetition, hence not needed

**Comment [U21]:** Please mention knowledge increased among whom and due to what reason?

## REFERENCES

1. P.P Reddy (2013). Bio-priming of seeds. **In:** Reddy, P.P. (Ed.), *Recent Advances in Crop Protection*. Springer, India, Pp. 83–90.
2. V. Sukanya, R.M.Patel, K.P. Suthar and D. Singh (2018). An Overview: Mechanism Involved in Bio-Priming Mediated Plant Growth Promotion. *International Journal of Pure and Applied Bioscience* 6, 771–783.
3. A.K. Singh (1986). Tests, Measurements and Research Methods in Behavioral Sciences. Tata McGraw-Hill publishing company Ltd., New Delhi.
4. P. Madhan(2002). Generation, dissemination and adoption of rice varieties of Reginal Research Station, Ambasamudram –An analysis. M.Sc. (Ag.) Thesis, TNAU, Coimbatore.
5. M. Rafique, A. Riaz, A. Anjum, M.A. Qureshi and F. Mujeeb(2018). Role of Bioinoculants for Improving Growth and Yield of Okra (*Abelmoshusesculentum*).*Universal Journal of Agricultural Research* , 6(3): 105-112
6. A. K. Rai, and A.K. Basu(2014). Pre-Sowing Seed Bio-Priming in Okra: Response for Seed Production. *The Bioscan*, 9(2): 643-647.
7. P. Ahmad, A. Hashem, E. F Abd-Allah, A. A Alqarawi, R. John, D. Egamberdieva and S. Gucel, (2015). Role of *Trichoderma harzianum* in mitigating NaCl stress in Indian mustard (*Brassicajuncea* L) through antioxidative defense system. *Frontiers in plant science*, 6, 868.
8. F. Mastouri, T. Bjorkman and G. E. Harman (2012). *Trichoderma harzianum* enhances antioxidant defense of tomato seedlings and resistance to water deficit. *Molecular plant-microbe interactions*, 25(9), 1264-1271.
9. L.W. Durrell, (1968). Hyphal invasion by *Trichoderma viridae*. *Mycopathol Mycol. Appl.*, 35: 138-144

10. H. L Barnett and H.A. Binder (1973). The fungal host parasite relationship. *Annual Review of Phytopathology*, **11**: 273-292
11. B.G.S BharathLokesh, and H.S. Shetty (2005). Effects of fungicides and bioagents on seed mycoflora, growth and yield of watermelon. *Integrative Biosciences*, **9**: 75-78
12. I. Janaki, S.Suresh and P.Karuppuchamy (2012). Efficacy and economics of bio-pesticides for the management of papaya mealybug, *Paracoccus marginatus* (Williams and Granara de Willink) in brinjal (*Solanum melongena* L.). *Journal of Biopesticides*, **5**(1): 87-90
13. K.S. Naveen Kumar, B.V.Sowmyamala, P.G.Sadhan Kumar, P.N. Vasudev, R. Vasantha Kumar and H.T. Nagaraj (2012). Effect of plant growth promoting rhizobacteria (PGPR) on growth and yield of bitter melon. *International Journal of Applied Biology & Pharmaceutical Technology*, **3**: 1-7
14. AbdAlla, M. A. and F. S. El-Shoraky (2017). Impact of biological agents and plant essential oils on growth, quality and productivity of cabbage and cauliflower plants correlated to some diseases control. *Journal of Sustainable Agricultural Sciences*, **43**(1): 27-38
15. Singh, H.B., 2016. Seed bio-priming: a comprehensive approach towards agricultural sustainability. *Indian Phytopath.* **69**, 203–209.
16. Singh, V., R.S. Upadhyay, B.K. Sarma and H.B. Singh (2016). Seed bio-priming with *Tirchodermaviridae asperillum* effectively modulate plant growth promotion in pea. *Int. J. Agric. Environ. Biotechnol.* **9**, 361–365.
17. Callan, N.W., D.Mathre and J.B., Miller (1990). Bio-priming seed treatment for biological control of *Pythium ultimum* preemergence damping-off in SH-2 sweet corn. *Plant Dis.* **74**, 368–372
18. Harman, G.E. and A.G. Taylor (1988). Improved seedling performance by integration of biological control agents at favorable pH levels with solid matrix priming. *Phytopathology* **78**, 520–525.
19. Callan, N.W., D.E. Mathre and J.B. Miller (1991). Field performance of sweet corn seed bio-primed and coated with *Pseudomonas fluorescens* AB254. *Hortic. Sci.*, **26**, 1163–1165
20. Jain, A., A. Singh, S. Singh and H.B. Singh (2015). Biological management of *Sclerotinia sclerotiorum* in pea using plant growth promoting microbial consortium. *J. Basic Microbiol.* **55**, 961–972.
21. Singh, A., B.K. Sarma, R.S. Upadhyay and H.S. Singh (2013). Compatible rhizosphere microbes mediated alleviation of biotic stress in chickpea through enhanced antioxidant and phenylpropanoid activities. *Microbiol. Res.* **168**, 33–40.