

Original Research Article

Effect of foliar application of different chemicals on yield attributes and yield of Black gram [*Vigna mungo* (L.) Hepper] under rainfed condition

Comment [H1]: You can modify the topic name

Comment [H2]: You can do- Effect of foliar application of different chemicals on yield and yield attributes of Black gram [*Vigna mungo* (L.) Hepper] under rainfed condition

Comment [H3]: Write full name of the university

ABSTRACT

A field experiment was performed at the “TCA., research farm of Dholi (Muzaffarpur), which is a unit of the Dr. RPCAU, Pusa (Samastipur), Bihar, during the *Kharif* season of 2021” to investigate the “**Effect of foliar application of different chemicals on yield attributes and yield of Black gram [*Vigna mungo* (L.) Hepper] under rainfed condition**”. The design used for making layout of experimental unit was RBD, that was replicated three times, with ten treatments in each replication for a total of 30 plots. The experiment conducted with 10 treatments *viz.*, Absolute Control (T₁), Water Spray (T₂), ZnSO₄ @ 0.5% Spray (T₃), NPK (19-19-19) @ 2% Spray (T₄), Urea @ 2% Spray (T₅), DAP @ 2% Spray (T₆), Salicylic acid @ 100 ppm Spray (T₇), Potassium Nitrate @ 2% Spray (T₈), Boron 0.25 % Spray (T₉), Potassium Chloride @ 0.2% Spray (T₁₀). The foliar spray of different treatments applied had a significant impact on the number of pod number plant⁻¹, seed count pod⁻¹ and the length of the pod. However, foliar chemicals had no appreciable impact on the weight of 100 seeds. At pre-flowering and pod-initiation stage, foliar treatment of “NPK (19-19-19) @ 2 % spray (T₄)” resulted in significantly greater seed production and stover output. Different chemical administration to the foliage had no significant effect on harvest index.

Keywords: *Foliar application, Rainfed, Spray*

Comment [H4]: Add minimum 4-5 keywords

INTRODUCTION

Pulses are marvellous gift of the mother nature with distinctive capability of biological nitrogen fixation (BNF). The overall requirement of protein content in a balanced diet of vegetarian population of the world is mainly fulfilled by pulses. Pulses also supply nutritive fodder, especially for milch cattle and maintain fertility of soil either through BNF or as a green manure crop and accordingly, play a key role in promoting sustainable agriculture. One of the most coveted pulse crops growing in rainfed areas across the nation is the Black gram or Urd bean [*Vigna mungo* (L.) Hepper]. Urd bean belongs to the “Leguminosae” family and “Papilionaceae” sub-family having chromosome number 2n=22. It is also known as “Mash bean and Urd bean”. Urd bean seeds are perfect blend of all nutrients that includes

Comment [H5]: Add references of these lines if possible.

approximately 23-27 % protein, 60-65 % carbohydrates, 1-1.4 % oil, 3.5-5 % fiber, 4-5.5 % ash on dry weight basis and minerals, amino acids, and vitamins.

Comment [H6]: Add references

The Urd bean is mostly grown in rainfed conditions under poor management strategies, as well as due to various physiological, bio-chemical, and genetic factors of crop itself, which ultimately contribute to its relatively low yield potential. In addition to genetic make-up, physiological factors such as insufficient assimilate partitioning, poor pod setting as a direct consequence of flower abscission, and a lack of nutrients during or after a crucial time of crop growth, including a number of diseases and pests, constitute the major factors limiting the yield of Urd bean (Mahala *et al.*, 2001).

Application of fertilizers to soil and due to formation of certain soil complexes, the uptake of necessary elements becomes difficult for the plants. The applied fertilizers are not fully utilized by the plants. One of them is the foliar administration of various nutrients to optimise the crop's genetic potential. Because less fertilizers are needed for foliar application than for soil application, therefore foliar application of nutrients seems to be preferable. Because fertilizer costs are rising daily, it is essential to boost the productivity of legume crops through lowering fertiliser costs by applying nutrients as foliar spray.

Comment [H7]: Add references if you can

The most effective treatment for increasing the Black gram's LAI, LAD, specific leaf weight (g), overall dry matter accumulation, and seed yield was foliar spray of nutrients mixed with salicylic acid at a concentration of 100 ppm at 20, 30, and 40 DAS (Amutha *et al.*, 2012). The slow growth & development of nodule senescence may be addressed by applying N to the leaves at specific stages, which ultimately led to greater growth and production of Black gram without involvement of root absorption at pivotal stages.

Comment [H8]: Add references

MATERIALS AND METHODS

The field research was performed at the "TCA., research farm of Dholi (Muzaffarpur), which is a unit of the Dr. RPCAU, Pusa (Samastipur), Bihar", during the *Kharif* season of 2021. For this, a field with a consistent textural make-up and homogeneous fertility was chosen. Soil in the research plot was alluvial & found to be calcareous in nature which developed due to the sedimentation of the river "Burhi Gandak" over the time. The soil on the experimental plot had low levels of accessible organic carbon (0.395), nitrogen (190.35 kg/ha), P₂O₅ (21.34 kg/ha), and K₂O (135.68 kg/ha), all of which suggested low fertility. The pH and EC of the soil was 8.44 and 0.31 dSm⁻¹ respectively. Data pertaining to the Physico-chemical properties of the soil is given in table 1. The mean maximum temperature ranged during crop season

Comment [H9]: Write Full name

was in between 27.9 °C to 33.4 °C. The mean minimum temperature ranged during crop season was in between 14.1 °C to 27.0 °C. The Relative Humidity (RH) observed during crop season varied from 98.0% - 100% at 7 AM whereas, at 2 PM it ranged between 73.7% - 94.7%. The total amount of rainfall (675.63 mm) was received during the crop season (August- November 2021). There were ten treatments and they replicated thrice. Under replications, the treatments were allocated randomly. RDF “20 kg/ha N, 20 kg/ha P₂O₅, 20 kg/ha K₂O” is applied as basal dose in all the treatments “except in Absolute control (T₁)”. According to experimental treatments in each plot, the appropriate amount of foliar nutrients/chemicals and water was measured and then prepared a uniform solution and same solution was uniformly sprayed using conical-shaped nozzles with knapsack sprayer. As per treatments, first foliar administration (Fig 1) of nutrients was applied at 33 DAS *i.e.*, at pre flowering stage and again it was applied at pod initiation stage of plant, which was coincided at 53 DAS of crop under study. Collected data at various growth stages and upon harvest were analysed statistically in accordance with Gomez and Gomez's recommendations (1984). The "F" test's significance threshold was set at $p=0.05$.



Fig. 1. Foliar spray of various nutrients

Table 1. Physico-chemical properties of the soil

Sl. No.	Detail	Value determined	Procedure followed	References
1	Sand (%)	49.75	International pipette Method	Piper, 1966
2	Silt (%)	38.65	„	„
3	Clay (%)	11.60	„	„
4	Textural class	Sandy loam	Triangular diagram	„
5.	EC (dSm ⁻¹ at 25 ⁰ C)	0.31	Conductivity bridge	Jackson, 1973

6	pH	8.44	Glass electrode pH meter	„
7	Organic Carbon (%)	0.39	Walkley and Black	Walkley and Black, 1934
8	Available N (kg/ha)	190.35	Alkaline permanganate	Subbiah and Asija (1956)
9	Available P ₂ O ₅ (kg/ha)	21.34	Olsen's Method	Olsen et al, 1954
10	Available K ₂ O (kg/ha)	135.68	Flame photometer	Jackson, 1973

RESULT AND DISCUSSION

Yield attributes:

A crop's yield is a collective function of multiple yield-attributing factors *viz.*, “number of pods plant⁻¹, number of seeds pod⁻¹, length of pod and seed index”. The results of the current investigation after harvest on yield attributes are reported in Table 2.

Number of pods plant⁻¹

Per plant seed yield and productivity is directly correlated with number of pods plant⁻¹. Foliar application of treatment at two critical growth stages *i.e.*, pre-flowering and pod initiation has significant effect on pod number plant⁻¹. The data presented in table 2 revealed that pod number plant⁻¹ after harvest were significantly higher in “NPK (19-19-19) @ 2% Spray (T₄)” *i.e.*, 25.31 as compared to all the remaining treatments applied besides that it was *at par* with following treatments *i.e.*, Urea @ 2% Spray (T₅), DAP @ 2% Spray (T₆) and Potassium Nitrate @ 2% Spray (T₈). Pod number plant⁻¹ recorded were lowest in absolute control (T₁) *i.e.*, 16.58 after harvest. Increased pod number was due to decrease in flower drop caused by prolonged assimilatory activity of the leaves. Furthermore, the nitrogen and phosphorus given to the foliage during the pre-flowering and pod-initiation stage may have been easily absorbed and readily translocated to the pods, resulting in a greater pod number.

Solaiappan et al., (2002) reported similar findings in red gram.

Comment [H10]: Maintain Spacing between Paragraphs

Number of seeds pod⁻¹

Foliar spray of different chemicals at two critical growth stages, first at 33 DAS (Pre-flowering stage) and second at 53 DAS (Pod-initiation stage) increased seed number pod⁻¹ significantly in “NPK (19-19-19) @ 2% Spray (T₄)” *i.e.*, 5.99 as compared to all other treatments applied besides that it was *at par* with following treatments *i.e.*, Urea @ 2% Spray (T₅), DAP @ 2% Spray (T₆) and Potassium Nitrate @ 2% Spray (T₈) and it is revealed by the data tabulated in Table 2. Lowest seed number pod⁻¹ were noted down in Absolute control (T₁) *i.e.*, 4.01 after harvest (Fig.2).



Fig. 2. Counting of number of seed per pod

Length of pod (cm)

This yield parameter is directly correlated with seed number pod⁻¹ and indirectly influenced economic yield of crop. The data shown in Table 2 revealed that different chemicals were applied as foliar spray at two critical growth stages, first at 33 DAS (Pre-flowering stage) and second at 53 DAS (Pod-initiation stage) increased pod length significantly in “NPK (19-19-19) @ 2% Spray (T₄)” *i.e.*, 4.98 cm as compared to all the remaining treatments applied besides that it was *at par* with following treatments *i.e.*, Urea @ 2% Spray (T₅), DAP @ 2% Spray (T₆) and Potassium Nitrate @ 2% Spray (T₈). Lowest pod length was noted down in Absolute control (T₁) *i.e.*, 3.93cm after harvest. It is due to the administration of nutrients during the reproductive phase, which aided to the transfer of photosynthates from the source to developing pods, resulting in an increase in pod length (4.98 cm). **Thakur *et al.*, (2017)** reported similar result.

Seed index (g)

100- seed weight data shown in Table 2 shows that there was no significant difference observed between various treatments applied at two critical growth stages, first at 33 DAS

Comment [H11]: Maintain Proper spacing between paragraphs.

(Pre-flowering stage) and second at 53 DAS (Pod-initiation stage). Maximum seed index was found in NPK (19-19-19) @ 2% Spray -T₄ (4.54 g) and lowest seed index was found in Absolute control -T₁ (3.81 g). The findings confirmed that the foliar spray of various treatments had no significant effect on hundred seed weight (Seed Index). Although seed index was mostly controlled by the genetic characteristics of the cultivar, as no significant differences between treatments were observed.

Table 2. Influence of different chemicals as foliar spray on Yield attributing characters of Black gram

Treatments	Yield Attributes			
	Pods plant ⁻¹	Seeds pod ⁻¹	Pod length(cm)	Seed index (g)
T ₁ - Absolute Control	16.58	4.01	3.93	3.81
T ₂ - Water Spray	18.90	4.93	4.01	4.12
T ₃ - ZnSO ₄ @ 0.5% Spray	22.11	5.45	4.29	4.39
T ₄ - NPK (19-19-19) @ 2% Spray	25.31	5.99	4.98	4.54
T ₅ - Urea @ 2% Spray	23.60	5.71	4.61	4.49
T ₆ - DAP @ 2% Spray	24.01	5.79	4.80	4.45
T ₇ - Salicylic acid @ 100 ppm Spray	22.01	5.32	4.26	4.37
T ₈ - Potassium Nitrate @ 2% Spray	23.12	5.50	4.44	4.45
T ₉ - Boron 0.25 % Spray	21.12	5.11	4.21	4.25
T ₁₀ - Potassium Chloride @ 0.2% Spray	20.32	4.99	4.12	4.20

SEm (\pm)	1.06	0.26	0.21	0.22
LSD ($p=0.05$)	3.14	0.77	0.64	NS

Comment [H12]: Draw the graphical representation of Table 2 i.e., Figure

YIELD STUDIES:

Seed Yield (kg ha^{-1})

The data tabulated in Table 3 revealed that foliar spray of different chemicals at two critical growth stages, first at 33 DAS (Pre-flowering stage) and second at 53 DAS (Pod-initiation stage) increased seed yield significantly in “NPK (19-19-19) @ 2% Spray (T_4)” i.e., 1120 kg ha^{-1} as compared to all other treatments applied besides that it was *at par* with following treatments i.e., Urea @ 2% Spray (T_5), DAP @ 2% Spray (T_6) and Potassium Nitrate @ 2% Spray (T_8). Lowest seed yield was reported in Absolute control (T_1) i.e., 605 kg ha^{-1} after harvest. This is because of the continuous supply of nutrients from foliar spray during the flowering and pod-initiation stage of the crop, which increased yield parameters such as “number of pods plant^{-1} , number of seeds pod^{-1} , pod length and 100-seed weight”, which had a direct impact on seed yield.

Comment [H13]: Add reference of these lines.

Stover Yield (kg ha^{-1})

Vegetative growth of the crop is directly correlated with stover yield. Data on stover yield after harvest in response to varied foliar chemicals spray are listed in Table 3. Foliar application of different chemicals at two critical growth stages, first at 33 DAS (Pre-flowering stage) and second at 53 DAS (Pod-initiation stage) increased stover yield significantly in “NPK (19-19-19) @ 2% Spray (T_4)” i.e., 2321 kg ha^{-1} as compared to all the remaining treatments applied besides that it was *at par* with following treatments i.e., Urea @ 2% Spray (T_5), DAP @ 2% Spray (T_6) and Potassium Nitrate @ 2% Spray (T_8). Lowest stover yield were reported in Absolute control (T_1) i.e., 1590 kg ha^{-1} after harvest. Stover yield directly dependent on vegetative growth of the crop. Stover yield follows the same pattern as of seed yield. It is primarily due to increased plant height and dry matter formation.

Harvest Index (%)

HI data shown in Table 3 shows that there was no significant difference observed between various treatments applied at two critical growth stages, first at 33 DAS (Pre-flowering stage) and second at 53 DAS (Pod-initiation stage). Maximum Harvest index was found in “NPK (19-19-19) @ 2% Spray - T_4 ” (32.55 %) and lowest Harvest index was found in Absolute control - T_1 (27.57 %).

Comment [H14]: Maintain Spacing

Table 3. Influence of different chemicals as foliar spray on Yield and HI of Black gram

Treatments		Seed Yield (kg ha ⁻¹)	Stover yield (kg ha ⁻¹)	Harvest Index (%)
T ₁ -	Absolute Control	605	1590	27.57
T ₂ -	Water Spray	751	1890	28.43
T ₃ -	ZnSO ₄ @ 0.5% Spray	900	1980	31.25
T ₄ -	NPK (19-19-19) @ 2% Spray	1120	2321	32.55
T ₅ -	Urea @ 2% Spray	1060	2242	32.11
T ₆ -	DAP @ 2% Spray	1091	2301	32.18
T ₇ -	Salicylic acid @ 100 ppm Spray	860	1941	30.72
T ₈ -	Potassium Nitrate @ 2% Spray	991	2131	31.73
T ₉ -	Boron 0.25 % Spray	825	1928	29.98
T ₁₀ -	Potassium Chloride @ 0.2% Spray	812	1920	29.73
	SEm (±)	45	100	1.54
	LSD (p=0.05)	134	297	NS

Comment [H15]: Add Figure of Table no. 3 for improving the quality.

CONCLUSION

Foliar application of “NPK (19-19-19) @ 2% Spray (T₄)” at pre flowering and pod initiation stage significantly increased seed (1120 kg/ha) and stover yield (2321 kg/ha) and yield attributing parameters like pod number plant⁻¹ (25.31), seeds pods⁻¹ (5.99) and pod length (4.98 cm) as compared to all the remaining treatments except treatments T₅, T₆ and T₆ that were *at par* along with T₄. Based on the aforementioned investigation, it could be concluded that foliar treatment of “NPK (19:19:19) 2 % spray at pre-flowering & pod-initiation stage” may be used to boost the productivity & profitability for Black gram under rainfed conditions.

REFERENCES

1. Amutha, R., Nithila, S., & Kumar, T. S. 2012. Management of source limitation by foliar spray of nutrients and growth regulators in Black gram. *International Journal of Plant Sciences (Muzaffarnagar)*, **7**(1): 65-68.
2. Gomez, K. A., & Gomez, A. A. 1984. *Statistical procedures for agricultural research*. John Wiley & Sons.
3. Jackson, M.L. 1973. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi.
4. Mahala, C.P.S., Dadheech, R.C. and Kulhari, R.K. 2001. Effect of plant growth regulators and yield of Black gram at varying levels of phosphorus. *Crop Research*, **18**(1):163-165.
5. Pandey, V., Dwivedi, D. K., Singh, H., & Prasad, S. S. Evaluation of Rhizobium along with Multifunctional Rhizobacteria for Improving the Growth and Yield of Black Gram [*Vigna mungo* (L.) Hepper]. *Legume Research-An International Journal*, **1**, **5**.
6. Piper, C.S 1966. Soil and plant analysis. *Inter Science Publication Corporation. Inc.* New York, PP. 91-92.
7. Solaiappan, U. and Ramiah, S. 2002. Effect of seed treatments, soil and foliar fertilization on N and P on yield attributes of pigeon pea grown under rainfed condition. *Indian J. Agron.*, **35**(3): 234-237.
8. Subbiah, B.V. and Asija, G.L. 1956. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*, **25**(8): 259-260.

Comment [H16]: Please Check it very carefully

Comment [H17]: Check it again, year is missing

Comment [H18]: This reference is missing from manuscript.

9. Thakur,V., Patil., R.P., Patil, J.R., Suma, T.C and Umesh, M.R. 2017. Influence of foliar nutrition on growth and yield of Black gram under rainfed condition. *Journal of Pharmacognosy and Phytochemistry*,6(6): 33-37.
10. Walkley, A. and Black, I.A. 1934. Estimation of soil organic carbon by the chromic acid titration method. *Soil Science*, 37: 29-38.

UNDER PEER REVIEW