

## Influence of Weed Control Measures and Nutrient Management on Nutrient Content, Uptake and Nutrient Use Efficiency of Fennel Crop (*Foeniculum vulgare* Mill.)

### Abstract

A research experiment was conducted during the *Rabi* season of 2021-22 at the College of Agriculture, SKRAU, Bikaner. The objective of the experiment was to evaluate the influence of different nutrient management and weed control treatments on the nutrient content, uptake and nutrient use efficiency of fennel crop. The experiment consisted of sixteen treatment combinations, which included four nutrient management treatments and four weed control measures. The nutrient management treatments included control (no fertilizer applied), 75 % RDF, 100 % RDF and 125 % RDF and four weed control measures- weed free, pendimethalin @ 0.75 kg ha<sup>-1</sup> (PE), oxyfluorfen @ 50g ha<sup>-1</sup> (POE at 25 DAS) and weedy check. These treatments were replicated three times in a factorial randomized block design (FRBD), ensuring the reliability and accuracy of the results. The weed-free treatment was found to be significantly better in increasing the nitrogen, phosphorus, and potassium content in both the seed and straw of the crop. Additionally, the weed-free treatment also resulted in higher nutrient uptake by the crop compared to the weedy check treatment. Among the nutrient management treatments, the application of 100% RDF showed the most promising results. It improved the nutrient content in both the seed and straw of the crop and recorded the highest nutrient use efficiency. This efficiency was found to be at par with the application of 125% RDF and significantly higher than the remaining treatments. Overall, the experiment demonstrated the importance of weed control and proper nutrient management in enhancing the nutrient content and uptake by the crop. The weed-free treatment and the application of 100% RDF were identified as the most effective strategies in achieving these goals. These findings can be valuable for farmers and agricultural practitioners in optimizing crop productivity and ensuring sustainable agricultural practices.

**Key words:** Weed Control Measures, Nutrient Management, Uptake, Content, Nutrient Use Efficiency, Fennel, RDF.

### Introduction

Fennel, also known as Saunf, is a highly important seed spice crop that belongs to the Umbelliferae (Apiaceae) family. It is believed to have originated in the Mediterranean and

Southern Europe. The fennel plant has a pleasant aroma and its all parts, including leaves, stalks, bulbs, and seeds, are edible. Fennel seeds are known for their fragrant scent and aromatic flavor, which is due to the presence of volatile oils like Anethole and Fenchone. These seeds are commonly used in various culinary preparations, such as meals, meat and fish dishes, ice cream, alcoholic beverages, and herb blends. They are also frequently used in the making of candies, soups, sauces, pastries, pickles, liquors, and bakery dishes. The seeds have approximately 9.5% protein, 10.0% fat, 42.3% carbohydrates, 18.5% fiber, and 13.4% minerals. The amount of volatile oil in the seeds can range from 0.7% to 6.0%, depending on the specific genotypes or botanical types.

In India, fennel is mostly produced in Gujarat, Rajasthan, Madhya Pradesh, Maharashtra, Haryana, Uttar Pradesh, and Punjab. In Rajasthan fennel cultivated on 26.25 thousand hectare area and produces 25.62 thousand tones with an average yield of 976 kg ha<sup>-1</sup> (Spice Board, India, 2019-20). It is primarily grown in the districts of Tonk, Pali, Sirohi and Jodhpur and to a limited extent in Bharatpur, Bikaner, Kota, Jaipur and Ajmer.

Fennel is a valuable commercial cash crop traditionally grown as *rabi* season crop in dry and semi-arid areas. It is a slow growing plant that takes longer time to germinate and leads heavy weed infestation in the initial days of growth. Looking to the importance of the crop the average productivity is low, so the efforts have made to enhance the productivity of the fennel by the management of agronomic practices and fertilizer. Macro nutrients such as N, P and K are essential to all crops. Nitrogen is the most important element that limits crop growth and yield. Most of the nitrogen in plants is present in organic form: nucleic acid, hormones, membrane component, coenzymes and pigment. Phosphorus is an essential component of the genetic information system, the energy transfer compounds (ATP and other nuclei-proteins), cell membranes and phospho-proteins. Potassium improves droughttolerance and water uptake in plants. Hence, balanced crop nutrition and good weed control practices is a key factor to achieving higher crop yields. Keeping the above facts in view, the study was conducted for knowing the suitable weed control measure and best nutrient management for maximizing nutrient content, uptake and nutrient use efficiency of fennel.

## **Materials and Methods**

A field experiment was conducted at Instructional farm, College of Agriculture, Swami Keshwanand Rajasthan Agricultural University, Bikaner during *Rabi* season 2021-22 which is situated at 28.01<sup>0</sup> N latitude and 73.22<sup>0</sup> E longitude and altitude of 234.70 meters above mean sea level. The region falls in Agro-climatic zone IC (Hyper Arid Partially Irrigated North Western Plain Zone) of Rajasthan. The soil of experimental site was loamy sand having pH 8.5, low organic carbon (0.18%), low available N (121.4 kg ha<sup>-1</sup>), medium P (19.08 kg ha<sup>-1</sup>) and low K (191.42 kg ha<sup>-1</sup>). The experiment was laid out in factorial randomized block design with three replications. The treatments comprising combinations of 16 treatments comprising four nutrient management treatments, viz. control (no fertilizer applied), 75 % RDF, 100 % RDF and 125 % RDF and four weed control measures, viz. weed free, pendimethalin @ 0.75 kg ha<sup>-1</sup> (PE), oxyfluorfen @ 50g ha<sup>-1</sup> (POE at 25 DAS) and weedy check. Fennel crop variety RF-143 was used as a test crop. Herbicides were sprayed by using knapsack sprayer at pre emergence (PE) and post emergence of both weed and crop. Urea, SSP and MOP was used as source of N, P and K, respectively. The nitrogen content was estimated by Nessler's reagent colorimeter method (Snell and Snell, 1959), Phosphorus by Ammonium 'vanadomolybdatephosphate' yellow color method (Richards, 1954) and potassium by Flame photometer method (Jackson, 1959). The nutrient uptake was calculated from N, P and K content in seed and straw at harvest using following equation.

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = \frac{\text{Nutrient content in seed (\%)} \times \text{Seed yield} + \text{Nutrient content in straw (\%)} \times \text{Straw yield}}{100}$$

The nutrient use efficiency was computed by using the following formula:

$$\text{NUE (kg kg}^{-1}\text{)} = \frac{\text{Grain yield of fertilizer plot} - \text{Grain yield of unfertilized plot}}{\text{Quantity of applied nutrient}}$$

## **Result and Discussion**

### **Effect of Weed Control Measures**

Nutrient content in seed and straw and their total uptake by crop were significantly influenced by different weed control measure. The lowest content of N, P and K in seed and straw was found in weedy check. Whereas, maximum content of N, P and K was obtained in weed free treatment followed by pendimethalin @ 0.75 kg ha<sup>-1</sup> (PE). Higher content of nutrients in crop ascribed great availability of nutrients in soil, and less weed population. It is

clear from the data (Table 1-3), that minimum uptake of N, P and K was observed under weedy check. Data further revealed that weed free recorded the significantly highest uptake of 40.50 kg N, 10.72 kg P and 58 kg K ha<sup>-1</sup>. Pendimethalin @ 0.75 kg ha<sup>-1</sup> (PE) and oxyfluorfen @ 50 g ha<sup>-1</sup> (POE at 25 DAS) also recorded more N, P and K uptake compared to weedy check. These treatments provided weed free environment to crop at early growth stage, wherein the major portion of the basal dose of fertilizer was applied to soil was available to crop in contrast to weedy check. Similarly, at later stages, the applied nutrients under weedy check were utilized by weeds due to their greater competition and better root system. More availability of nutrients to crop under weed free condition under superior treatments increased their content in the plants, which resulted in higher crop dry matter and yields. Similar findings have also reported by Mehriya *et al.* (2008) and as well as Yadav and Sharma (2004) in cumin, Yadav *et al.* (2015) in coriander and Choudhary *et al.* (2022) in fennel.

### **Effect of Nutrient Management**

A significant increase in N, P and K content in seed and straw and their total uptake by crop recorded with the application of 100 % RDF over control but was found at par with 125 % RDF (Table 1-3). Significant improvement in uptake of nitrogen, phosphorus and potassium might be attributed to their respective higher content in seed and straw. Data in table 4 indicated that increasing level of fertility affects the NUE. Significantly, highest N P K use efficiency was obtained with 100 % RDF 3.18, 7.90 and 15.81 kg kg<sup>-1</sup>. Similar results have also been reported by Naher *et al.* (2016) in fenugreek, Javiya *et al.* (2017) in coriander, Mevada *et al.* (2017) and Kalasare *et al.* (2021) in fennel.

### **Conclusion**

The study suggests that weed control measures and proper nutrient management are crucial for achieving high nutrient content and uptake in fennel crops. The weed free treatment and the application of pendimethalin at a rate of 0.75 kg ha<sup>-1</sup> were found to be effective in controlling weeds and improving nutrient levels. Moreover, the use of 100% recommended dose of fertilizer was found to be the most effective nutrient management strategy, resulting in superior nutrient content, uptake, and nutrient use efficiency. These findings highlight the importance of integrated weed and nutrient management practices for optimizing crop productivity and quality.

### **ACKNOWLEDGEMENT**

I express my gratitude to my advisor Dr. Ramesh Chand Bairwa for constant support, guidance and for his valuable suggestions for improving the quality of this Research work and to all the faculty members of Department of Agronomy, SKRAU, Bikaner, Rajasthan India for providing necessary facilities, for their cooperation, encouragement and support.

## References

- Choudhary, M., Shivran, A., Bana, R. C., Choudhary, S., Kakraliya, M. and Lakhran, H. 2022. Influence of weed management practices under different plant rectangularity on fennel (*Foeniculum vulgare* Mill.) yield performance and nutrient effectiveness with special reference to semi-arid ecologies of Rajasthan. *Agricultural Mechanization in Asia* **53**(2): 5525-5533.
- Jackson, M. L. 1959. Soil chemical analysis, *Journal of Plant Nutrition and Soil Science* **85**(3): 251-252.
- Javiya, P. P., Solanki, J. N., Kaneria, S. C. and Rupareliya, V. V. 2017. Response of coriander (*Coriandrum sativum* L.) to nitrogen and phosphorus in South Saurashtra condition, *International Journal of Pure and Applied Bioscience* **5**(4): 860-866.
- Kalasare, R. S., Gaikwad, D. J., Gaikwad, P. P., Maitra, S., Mahapatra, A. and Yadav, M. K. 2021. Studies on interspersed nutrient management on uptake of nutrients seeds, stover and soil biota status of fennel. *Biological forum-An International Journal* **13**(3a): 760-764.
- Mehriya, M.L., Yadav, R.S., Jangir, R.P. and Poonia, B.L. 2008. Effect of different weed management practices on weeds and yield of cumin. *Annals of Arid Zone* **47**(2): 139-144
- Mevada, K. D., Gamar, P., Ombase, K. C., Bhadane, R. S., and Patel, P. D. 2017. Influence of integrated nutrient management on yield, quality and nutrient status of drilled rabi fennel (*Foeniculum vulgare* Mill.). *Journal of Pure and Applied Microbiology* **11**(3): 1559-1565.

- Naher, M. S., Fahim, A. H. F. and Wadud, M. A. 2016. Crop productivity and nutrient uptake of fenugreek (*Trigonella foenum-graecum* L.) as influenced by integrated nutrient management. *SAARC Journal of Agriculture* **14**(1): 71-79.
- Nalini, N., Srinivas, A., Ramprakash, T., & Rao, V. P. (2017). Weed Management in Ajwain (*Trachyspermum ammi*). *International journal of pure and applied bioscience* **5**(4): 932-939.
- Richard, L. A. (1954). Diagnosis and improvement of saline and alkaline soils. Handbook No. 60. *US Department of Agriculture*
- Sharma, N. and Kumar, A. 2022. Effect of integrated nutrient management on growth and yield of fennel (*Foeniculum vulgare*L.). *Indian Journal of Agricultural Sciences***92**(1): 00-00.
- Snell, F.D. and Snell, C.T. 1959. Colorimetric methods of analysis, *The Journal of Physical Chemistry***52**(7): 1265-1266.
- Yadav, S. S., Choudhary, I., Yadav, L. R. and Sharma, O. P. 2015. Weed management in coriander (*Coriandrum sativum* L.) at varying levels of nitrogen. *Journal of Spices and Aromatic Crops* **25**(1):18-25
- Yadav, S.S. and Sharma, O.P. (2004). Growth analysis in cumin (*Cuminum cyminum* L.) under different weed control methods and nitrogen levels. *Journal of Spices and Aromatic Crops***13**(2): 92–96.

**Table 1: Effect of Weed Control Measures and Nutrient Management on Nitrogen Content and Uptake by Crop**

Treatment	Nitrogen Content (%)		Total N Uptake (Kg ha <sup>-1</sup> )
	Seed	Straw	
<b>Nutrient management</b>			
Control	1.19	0.562	21.79
75% RDF	1.31	0.611	29.19
100 % RDF	1.34	0.651	34.18
125 % RDF	1.35	0.653	35.56
SEm±	0.02	0.008	0.46
CD (p=0.05)	0.05	0.023	1.34
<b>Weed control measures</b>			
Weed free	1.39	0.672	40.50

Pendimethelin 750g ha <sup>-1</sup> PE	1.33	0.630	32.57
Oxyfluorfan50g ha <sup>-1</sup> PoE at 25 DAS	1.31	0.626	30.39
Weedy check	1.16	0.549	17.26
SEm±	0.02	0.008	0.46
CD (p=0.05)	0.05	0.023	1.34

**Table 2: Effect of Weed Control Measures and Nutrient Management on Phosphorus Content and Uptake by Crop**

Treatment	Phosphorus Content (%)		Total P Uptake (Kg ha <sup>-1</sup> )
	Seed	Straw	
<b>Nutrient management</b>			
Control	0.404	0.157	5.60
75% RDF	0.440	0.177	7.63
100 % RDF	0.464	0.190	9.11
125 % RDF	0.469	0.192	9.52
SEm±	0.006	0.002	0.14
CD (p=0.05)	0.018	0.007	0.42
<b>Weed control measures</b>			
Weed free	0.472	0.193	10.72
Pendimethalin 750g ha <sup>-1</sup> PE	0.468	0.191	8.60
Oxyfluorfan50g ha <sup>-1</sup> PoE at 25 DAS	0.444	0.179	8.03
Weedy check	0.393	0.152	4.52
SEm±	0.006	0.002	0.14
CD (p=0.05)	0.018	0.007	0.42


**Table 3: Effect of Weed Control Measures and Nutrient Management on Potassium Content and Uptake by Crop**

Treatment	Potassium Content (%)		Total K Uptake (kg ha <sup>-1</sup> )
	Seed	Straw	
<b>Nutrient management</b>			
Control	0.520	1.38	30.02
75% RDF	0.587	1.54	41.43
100 % RDF	0.602	1.64	49.04
125 % RDF	0.608	1.66	51.22
SEm±	0.009	0.02	1.03
CD (p=0.05)	0.026	0.06	2.99
<b>Weed control measures</b>			

Weed free	0.632	1.67	58.00
Pendimethalin 750g ha <sup>-1</sup> PE	0.596	1.65	46.32
Oxyfluorfan50g ha <sup>-1</sup> PoE at 25 DAS	0.588	1.55	43.45
Weedy check	0.502	1.34	23.93
SEm±	0.009	0.02	1.03
CD (p=0.05)	0.026	0.06	2.99

**Table 4: Effect of Weed Control Measures and Nutrient Management on Nutrient Use Efficiency of Crop**

Treatment	Nitrogen Use Efficiency	Phosphorus Use Efficiency	Potassium Use Efficiency
<b>Nutrient management</b>			
Control	0.00	0.00	0.00
75% RDF	2.85	6.41	12.82
100 % RDF	3.51	7.90	15.81
125 % RDF	3.18	7.16	14.32
SEm±	-	-	-
CD (p=0.05)	-	-	-



**Fig. 1 Effect of Nutrient Management and Weed Control Measures on Total Uptake**