

Effectiveness of Proximal Massage Versus Palm Fisting Exercise on Reducing the Risk of Thrombophlebitis among IV Cannulated Patients Receiving Chemotherapy

Abstract

Introduction

Peripheral venous catheters (PVCs) are frequently used for vascular access cannulation, which is thought to be the most frequent invasive treatment carried out on hospitalized patients. Chemotherapy, which uses some "Cytotoxic" medications, is a common systemic approach to treating cancer. One known side effect of some intravenous chemotherapy administered peripherally is venous irritation, which can result in phlebitis and thrombophlebitis due to the drug's chemical actions on the vein wall. Hands ~~exercise~~ exercise and massage are the ~~simplest way~~ simplest ways to improve blood's ability to circulate through the hands. The muscles of the hands and ~~the surrounding~~ surrounding blood vessels ~~will dilate~~ will dilate to allow more of oxygenated blood to flow

Methods

Seventy-five ~~chemotherapy patients~~ chemotherapy patients from were recruited and assigned randomly to group I (n=25) and group II (n= 25) and group III (n= 25) ~~).~~. The researcher provided massage on the proximal area for group I and group II patients performed palm fisting exercise under the supervision of researcher by squeezing the soft ball where as group III received only routine care. Results: Post-test ~~was carried~~ was carried out to assess the risk of ~~thrombophlebitis in~~ thrombophlebitis in Group I, Group II and Group III using VIP scale after 24 hours, 48 hours and 72 hours of cannulation. The study findings revealed that on day 1 in the Group I, ~~majority, majority~~ majority (60%) ~~of~~ of the samples were with Grade 1 level of thrombophlebitis. While in the, Group II (56%) Group ~~III~~ III (64%) of the samples had

Comment [M1]: Change the title to Background
Try make it more concise
Add the purpose or aim of you study

Grade 2 level of thrombophlebitis and 34% of the samples had Grade 1 level of thrombophlebitis. ~~Where as~~Whereas on day ~~3 in~~3 in the Group I, ~~most~~ (most (88%) ~~of~~ of the samples were with Grade 0 level of thrombophlebitis. While in the, Group II ~~32%~~32% ~~of~~ of the samples had Grade 0 level of thrombophlebitis where as in Group III only 16% of the samples had Grade ~~0 level~~0 level of thrombophlebitis. There was statistically very high significant difference was observed in thrombophlebitis scores in all 3 ~~days~~among days among samples in 3 groups. (p value is =0.001). **Conclusion:** Proximal massage and palm fisting exercises are cost-effective, safe, complementary, non - pharmacological intervention for prevention ~~of thrombophlebitis~~of thrombophlebitis which can be performed by nursing ~~personnel after~~personnel after cannulation.

~~—~~**Keywords:** Chemotherapy; IV Cannulated Patients; Palm fisting exercises; Proximal massage ;Thrombophlebitis.

Proximal massage

Palm fisting exercises

Thrombophlebitis

IV Cannulated Patients

Chemotherapy

Introduction

Chemotherapy, which uses "Cytotoxic" medications, is a common systemic approach to treat cancer¹. Any chemotherapy that lasts longer than 24 hours acts as a very potent intravenous stimulant, typically resulting in "Phlebitis," which then results in the loss of superficial veins.² Between 27% and 70% of all patients undergoing IV therapy experience thrombophlebitis, which is a typical side effect of the treatment.³

Pain, erythema, swelling, and visible thrombosis of the cannulated vein are the symptoms of thrombophlebitis. A study finds that 67.2 percent of cannula removals for IV infusions are caused by problems, necessitating the removal of the cannula and its replacement.⁴ However, the complete prevention of thrombophlebitis incidence due to chemotherapy administration is difficult. There are some interventions which may help to reduce the incidence of thrombophlebitis.⁵ Chemotherapy-related infusion ~~phlebitis is~~ phlebitis is a frequent and severe consequence.⁶ Phlebitis is an inflammatory reaction to chemotherapy medications administered intravenously that may linger for weeks or months.⁷ Phlebitis⁷ Phlebitis caused by chemotherapy has been reported to occur 70% of the time,⁸ ~~which~~⁸ which raises the risk of thrombophlebitis and embolism danger and jeopardizes patient health.⁹ ~~However~~⁹ However, by the standards of the Intravenous Nurses Society, phlebitis should only occur at a rate of 5% or less in every society. Phlebitis incidence ranges from 0.5 percent to 59.1 percent at the moment.¹⁰ ~~The~~¹⁰ The peripheral venous catheter is a common and necessary intravenous device that is commonly used in medical procedures¹¹ and is an essential component of professional nursing practise in all healthcare facilities.¹²

Even when nurses administer chemotherapy with care, all difficulties, including drug extravasations into surrounding tissues, cannot always be prevented.¹³ The use of a peripheral IV catheter (PIVC), the length of the procedure, and the type of therapy are the main variables that affect how well peripheral IV chemotherapy is administered.¹⁴ Nurses

~~should~~Nurses should also be aware of the dynamic, evidence-based practice guidelines for the prevention, management, and treatment of local and systemic problems.¹⁵ ~~However~~¹⁵

However, it is challenging to completely prevent the occurrence of thrombophlebitis as a result of chemotherapy administration. Several interventions have the potential to lower the prevalence of thrombophlebitis.⁵

According to the findings of a meta-analysis, any type of upper limb movement can increase the blood flow rate during venous catheterization and lower the incidence of thrombus in patients with PICC. Although there are many distinct types of upper limb motions, they can all help to speed up the blood flow, increase the limb's flow rate on the catheterization side and encourage lymphatic and venous circulation to decrease the likelihood of thrombosis.¹⁶

Hand exercises and massage are two of the most basic methods for increasing circulation flow through the hands. While performing hand exercises, the muscles in the hands and the surrounding blood vessels will relax, allowing more oxygenated blood to flow through.¹⁶

~~One~~¹⁶ One of the most widely publicized advantages of massage is an increase in blood flow to muscular tissue. Improved blood circulation is thought to increase the amount of oxygen and other nutrients delivered to the muscular tissue.¹⁷

Considering the importance of phlebitis induced by intravenous injection of chemotherapy drugs and the consequence of imposing additional treatment costs for the patient and the health system, the present study was designed to compare the effectiveness of proximal massage versus palm fisting exercise on reducing the risk of thrombophlebitis among IV cannulated patients receiving chemotherapy in selected hospital in Mangalore.

Methods

Study design, duration and ethical considerations

A quasi-experimental repeated treatment time series design was used to accomplish the aim to evaluate the effectiveness of proximal massage versus palm fisting exercise on reducing the risk of thrombophlebitis among IV cannulated patients receiving ~~chemotherapy~~ chemotherapy.

Sample size was calculated based on Power ~~analysis with~~ analysis with a power of 0.80, level of significance 0.05 ~~and the~~ and the effect size was ~~0.5~~ 0.5. Recommended sample size was 25 sample in each group. Using purposive sampling technique 75 chemotherapy patients newly cannulated with an IV cannula in the upper extremities, conscious and admitted for more than 3 days were recruited and assigned randomly to group I (n=25) and group II (n=25) and group III (n=25).

Research Design:

The research design selected for this study in quasi experimental post-test only repeated treatment measure time series design

Comment [M2]: Add under title study design, duration and ethical considerations

Variables Under The Study :

Independent variable:

- Proximal massage
- Palm fisting exercise

Dependent Variable :

Risk of thrombophlebitis

Demographic variables : Age, gender, previous history of smoking and alcohol, current alcoholic status, and current smoking status.

Clinical variables : cannula size, number of needle prick, previous hospitalization previous history of thrombophlebitis ,anti-coagulant therapy

Setting :Study was conducted in oncology wards of selected hospitals at Mangalore.

Comment [M3]: Replace this section by "outcome measures " and for each outcome add details about the test or equipment used

Data Collection Method Participants and randomization

Comment [M4]: Move this section before outcome measures

Prior to the data collection formal permission was obtained from hospital ~~authority~~. authority. An ethical clearance was obtained from institutional ethical committee ~~board~~. board. Subject were selected according to inclusion criteria and randomly assigned to Group I, Group II and Group III. . Informed consent was obtained from subject after explaining the purpose of the study. Base line information was collected using socio demographic and clinical proforma.

Comment [M5]: What are the inclusion and exclusion criteria, how did you apply randomization

Intervention

Group I-I: Gentle massage was provided on the proximal area about two ~~centimeters~~centimetres away from cannulation site in the direction of blood flow that allows the blood to move in one direction after pouring a drop of coconut oil on the site for 5 minutes about 20 strokes per minute using palmer surface of the fingers and given once a day on day of cannulation and thrice a day for next 2 days with 3hours interval .

Group II-II: Palm fistng exercise ~~was demonstrated~~was demonstrated by the researcher and asked patients to perform exercise under the supervision of researcher by squeezing the soft ball in the cannulated hand for 5 minutes about 20 times per minute once a day on day of cannulation and thrice a day for next 2 days with 3hours ~~interval~~interval.

Group III-III: Routine care ~~was given~~was given by the ward staff which includes changing the cannula plaster and observing for thrombophlebitis

Post-test ~~was carried~~was carried out to assess the risk of thrombophlebitis in Group I, Group II and Group III using VIP scale after 24 hours, 48 hours and 72 hours of cannulation.

(Figure 1 ~~Flow~~Flow Diagram – Methodology)

Comment [M6]: There is no diagram in the manuscript

UNDER PEER REVIEW

VIP scale is a standardized zero to five points rating scale and used to assess the phlebitis and thrombophlebitis. VIP scale was developed by Andrew Jackson (1998).

The **VIP scale provides a score from 0 to 5**, in ascending order of severity of inflammation.

Score 0 - the insertion site appears healthy and there are no signs of phlebitis.

Score 1 - one of the following signs is evident: slight pain or slight redness near IV insertion site. These are possible early signs of phlebitis.

Score 2 - two of the following signs are evident: pain at IV site, redness or swelling.

Score 3 - all of the following signs are evident: pain along the path of the cannula, redness around the insertion site and swelling.

Score 4 - all of the following signs are evident and extensive: pain along the path of the cannula, redness around the insertion site, swelling, palpable venous cord.

Score 5 - all of the following signs are evident and extensive: pain along the path of the cannula, redness around the insertion site, swelling, palpable venous cord, pyrexia.

Comment [M7]: Write this in the outcome measures section for each outcome write the used test or tool demonstrate its validity scoring and interpretation

Ethical consideration

Institutional review board approval was obtained from Institutional Ethics Committee. Permission has been obtained from concerned authority to collect data. Informed consent was obtained from all the participants of the study. The participants were informed that their participation in the study was entirely voluntary and they may withdraw from the study anytime.

Comment [M8]: Add under title study design, duration and ethical considerations

Data Analysis

SPSS version ~~20~~ was 20 was used for the statistical analysis of the data. Descriptive statistical parameters of mean, standard deviation and percentage were calculated for socio-

demographic and clinical information ~~and repeated~~ repeated measures ANOVA test for intra group comparison.

Results

Descriptive characteristics

Equal percentage of the respondents in Group 1 (28%) were in the age group of 25-35 and >55, and Group 3 (36%) were in the age group of 45-55 and >55 where as the highest percentage of the respondents in Group 2 (44%) were in the age group of 45-55. Majority of the respondents in Group 1 (76%), Group 2 (76%) and Group 3 (64%) were males. The calculated p value reveals that the 3 group didn't vary in demographic characteristics except for past history of smoking and alcohol.. (Table 1).

Table 1. Analysis of Demographic Variables

	Group I(N=25)		Group II(N=25)		Group III(N=25)		P value
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage	
	(f)	(%)	(f)	(%)	(f)	(%)	
Agein Years							
25-35	7	28%	4	16%	3	12%	.3386
35-45	5	20%	5	20%	4	16%	
45-55	6	24%	11	44%	9	36%	
>55	7	28%	5	20%	9	36%	
Gender							
Male	19	76%	19	76%	16	64%	.551
Female	6	24%	6	24%	9	36%	
History of Alcohol							
Yes	0	0%	7	28%	6	24%	0.017
No	25	100%	18	72%	19	76%	
History of Smoking							
Yes	0	0%	6	24%	6	24%	0.009
No	25	100%	19	76%	19	76%	
CurrentAlcoholic Status							
Yes	0	0%	0	0%	0	0%	1
No	25	100%	25	100%	25	100%	
Current SmokingStatus							
Yes	0	0%	0	0%	0	0%	1
No	25	100%	25	100%	25	100%	
Body Mass Index							

Formatted: Centered, Indent: Left: -0.11", Hanging: 0.04", Right: -0.08", Space Before: 0 pt

Formatted: Centered, Indent: Left: -0.07", First line: 0.06", Right: 0.32"

<18.5:- underweight	8	32%	3	12%	1	4%	1
18.5-24.9:- Normal	15	60%	18	72%	23	92%	
25-29.9:- Preobesity	2	8%	2	8%	1	4%	
30-34.9 :- ObesityI	0	0%	2	8%	0	0%	
>40:- ObesityIII	0	0%	0	0%	0	0%	

Analysis of clinical proforma showed that highest percentage of the respondents shows that majority of the respondents duration of ~~illness in~~ illness in group I Group I (60%) and Group ~~3II~~ (68%) from 6 months - 1 year, where ~~as in~~ as in group Group 2 II (48%) duration of ~~illness was~~ illness less ~~less than 6~~ than 6 months. Majority of the respondents in group Group 1 I (80%), ~~Group group 2 II~~ (68%) and and group Group 3 III (84%) had ~~single~~ single needle prick for cannulation. Majority of the respondents in group Group 1 I (52%) and ~~Group and Group 2 II~~ (40%) had the frequency of Infusion more than thrice and majority of the respondents in Group group 3 III (52%) had the frequency of ~~infusion~~ infusion more than ~~twice~~ twice. Majority of the respondents in group Group 1 (68%), group Group 2 (72%) and Group group 3 (64%) do not have the history of thrombophlebitis. The calculated p value reveals that the 3 ~~groups didn't~~ groups didn't vary in clinical variables (Table 2)

Table 2 Analysis of Clinical Variables

	Group I(N=25)		Group II(N=25)		Group III(N=25)		P value
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage	
	(f)	(%)	(f)	(%)	(f)	(%)	
Current Diagnosis							
Breastcancer	0	0%	3	12%	3	12%	0.19 1
Thyroidcancer	0	0%	1	4%	0	0%	
Leukaemia	4	16%	5	20%	5	20%	
Laryngeal cancer	1	4%	1	4%	1	4%	
Prostatecancer	5	20%	6	24%	5	20%	
Lungcancer	10	40%	3	12%	5	20%	
Oral Cancer	5	20%	6	24%	6	24%	
Duration of Illness							
<6months	3	12%	12	48%	3	12%	0.80 3
6months-1year	15	60%	7	28%	17	68%	
1year- 2year	6	24%	1	4%	5	20%	
>2years	1	4%	5	20%	0	0%	
Cannula size							
18G	0	0%	0	0%	0	0%	
20G	25	100%	22	88%	20	80%	
22G	0	0%	3	12%	5	20%	
Site of Cannulation							
a. Brachialvein	0	0%	5	20%	1	4%	.075 7
b. Basalic vein	4	16%	9	36%	3	12%	
c.Cephalic vein	11	44%	7	28%	11	44%	
d. Dorsalmetacarpal	10	40%	4	16%	10	40%	
Frequency of Infusion							
a.Once	3	12%	7	28%	5	20%	0.22 3
b. Twice	5	20%	5	20%	13	52%	
c.Thrice	4	16%	3	12%	6	24%	
d. Morethan thrice	13	52%	10	40%	1	4%	
Number of needle prick							
a.One	20	80%	17	68%	21	84%	.372
b. Two	4	16%	3	12%	4	16%	
c.Three	1	4%	5	20%	0	0%	
Previous Hospitalisation							
a.None	0	0%	0	0%	0	0%	0.46 5
b.Once	11	44%	8	32%	7	28%	

Formatted Table

Formatted: Centered, Indent: Left: -0.07", First line: 0.06", Right: 0.32", Space Before: 0 pt, Line spacing: Multiple 1.15 li

Formatted: Left, Indent: Left: -0.07", Line spacing: Multiple 1.15 li

c. Twice	8	32%	7	28%	9	36%	
d. More than twice	6	24%	5	20%	9	36%	
a. None	0	0%	5	20%	0	0%	
Previous history of Thrombophlebitis							
a. Present	8	32%	7	28%	9	36%	0.83
b. Absent	17	68%	18	72%	16	64%	2

On day 1 in the Group-group I, majority (60%) of the samples were with grade Grade-1 level of thrombophlebitis. While in the, Group-group II (56%) Group-group III (64%) of the samples had grade Grade-2 level of thrombophlebitis and 34% of the samples had Grade 1 level of thrombophlebitis. Whereas on day 3 in the group Group-I, ~~most (88%) of~~ of the samples were with Grade-grade 0 level of thrombophlebitis. While in the, group Group-II 32% of the samples had grade Grade-0 level of thrombophlebitis where as in group Group III only 16% of the samples had grade Grade-0 level of thrombophlebitis. ~~(figure2Figure2)~~
-).




Figure 2. Grading of Risk For Thrombophlebitis in Group I, II And III

From the ~~data~~ ~~it~~ ~~can~~ ~~be~~ ~~concluded~~ ~~that~~ ~~concluded~~ ~~that~~, in group ~~1~~ ~~there~~ ~~there~~ was significant ~~difference~~ ~~difference~~, from ~~Day~~ ~~day~~ ~~1~~ to ~~Day~~ ~~day~~ ~~2~~ (t=3.36, p=0.000) and from ~~Day~~ ~~day~~ ~~2~~ to ~~Day~~ ~~day~~ ~~3~~ (t=2.29, p=0.03) and from ~~Day~~ ~~day~~ ~~1~~ to ~~Day~~ ~~day~~ ~~3~~ (t=2.28, p=0.03). In ~~Group~~ ~~group~~ ~~2~~ ~~2~~, there was significant ~~difference~~ ~~from~~ ~~difference~~ ~~from~~ ~~Day~~ ~~day~~ ~~1~~ to ~~Day~~ ~~day~~ ~~2~~ (t=3.05, p=0.005) ~~and~~ ~~and~~ from ~~Day~~ ~~day~~ ~~2~~ to ~~Day~~ ~~day~~ ~~3~~ (t=5.19, p=0.000) ~~and~~ ~~and~~ ~~Day~~ ~~day~~ ~~1~~ - Day 3. (t=3.05, p=0.005). In Group 3, there was significant ~~difference~~ ~~from~~ ~~difference~~ ~~from~~ ~~Day~~ ~~day~~ ~~1~~ - ~~Day~~ ~~day~~ ~~2~~ (t=5.19, p=0.000) ~~from~~ ~~from~~ Day 2 - Day 3 (t=7.58, p=0.000) ~~and~~ ~~and~~ ~~Day~~ ~~day~~ ~~1~~ - ~~Day~~ ~~day~~ ~~3~~. (t=3.41, p=0.002) ~~0.002~~. (Table 3)

Table 3-3: Effect of proximal massage and palm fisting exercise in reducing the risk of thrombophlebitis in 3 groups

Day	Mean+_SD	Group 1		Mean+_SD	Group 2		Mean+_SD	Group 3	
		t value	p value		t value	p value		t value	p value
Day 1- Day 2	0.72 ± 0.61	3.36	0.000	1.72 ± 0.79	3.05	0.005	1.88 ± 0.78	5.19	0.000
Day 2- Day 3	0.4 ± 0.5	2.29	0.03	1.48 ± 0.77	5.19	0.000	1.6 ± 0.58	7.58	0.000
Day 1- Day 3	0.12 ± 0.33	2.28	0.03	0.88 ± 0.73	3.05	0.005	1.32 ± 0.8	3.41	0.002

To compare the risk of Thrombophlebitis in IV cannulated patients receiving ~~chemotherapy~~ ~~between~~ ~~chemotherapy~~ ~~between~~ ~~Group~~ ~~group~~ ~~I~~, ~~Group~~ ~~group~~ ~~II~~ and ~~Group~~ ~~group~~ ~~III~~, one way ANOVA was computed which showed a ~~high~~ ~~statistical~~ ~~high~~ ~~statistical~~ significance. (p = 0.001). The mean VIP score of the ~~Group~~ ~~group~~ ~~1~~ was ~~lower~~ ~~than~~ ~~lower~~ ~~than~~ the mean VIP scores of the ~~Group~~ ~~group~~ ~~2~~ ~~II~~ and group ~~3~~ ~~III~~. Thus, indicating that proximal massage was effective in reducing the risk of thrombophlebitis in comparison with palm fisting exercises. (Table 4)

Table 4-4: Comparison of risk of thrombophlebitis between 3 groups

	Mean+_SD	F value
Day 1		

Group 1	0.72±_0.61	< 0.0001
Group 2	1.72±_0.79	
Group 3	1.88±_0.78	
Day 2		
Group 1	0.4±_0.5	< 0.00001
Group 2	1.48±_0.77	
Group 3	1.6±_0.58	
Day 3		
Group 1	0.12±_0.33	< 0.00001
Group 2	0.88±_0.73	
Group 3	1.32±_0.8	

Discussion:-

The first objective is to assess the level of phlebitis among IV cannulated patients before and after the palm fisting and proximal massage. Majority of the respondents in Group 1 (76%), Group 2 (76%) and Group 3 (64%) were males. This finding were similar with the other studies where most of patients were males. None of the respondents in group 1 had history of alcohol and smoking and majority of respondents in group 2 and 3 had no history of alcohol and smoking. None of the respondents in group 1, group 2 and 3 had current alcoholic and smoking status where as in other study conducted showed that most 60.0% in group uses alcohol and 86.7% has smoking history.¹⁸ In the present study, majority of samples in all 3 groups did not had previous history of phlebitis . This finding was similar with the study¹⁹ which showed that, very few (19%) of patients were having previous history of phlebitis.

In present study on day 3 in the Group 1, most (88%) of the samples were with Grade 0 level of thrombophlebitis. While in the, Group 2 32% of%

of the samples had ~~Grade-grade~~ 0 level of thrombophlebitis where as in ~~Group-group~~ 3 ~~III~~ only 16% of the samples had ~~Grade-grade~~ 0 level of thrombophlebitis. This study was similar with the study conducted showed 91% of the samples with ~~Grade-grade~~ 0 level of thrombophlebitis. While in the ~~control group~~, 41% of the samples had ~~Grade-grade~~ 0 level of thrombophlebitis and 34% of the samples had Grade 1 level of thrombophlebitis.²⁰ This study was also consistent similar with another study²¹ showed that 90% of the patients had no phlebitis in the post-test.

Comment [M9]: Which group do you mean?
Group III

The second objective is to assess the effectiveness of palm fisting and proximal massage among IV cannulated patients. In the present study the mean ~~VIP-score~~ ~~VIP score~~ of group 4 ~~I~~ was less than the mean VIP score of ~~group2-group II~~ and ~~groupand group3III~~. The findings are consistent with another study ~~where-the~~ ~~where the~~ mean VIP score (1.67) of the control group was higher than the mean VIP score (0.8) of the experimental group.

In the current study there was statistically very high significant difference was observed in thrombophlebitis scores in all 3 ~~days-among~~ ~~days among~~ samples in 3-III groups. (~~p=~~ ~~value is~~ <0.001). The present study findings are consistent ~~with-other~~ ~~with other~~ study where paired “t” value is -1.648 which is significant at 0.05 level.¹⁹ The study finding is also supported by⁵, a study ,among group 60 samples in experimental and control group. Both the groups were compared by using Fisher’s Exact Test ($p=0.027$) at the level of <0.05 for the grades of thrombophlebitis and it was found significant. This indicates proximal massage and palm fisting exercise was effective in reducing the risk of thrombophlebitis¹⁷

REFERENCES

1. DeVita VT, Lawrence TS, Rosenberg SA. Cancer: Principles and Practice of Oncology. 8th ed. Philadelphia: Lippincott; 2008. pp. 330–380
2. Hecker JF. Potential for extending survival of peripheral intravenous infusions. Br. Med. J. 1992;304:619–624.

3. Zingg W, Pittet D. Peripheral venouse catheters: An under-evaluated problem. *Int J Antimicrob Agents* 2009;34:38-42
4. Enes MS, Opitz SP. Phlebitis associated with peripheral intravenous catheters in adults admitted to hospital in the Western Brazilian Amazon. *J SchNurs* 2016;50:261-9.
5. Anjum U, Sharma V, Tamang EL et al. Effect of effectiveness of Palm Fisting Exercise on Occurrence of Thrombophlebitis among the IV Cannulated Patients Receiving Chemotherapy – A Selected Hospital of Delhi. *Int J NursMidwif Res* 2017; 4(3): 61-65
6. Urbanetto JS, Peixoto CG, May TA. Incidencia de flebitis durante el uso y después de la retirada de catéter intravenoso periférico. *Rev. Latino-Am. Enfermagem*. 2016;24: 2746.
7. Mermel LA. Prevention of intravascular catheter-related infections. *Ann. Int. Med.* 2000;132:391–402) doi: 10.1590/1518-8345.0604.2746.
8. Rahmani R. Effect of Topical TNG to Prevent From Phlebitis through Using Venous Catheter on Patients Who Were Under Chemotherapy Treatment [Dissertation] Iran: Bagiyatallah University of Medical Sciences; 2008) (Rahmani R. Effect of Topical TNG to Prevent From Phlebitis through Using Venous Catheter on Patients Who Were Under Chemotherapy Treatment [Dissertation] Iran: Bagiyatallah University of Medical Sciences; 2008
9. Tagalakis V, Kahn SR, Libman M, Blostein M. The epidemiology of peripheral vein infusion thrombophlebitis: a critical review. *Am. J. Med.* 2002;113:146–5
10. Tork-Torabi M, Namnabati M, Allameh Z, Talakoub S. Vancomycin Infusion Methods on Phlebitis Prevention in Children. *Iran. J. Nurs. Midwifery Res.* 2019;24:432–436. doi: 10.4103/ijnmr.IJNMR_149_18

11. Bijayalaxmi B, Urmila A, Prasad PS. Knowledge of staff nurses regarding intravenous catheter related infection working in Orissa. *Nursing Journal of India*, (2010); 101(6): 122–124
12. Arbaee I. Nurses knowledge and practice towards care and maintenance of. *Qualitative Research*, 2016; 1(3): 385–405.
13. Fidalgo JA, Fabregat LG, Cervantes A, Margulies A, Vidall C, Roila F. ESMO Guidelines Working Group. Management of chemotherapy extravasation: ESMO-EONS clinical practice guidelines. *Annals of Oncology*, 2012; 23: 167–174. doi:10.1093/annonc/mds29
14. Centre for Healthcare Related Infection Surveillance and Prevention and Tuberculosis Control. (2013). Guideline: Peripheral intravenous catheter (PIVC) [v.2]. Retrieved from <http://www.cclin-arlin.fr/nosopdf/doc13/364068.pdf>
15. Osti C, Khadka M, Wosti D, Gurung G, Zhao Q. Knowledge and practice towards care and maintenance of peripheral intravenous cannula among nurses in Chitwan Medical College Teaching Hospital, Nepal. *Nurs Open*. 2019 Apr 15;6(3):1006-1012. doi: 10.1002/nop2.288. PMID: 31367425; PMCID: PMC6650670.
16. Li CM, Liang X, Cao SL, Zhou H, Zhang HY. Effect of Physical Intervention on Preventing Upper Limb Venous Thrombosis of Patients with PICC: A Systematic Review and MetaAnalysis of Randomized Trials. *Yangtze Medicine*. 2020; 4: 241-253. <https://doi.org/10.4236/ym.2>
17. Bakhtiar J, Sengupta M. Effect of Proximal Massage and Palm Fisting Exercise in Reducing the risk of Thrombophlebitis among IV cannulated patients in selected hospital, Guwahati, Assam. *Int. J. Nur. Edu. and Research*. 2021; 9(2):202-206. doi: 10.5958/2454-2660.2021.00049

18. Varghese AT, Kt M. Effectiveness of magnesium sulfate with glycerine versus cold compress on patients with peripheral intravenous cannula induced phlebitis. *Asian J Pharm Clin Res* 2018;11:275-8.
19. Daud A, Mohamad F, Noorkhairina S. Incidence of Phlebitis among Adult Patients with Peripheral Intravenous Catheter in an East Coast Hospital Malaysia. *Int J Care Sch* [Internet]. 2018; 1(2):5–8. Available from:<http://journals.iium.edu.my/ijcs/index.php/ijcs/article/view/62/13>
20. Bai RR. Effectiveness of Proximal Massage and Palm Fisting in Reducing the Risk of Thrombophlebitis among Intravenous Cannulated Inpatients. *Indian J of Nurs Sci*. 2022;7(1):14-18.
21. Packialakshmi K, Vidhya T. Effectiveness of warm compress in the management of complications among patients with peripheral catheters. *Int J Pharma Bio Sci* 2017;8:208-13.

UNDER

UNDER PEER REVIEW