

Original Research Article

Efficacy of Fungicides and Herbicides for the management of *Sclerotium rolfsii* of wheat

ABSTRACT

The present research was conducted to investigate and assess the effectiveness of fungicides and herbicides in controlling the disease caused by the fungus *Sclerotium rolfsii* in wheat crops. Different fungicides against *S. rolfsii* were tested in vitro: The efficacy of two systemic fungicides (Tricyclazole 75% W.P. & Carbendazim 50% WP); two non-systemic fungicides (Mancozeb 75% WP & Copper oxychloride 50% WP); one contact fungicides Propineb 70% WP and one combo fungicides (Metalaxyl 8%+Mancozeb 64%) were evaluated at different concentrations (50,150 and 250 ppm) on the development of *S. rolfsii* on Potato dextrose agar (PDA) medium using poisoned food technique (Nene and Thapliyal, 1982). Among the six fungicides, Mancozeb was found best at all the concentrations followed by Tricyclazole at higher concentrations which inhibit the growth of *S. rolfsii*. Another method used for management strategy was herbicide for inhibiting the mycelial growth of *S. Rolfsii*. *In vitro* efficacy of selective herbicides viz. (Metribuzin 70% WP& Oxadiargyl 80% WP), systemic herbicide viz. (Pyrazosulfuron Ethyl 10% WP & Metsulfuron Methyl 20% WP) and one combo herbicides viz. (Bensulfuron Methyl 0.6% + Pretilachlor 6% GR) were tested at their recommended concentrations. Among them metribuzin showed 80.00 % growth inhibition, succeeded by Pyrazosulfuron ethyl (51.85%) and Bensulfuron methyl + Pretilachlor (45.92%).

Introduction

Wheat is an important cereal crop and a staple food for majority of the human population. Presently in the world, wheat is grown over an area of 240.4 m ha with a production of 757.92 mt and a productivity of 3,438 kg ha⁻¹. India stands fourth among wheat producing countries both with respect to area and production. In India, it is grown over an area of 30.71 m ha with a production of 101.20 mt and a productivity of 3,295 kg ha⁻¹ (Anon., 2019). Karnataka is unique in wheat

cultivation where in all three cultivated species, viz., *Triticum aestivum* L., (Bread wheat), *T. durum* (Marconi wheat) and *T. dicoccum* (Khapli, Sadaka or Emmer wheat) are grown in tropical climates characterized by the prevalence of high temperature during the crop growth. *Sclerotium rolfsii*, also known as southern blight or white mold, is a devastating soil-borne fungal pathogen that affects a wide range of crops, including wheat. It can lead to significant yield losses and reduce the overall quality of the crop. To mitigate the impact of this pathogen on wheat production, farmers and researchers explore various management strategies, and chemical control using fungicides and herbicides is one of the common approaches. Efficient management strategies are crucial to combat *Sclerotium rolfsii* and protect wheat crops from its destructive impact. Among the diverse approaches available, the application of fungicides and herbicides has shown promise in mitigating the spread and severity of this fungal pathogen.

The primary objective of this study is to evaluate the efficacy of fungicides and herbicides for the management of *Sclerotium rolfsii* in wheat crops. The research will focus on understanding the impact of these chemical treatments on disease suppression, overall plant health, and crop yield. Additionally, we will investigate the potential ecological implications and safety concerns associated with their use.

To accomplish this, a series of field trials will be conducted over multiple cropping seasons in diverse geographical regions. Various commercial fungicides and herbicides, recommended for fungal and weed control, respectively, will be assessed for their effectiveness against *Sclerotium rolfsii*. Data on disease incidence, severity, and crop yield will be recorded, and statistical analyses will be employed to determine significant differences among treatments. Furthermore, this research aims to address the issue of resistance development in the fungal population against fungicides and explore alternative strategies to minimize resistance risks. In doing so, the study seeks to devise integrated disease

management approaches that encompass cultural practices, biological control agents, and reduced-risk chemicals.

The findings of this study are anticipated to contribute valuable insights to the scientific community, agricultural stakeholders, and farmers alike. Ultimately, the goal is to provide evidence-based recommendations for sustainable and effective management practices against *Sclerotium rolfsii* in wheat crops, ensuring food security and safeguarding agricultural productivity in the face of fungal challenges.

Methodology

The present studies were accomplished at the Department of Plant Pathology, College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.). The efficacy of two systemic fungicides (Tricyclazole 75% W.P. & Carbendazim 50% WP); two non-systemic fungicides (Mancozeb 75% WP & Copper oxychloride 50% WP); one contact fungicides Propineb 70% WP and one combo fungicides (Metalaxyl 8%+Mancozeb 64%) and herbicides viz (Metribuzin 70% WP & Oxadiargyl 80% WP), systemic herbicide viz. (Pyrazosulfuron Ethyl 10% WP & Metsulfuron Methyl 20% WP) and one combo herbicides viz. (Bensulfuron Methyl 0.6% + Pretilachlor 6% GR) were tested at their recommended concentrations.

1. *In vitro* evaluation of fungicides against *S. rolfsii* by poison food technique

Fungicides against *S. rolfsii* were tested in vitro: The efficacy of two systemic fungicides (Tricyclazole 75% W.P. & Carbendazim 50% WP); two non-systemic fungicides (Mancozeb 75% WP & Copper oxychloride 50% WP); one contact fungicides Propineb 70% WP and one combo fungicides (Metalaxyl 8%+Mancozeb 64%) were evaluated at different concentrations (50,150 and 250 ppm) on the development of *S. rolfsii* on Potato dextrose agar (PDA) medium using poisoned food technique (Nene and Thapliyal, 1982).

To get the appropriate fungicide concentration, the required quantity of each fungicide was put individually into sterilized molten and lukewarm potato dextrose agar. The poisoned medium was then put into sterilized Petri plates in a volume of 20 ml. A sterile cork borer was used to cut 5 mm mycelium discs from a

5-day-old culture, and one disc was put in the center of each plate. As a control, a plate without any fungicide was used. For each concentration, three replications were kept. The plates were cultured at 25±1°C temperature and the radial growth was measured when the fungus reached maximal growth in control plates. The fungicide effectiveness was evaluated as a percentage suppression of mycelial growth above control, calculated by using the formula given by Vincent (1947).

$$I = \frac{C - T}{C} \times 100$$

Where I = per cent inhibition

C = growth in control

T = growth in treatment

Table: - 1 *In vitro* evaluation of fungicides used against *S. rolfsii*

S.N.	Name of fungicides	Formulation	Doses (ppm)
1.	Mancozeb	Wettable powder	50, 150, 250
2.	Carbendazim	Wettable powder	50, 150, 250
3.	Propineb	Wettable powder	50, 150, 250
4.	Metalaxyl + Mancozeb	Wettable powder	50, 150, 250
5.	Copper oxychloride	Wettable powder	50, 150, 250
6.	Tricyclazole	Wettable powder	50, 150, 250
7.	Control	-	-

2. *In vitro* evaluation of herbicides against *S. rolfsii* by poison food technique

In vitro efficacy of selective herbicides viz. (Metribuzin 70% WP & Oxadiargyl 80% WP), systemic herbicide viz. (Pyrazosulfuron Ethyl 10% WP & Metsulfuron Methyl 20% WP) and one combo herbicides viz. (Bensulfuron Methyl 0.6% + Pretilachlor 6% GR) were tested at their recommended concentrations. For each treatment, 100 ml PDA was autoclaved in a 250 ml conical flask. At a lukewarm temperature, a specific concentration of herbicide was added to the

medium and well mixed by shaking the flask. In 9 cm Petri plates, 20 ml of this medium was poured. Five mm diameter mycelial disc from a 5-days old pathogen culture were inoculated in the center and incubated at 25±1°C. A suitable control was maintained by growing the pathogen on herbicides free PDA medium. For each treatment, three replications were kept, and the percentage of growth inhibition was determined using the formula below.

$$I = \frac{C - T}{C} \times 100$$

Where, I = Percent inhibition,

C = Colony diameter of the test fungus in Control and

T = Colony diameter of the test fungus in Treatment

Table: - 2 *In vitro* evaluation of herbicides used against *S. rolf sii*

S.N.	Name of herbicides	Formulation	Doses (gm/ lit.)
1.	Bensulfuron methyl + Pretilachlor	Granule	20.00
2.	Oxadiargyl	Wettable powder	0.20
3.	Metsulfuron methyl	Wettable powder	0.008
4.	Pyrazosulfuron ethyl	Wettable powder	4.00
5.	Metribuzin	Wettable powder	0.65
6.	Control	-	-

Result

1. *In-vitro* evaluation of fungicides against *S. rolf sii*: -

To evaluate the efficacy of fungicide against the *S. rolf sii*. In present investigations, six different fungicides, including mancozeb (75 % WP), carbendazim (% WP), propineb (% WP), metalaxyl 4 percent + mancozeb (64 % WP), and copper oxychloride (% WP), on radial growth and inhibition (%) of *S. rolf sii*, were evaluated and the results were presented in Table-3 and illustrated in Plate-: 1.

At 50 ppm, significantly minimum radial growth (38.67mm) and maximum percent inhibition was observed in Mancozeb (57.03%) growth inhibition while other fungicide not effect the growth of *S. rolfsii*.

At 150 ppm, significantly minimum radial growth (0.00mm) and maximum percent inhibition was observed in Mancozeb 100% while other fungicide like Tricyclazole, Metalaxyl+ Mancozeb, Propineb, had least effective as compared to control.

At 250 ppm showed that significant minimum radial growth (0.00mm) and maximum percent inhibition (100%) was observed in Mancozeb while other fungicide like Tricyclazole, Metalaxyl+ Mancozeb, Propineb, had least effective as compared to control. Carbendazim, Copper oxychloride and Tricyclazole not effect the growth of *S. rolfsii*.

Manu *et al.*, (2012) also revealed that 14 fungicides viz., (Tricyclazole, Propiconazole, Difenconazole, Hexaconazole, Carbendazim, Thiophanate methyl, Captan, Chlorothalonil, Mancozeb, Thiram, Hexaconazole 4% + Zineb 68%, Tricyclazole 18% + Mancozeb 62%, Tebuconazole 50% + Trifloxystrobin 25% and Carboxin 37.5% + Thiram 37.5%) were screened *in-vitro* against *S. rolfsii* causing foot of ragi. Hexaconazole, propiconazole, difenoconazole, and combi products such as Avatar, Nativo, and Vitavax power were found to be effective as systemic fungicides, while mancozeb, a contact fungicide, was found to be effective only at higher dosages. Bhuiyan *et al.* (2012) reported that Carbendazim was not effective in inhibiting the radial growth of *S. rolfsii* which support the present finding. Bhatt (2015) also reported that Mancozeb and Captan completely inhibited *S. rolfsii* growth at 125 and 250 ppm. Mancozeb were found very effective in present study also.

S.N.	Treatment	Radial growth (mm) at different Concentrations					
		50 ppm		150 ppm		250 ppm	
		Radial growth (mm) *	Inhibition (%)	Radial growth (mm)	Inhibition (%)	Radial growth (mm)	Inhibition (%)
1.	Mancozeb (75% WP)	38.67	57.03	0.000	100.00	0.000	100.00

2.	Carbendazim (50 % WP)	90.00	0.00	90.00	0.00	90.00	0.00
3.	Propineb (70% WP)	90.00	0.00	74.00	17.77	53.67	40.36
4.	Metalaxyl 4%+ Mancozeb 64% WP	90.00	0.00	71.67	20.36	43.33	51.85
5.	Copper oxychloride (50 % WP)	90.00	0.00	90.00	0.00	90.00	0.00
6.	Tricyclazole (75% WP)	90.00	0.00	42.67	52.58	16.67	81.47
7.	Control	90.00	-	90.00	-	90.00	-
	SEm±	0.025		0.028		0.022	
	C.D.(P=0.05)	0.077		0.086		0.067	
	C.V.	0.528		0.745		0.69	

Table- 3. *In vitro* evaluation of fungicides against *S. rolfsii*.

* Average of 3 replications. 50 PPM


150 PPM

250 PPM

MANCOZEB



CARBENDAZIM



**METALAXYL
+ MANCOZEB**



TRICYCLAZOLE



Plate 1 : Different herbicides in different doses.

2. Efficacy of different Herbicides on *S. rolfsii*.

To evaluate the efficacy of herbicide on *S. rolfsii*. In present study five herbicides, namely Bensulfuron Methyl 0.6 % + Pretilachlor 6 % GR, Oxadiargyl 80 % WP, Metsulfuron Methyl 20% WP, Pyrazosulfuron Ethyl 10 % WP, and Metribuzin 70 % WP, were studied on the mycelial growth of *S. rolfsii in-vitro* by using the poisoned food technique.

Significantly minimum mycelial growth(18.00mm) was recorded in Metribuzin with 80% growth inhibition which was followed by Pyrazosulfuron ethyl (51.85%) and Bensulfuron methyl + Pretilachlor (45.92%). However, the maximum mycelial growth and minimum % inhibition was observed 65.33mm (27.41%) and 81.33mm (9.63%) in Oxadiargyl and Metsulfuron methyl respectively as compare to control.

Rangarani *et al.*, (2017) also evaluated three herbicides and found that Pendimethalin and Quizalofop p-ethyl completely inhibited the pathogen. The growth of *S. rolfsii* was inhibited by 65.6 percent in 2,4-D Na salt. While Sandhya *et al.* used the poisoned food technique to test herbicides such as Glyphosate, 2,4-D Sodium salt, Butachlor, Pretilachlor, Oxadiargyl, Pyrazosulfuron ethyl, Bensulfuron methyl 0.6 %+ Pretilachlor 6 %, Cyhalofop-butyl, Bispyribac sodium, and Ethoxy sulfuron at recommended concentrations Glyphosate, Butachlor, and Pretilachlor all inhibited the growth of *R. solani* to 100%.

Table 4- Efficacy of different Herbicides on *S. rolfsii*.

S.N.	Treatment	Concentrations	Radial growth (mm)*	Inhibition (%)
1.	Bensulfuron methyl 0.6% + Pretilachlor (6.0% GR)	20.00 gm/lit.	48.67	45.92
2.	Oxadiargyl (80% WP)	0.20 gm/lit.	65.33	27.41
3.	Metsulfuron methyl (20% WP)	0.008 gm/lit	81.33	9.63
4.	Pyrazosulfuron ethyl (10% WP)	4.00 gm/lit.	43.33	51.85
5.	Metribuzin (70% WP)	0.65 gm/lit.	18.00	80.00
6.	Control		90.00	-
	SEm±		0.027	-
	C.D. (P= 0.05)		0.085	-
	C.V.		0.816	-

*Average of 3 replications.

Conclusion


Defferent fungicides were tested for the management of *S. rolfsii* among them mancozeb was found to be the best fungicide against *S. rolfsii*, entirely inhibiting radial growth of *S. rolfsii*, followed by Tricyclazole at higher concentrations. Five herbicides were evaluated on the mycelial development of *S. rolfsii*, with metribuzin showing the maximum growth inhibition at 80 %, followed by Pyrazosulfuron ethyl (51.85 %).




**Bensulfuron methyl +
Metsulfuron
methyl**



Pretilachlor



Oxadiargyl



Pyrazosulfuron ethyl



Metribuzin



Control

Plate 2 : Different herbicides

References:

- Archana T. S., Pankaj B. Deore, Jagtap S. D. and Patil B. S. 2018. *In vitro* Evaluation of Fungicides and Bioagents against Root Rot of Chili Caused by *Sclerotium rolfsii* Sacc. International Journal of Pure and Applied Bioscience, 6 (1): 982-986.
- Awasthi, D.P. and Dasgupta, B. 2011. Studies on bio-efficacy of herbicides against *Sclerotium rolfsii* Sacc. causing stem rot of groundnut (*Arachis hypogaea* L.) under *in vitro* condition. Journal of Mycopathological Research, 49(2): 365-366.
- Begum, A., Dadke, M.S., Wagh, S.S., Kuldhar, D.P., Pawar, D.V., Chavan and Thaware, D.S. 2014. *In vitro* evaluation of fungicides and botanicals against stem rot of chilli caused by *Sclerotium rolfsii*. International Journal of Plant Protection, 7(2): 437-440.
- Bhatt, M.N., Sardana, H.R., Singh, D., Srivastava, C. and Ahmad, M. 2015. Evaluation of chemicals and bioagents against *Sclerotium rolfsii* causing southern blight of bell pepper (*Capsicum annuum*). Indian Phytopathology, 68 (1): 97-100.
- Bhuiyan, M.A.H.B., Rahman, M.T. and Bhuiyan, K.A. 2012. *In vitro* screening of fungicides and antagonists against *Sclerotium rolfsii*. African Journal of Biotechnology, 11(82): 14822-14827.
- Das, N.C. Dutta, B.K and Ray, D.C. 2014. Potential of some fungicides on the growth and development of *Sclerotium rolfsii* Sacc. *in vitro*. International Journal of Scientific and Research Publications, 4(12): 1-5.
- Dutta, P. and Das, B.C. 2002. Management of collar rot of tomato by *Trichoderma* spp. and chemicals. Indian Phytopathology, 55(2): 235-237.
- Madhuri, V. and Reddy, P.N. 2013. Compatibility of herbicides with fungicides against *Sclerotium rolfsii*, *Rhizoctonia solani* and *Fusarium udum*. BIOINFOLET-A Quarterly Journal of Life Sciences, 10(3b) :1032-1036.
- Mahato, A., Mondal, B., Dhakre, D.S. and Khatua, D.C. 2014. *In vitro* sensitivity of *Sclerotium rolfsii* towards some fungicides and botanicals. Scholars Academic Journal of Bio Sciences, 2(7): 467-471.
- Manu, T.G., Nagaraja, A., Chetan, S. and Janawad Venayaka, H. 2012. Efficacy of fungicides and biocontrol agents *Sclerotium rolfsii* causing root disease of finger millet under *in vitro* conditions. Global Journal of Biology, Agricultural and Health Science. 1(2):46-50.
- Nene, Y.L. and Thaplial, P.N. 1982. Fungicides in Plant Disease Control. Oxford and IBH Publishing House, New Delhi, p.163
- Pastro, S. and March, G.J. 1999. *In vitro* effect of residual herbicides used in peanut on *Sclerotium rolfsii*. Fitopatologia, 34 (3): 116-121.

- Pranab, D., Das, B.C. and Dutta, P. 2002. Management of collar rot of tomato by *Trichoderma spp.* and chemicals. *Indian Phytopathology*, 5(2): 235- 237.
- Rangarani, A., Rajan, C.P.D., Harathi, P.N., Bhaskar, B. and Sandhya, Y. 2017. Evaluation of fungicides and herbicides on *Sclerotium rolfsii*, incitant of stem rot diseases in groundnut (*Arachis hypogea* L.). *International Journal of Pure & Applied Bioscience*, 5(3): 92-97.
- Sharma, R.K. and Dhruj, I.U., 2018. *In vitro* Evaluation of Some Fungicides against Indian Isolates of *Sclerotium rolfsii* Sacc. [Teleomorph: *Athelia rolfsii* (Curzi) Tu & Kimbrough]. *International Journal of Current Microbiology and Applied Sciences*, 7(11) :1561-1586.
- Singh, S., Prajapati, R.K. and Srivastava, S.S.L. 2005. Efficacy of fungicides against *Sclerotium rolfsii* causing collar rot in lentil, *Journal of farm sciences*, 14(2): 68- 69.
- Tripathi, S.K., Sathie, K.B and Vyas, S.C. 1988. Chemical control of collar rot of pea. *Indian Phytopathology*, 41(1): 143-144.
- Yaqub, F. and Saleem, S. 2006. Effect of fungicides on *in vitro* growth of *Sclerotium rolfsii*. *Pakistan Journal Botany*, 38(3): 881-883.