

**Original Research Article**

**EFFICACY OF FEEDBACK BASED THERAPY FOR DYSPHAGIA IN POST STROKE PATIENTS – QUASI EXPERIMENTAL STUDY.**

**ABSTRACT**

**Background and Purpose**

Nearly one third of stroke survivors experiencing swallowing dysfunction called Dysphagia, in addition to the motor and sensory impairments. Post stroke dysphagia is associated with increased mortality and morbidity due to aspiration pneumonia, and malnutrition. About 11 to 13 % of stroke patients suffer from dysphagia even after 6 months. Feedback can be advocated for enhancing outcomes in dysphagia rehabilitation. However, biofeedback in swallowing therapy is not routinely used. Hence, we tested the functional outcome of cumulative effect of SEMG biofeedback adjunct to swallowing exercises summated with visual, auditory and tactile feedback which engross and stimulates active participation of the patients during the therapy of dysphagia rehabilitation.

**Methods**

This study included 12 stroke patients with swallowing difficulty allocated into control and interventional group with 6 subjects in each. Swallowing function was evaluated using Functional Oral Intake Scale (FOIS), Baseline SEMG of Suprahyoid muscle and Presence of Nasogastric tube. Interventional group received Feedback based therapy along with conventional therapy for 30 to 40 minutes a day for 10 days and control group received conventional therapy for 10 days. The post interventional outcome of FOIS, Average of SEMG of suprahyoid muscle was evaluated.

## **Results**

The Nasogastric tube was removed in all patients of experimental group after intervention and established initiation of oral feeding. The result showed improvement in functional oral intake scale after intervention. The muscle activity noted in experimental group was greater than the control group ( $p < 0.05$ ).

## **Conclusion**

We conclude that Feedback based therapy protocol shows good improvement in post stroke patients with dysphagia. Inclusion of this protocol in therapy may influence standards of care and best practices for post stroke dysphagia patients.

**THE REGISTRATION NUMBER :** This trial was registered in Clinical Trials Registry – India and the trial registration number is CTRI/2020/05/025356

**Key words :** Dysphagia, feedback, FOIS, SEMG, post stroke.

## INTRODUCTION

Stroke is one of the leading cause of disability in India. The incidence rate is 119-145/100,000 based on the recent population based studies (1). In addition to the motor and sensory impairments, dysphagia is a common impairment after stroke. The incidence of dysphagia following stroke was reported between 30% to 50% during recent times (2). Most of stroke survivors will recover from swallowing dysfunction within 7 days, but about 11–13 % will continue to have swallowing dysfunction upto 6 months (3). Post stroke dysphagia increases the risk for dehydration, malnutrition, pulmonary complications, and mortality, which all lead to a poor prognosis and affects the quality of life. As a result of dysphagia, 43%-50% of stroke patients land up with aspiration pneumonia which increases the mortality rate. Only 10% of stroke related deaths are caused by neurological deficits, while 30% of post-stroke deaths are due to pneumonia resulted from dysphagia (4).

Generally, a healthy adult swallows between 800 and 2400 times a day. When timing, coordination, feeling and muscular strength for swallowing are disturbed, this is called dysphagia (5). Impact of Dysphagia include following symptoms, food spillage from lips, taking a long time to finish a meal, poor chewing ability, dry mouth, drooling, nasal regurgitation, food sticking in the throat, poor oral hygiene, coughing and choking, regurgitation, weight loss, repeated chest infections (4). Some patients with severe dysphagia require a nasogastric feeding tube for months, even to years to compensate for adequate nutrition and water intake (6).

Physiotherapeutic interventions of orofacial rehabilitation which aim to maintain and rehabilitate swallowing vary from breath control, postural adjustment, electrical

stimulation, tongue exercises, lip exercises. van de Port IG et al (2007) stated that Biofeedback is often used in physiotherapy to augment skill based therapy. However, biofeedback in swallowing therapy is not routinely used to augment dysphagia rehabilitation. This study was conducted to elaborate the strategies of conventional physiotherapy program for dysphagia rehabilitation in post stroke, as physiotherapist are the part of multidisciplinary team addressing the swallowing dysfunction in stroke population.

The process of squeezing and holding the larynx at the peak of the swallow in Mendelsohn maneuver and rapid jaw opening using surface electromyography (SEMG) biofeedback providing visual feedback from the display as well as tactile feedback by palpating the movement of larynx (the rise, squeeze, and fall of larynx) (7). The use of SEMG biofeedback to facilitate increased effort and duration of attempted swallow (8). Visual Feedback allows the participant to know real time information of current force production and can aid relearning a skill by reducing the errors. Participants are motivate to make more efforts when try modulating the signals thereby indirectly working on their internal physiological events like improving the muscle recruitment. (9)

Generally, the studies were designed to find out the effectiveness of single interventions but none has documented the outcome of combining feedback therapy. while the outcome data from these studies reported improved oral intake in most patients, leaving open questions regarding the functional outcome and improvement in muscle activity which may have occurred. Evident data is extremely limited for dysphagia rehabilitation of conventional therapy augmented with feedback based strategies by biofeedback using SEMG while performing mendelsohn maneuver, effortful swallow and jaw opening to enhance the activity of suprahyoid muscle along with improvement in swallow function. We felt a need to formulate a sole dysphagia rehabilitation protocol in post stroke and to

study functional outcome of cumulative effect of SEMG biofeedback adjunct to swallowing exercises summated with visual, auditory and tactile feedback which engross and stimulates active participation of the patients during the therapy.

UNDER PEER REVIEW

## METHODOLOGY

This Interventional study was approved by the Ethics committee for the students proposals (**REF: CSP/19/NOV/81/369**), Sri Ramachandra Institute of Higher Education and Research. **Inclusion criteria** : Post stroke dysphagia female and male of age 40 to 70 years referred for physiotherapy, Medically stable, Complaints of swallowing and non-oral feeding, Functional oral intake scale less than or equal to two, Mini-mental state examination score of 21 or more than 21. **Exclusion criteria**: Neurological disease other than stroke, Upper gastrointestinal diseases.

## PROCEDURE

This Quasi Experimental study included 12 subjects diagnosed as Stroke with dysphagia from Sri Ramachandra Medical Centre and Hospital. Informed consent was obtained from the participants. The consent was obtained from the care givers if the patient could not sign their written consent. Convenient sampling method was used to allocate the patients who met the inclusion criteria into control and interventional group with 6 subjects in each. Swallowing function was evaluated using a 7 point ordinal Functional Oral intake Scale (FOIS) to document the level of swallowing function in post stroke dysphagia for both the control and experimental groups. Baseline SEMG of Suprahyoid muscle activity was recorded prior to the treatment. Presence of Nasogastric tube was noted.

The skin was prepared by standard skin preparation techniques of cleaning with 70% alcohol (ethanol). SEMG data was recorded using Neurotrac software 4.0 Of VM (Verity Medical Ltd, United Kingdom). Silver chloride adhesive electrodes were used with 10 mm diameter and 20 mm inter electrode distance. The electrodes are placed under the

chin, just lateral to midline, anterior to the hyoid bone thus recording the muscle activity in the submental muscles (m. stylohyoid, m. mylohyoid, m. digastricus). The reference (ground) electrode is positioned away from the muscles of interest. The electrodes are attached to an EMG device using a sensor cable and the signal from the device is registered on a computer using specialized software. On the horizontal axis the time frame is 30s, the vertical axis shows the electrical activity (measured in microvolt) of the submental muscles. During swallowing the larynx is elevated and moved forward by contraction of the supralaryngeal, suprahyoid and submental muscles. SEMG was recorded during dry swallowing.

Patients in both the group received conventional Neuromuscular Electrical stimulation, shaker exercise, postural correction, tongue and lip exercises for 20 minutes, once in a day for 10 days. SEMG was recorded in both the groups after 10 sessions of therapy. Experimental group received feedback based therapy of hyoid lift maneuver, mendelsohn maneuver, effortful swallow and rapid jaw opening along with biofeedback using SEMG in addition to conventional therapy. On completion of the treatment duration of 10 days, the post interventional outcome of Functional Oral Intake Scale (FOIS). Average EMG output from suprahyoid muscles, presence or absence of Nasogastric tube after intervention was evaluated.

## **STATISTICAL ANALYSIS**

Statistical analysis was performed using SPSS version 23.

Within group analysis of SEMG recording of suprahyoid muscles for both control and experimental group was done using Paired t test. Between group analysis of pre and post recordings of SEMG was done using independent t test.

Wilcoxon sign rank test was used to find out the changes associated with FOIS within the groups.

Mann-Whitney u test was used to compare changes associated with FOIS between the control and intervention group.

## RESULTS

**Table 1 : Demographic profile**

<b>Variables</b>	<b>Control</b>	<b>Experimental</b>
<b>Age [mean(SD)]</b>	<b>63(10.08)</b>	<b>61(13.15)</b>
<b>Male</b>	<b>3</b>	<b>1</b>
<b>Female</b>	<b>3</b>	<b>5</b>
<b>Hemorrhagic CVA</b>	<b>4</b>	<b>2</b>
<b>Ischemic CVA</b>	<b>2</b>	<b>3</b>
<b>Lacunar infarcts</b>	<b>-</b>	<b>1</b>

Table 1: The profile of the stroke patients participated in the study.

**Table 2 : Dependency on Nasogastric tube before and after therapy**

Control group			Experimental group		
Patient	Presence of Nasogastric tube		Patient	Presence of Nasogastric tube	
	Pre	Post		Pre	Post
1	Yes	No	1	Yes	No
2	Yes	Yes	2	Yes	No
3	Yes	No	3	Yes	No
4	Yes	No	4	Yes	No
5	Yes	No	5	Yes	No
6	Yes	Yes	6	Yes	No

Table 3: Nasogastric tube was removed for all participants of experimental group after therapy. Whereas in control group 4 were removed from feeding tube and 2 of the participants remained with nasogastric tube even after therapy.

**Table 4 : Functional Oral Intake Scale of post stroke dysphagia**

Variables - Control and Experimental group								
	Within group						Between group	
	Control			Experimental			P <sup>b</sup>	P <sup>c</sup>
Outcome	Pre Mean (SD)	Post Mean (SD)	P <sup>a</sup>	Pre Mean (SD)	Post Mean (SD)	P <sup>a</sup>		
FOIS	1.67 (0.52)	3.67 (0.52)	0.014*	1.67 (0.52)	5 (0.63)	0.023*	1.00	0.01*

Table 3: P<sup>a</sup> - Wilcoxon sign rank test for pre and post analysis within Control and Experimental group.

P<sup>b</sup> - Mann-whitney U test for pre intervention scores between the groups.

P<sup>c</sup> - Mann-whitney U for post intervention scores between the groups

The FOIS was significantly improved in both control and experimental group compared with pre intervention results ( $p < 0.05$ ). The pre intervention values are not statistically different between the groups, whereas ( $p < 0.05$ ) post intervention scores between the groups were statistically different at the end of the treatment period.

**Table 5 : Average SEMG output from suprahyoid muscles in stroke patients and within group results**

<b>Group</b>	<b>Dry swallowing Average output Mean <math>\mu</math>v (SD)</b>	<b>Pre mean (SD)</b>	<b>Post mean (SD)</b>	<b>p value</b>
<b>Control</b>	<b>Right suprahyoid</b>	<b>3.40 (0.64)</b>	<b>8.01 (0.93)</b>	<b>0.000*</b>
	<b>Left suprahyoid</b>	<b>3.48 (0.75)</b>	<b>7.79 (0.94)</b>	<b>0.000*</b>
<b>Experimental</b>	<b>Right suprahyoid</b>	<b>3.44 (0.75)</b>	<b>11.59(2.46)</b>	<b>0.000*</b>
	<b>Left suprahyoid</b>	<b>3.51 (0.72)</b>	<b>12.56(3.97)</b>	<b>0.004*</b>


Table 4: Paired t test was used to analyze the pre and post intervention scores within the groups. The result shows increase in muscle activity in both control and experimental groups following intervention when tested within the groups.

**Table 6 : Average SEMG output from suprahyoid muscles in stroke patients and between group results**


Dry swallowing	Control		Experimental		p value
	Pre Mean (SD)	Post Mean (SD)	Pre Mean (SD)	Post Mean (SD)	
Right suprahyoid	3.4(0.642)	8.00(0.931)	3.443(0.75)	11.59(2.46)	0.01
Left suprahyoid	3.475(0.750)	7.7(0.941)	3.513(0.718)	12.56(3.968)	0.02

Table 5: Independent t test was used to analyze the pre and post intervention scores between two groups. A greater muscle activity was recorded in experimental group compared to control group following intervention. ( $p < 0.005$ ) post intervention scores between the groups were statistically different.


**Graph 1 : Within group analysis of Interventional group for SEMG recording of Right suprahyoid**




**Graph 2 : Within group analysis of Interventional group for SEMG recording of Left suprahyoid**



**Graph 3 : Within group analysis of Control group for SEMG recording of Right suprahyoid**



**Graph 4 : Within group analysis of Control group for SEMG recording of Left suprahyoid**



## DISCUSSION

This study was conducted to elaborate the strategies of conventional physiotherapy program for dysphagia rehabilitation in post stroke, as physiotherapist are the part of multidisciplinary team addressing the swallowing dysfunction in stroke population (25) but the therapeutic strategies of physiotherapy which aim to maintain and rehabilitate swallowing may vary from postural adjustment in sitting, positioning of head and neck to electrical stimulation of pharyngeal muscles. Evident data is extremely limited for dysphagia rehabilitation of conventional therapy augmented with feedback based strategies by biofeedback using SEMG while performing mendelsohn maneuver, effortful swallow and jaw opening to enhance the activity of suprahyoid muscle along with improvement in swallow function.

In this study, we have included 12 participants with swallowing dysfunction who scored less than or equal to 2 in FOIS were allocated to control and experimental group with 6 participants in each group. The demographic profile from table 1 shows more female patients 8 out of 12 participated in this study and the mean(SD) age group in control group was 63(10.08) and in the intervention group was 61(13.15).

### **Dependency on Nasogastric tube :**

At the start of the treatment all 12 patients were fully dependent on Nasogastric tube (FOIS  $\leq$  3). The level was increased by 2 or 3 points on the FOIS in both groups but, the feeding tube was removed in all patients of experimental group after intervention when compared with control group, establishing 100% success rate. This is also explained in a similar study done by Bogaardt et al (2009) where removal of tube established to 75% after the use of surface EMG feedback in chronic dysphagia of stroke patients (17). Whereas in control group, 4 patients became independent of

feeding tube, leaving 2 patients remained on Nasogastric tube for non oral feeding even after treatment.

### **Functional Oral Intake Scale :**

In this study the FOIS of both the group was statistically significant. This may be attributed to the employment of conventional treatment of neuromuscular electrical stimulation of suprahyoid and infrahyoid muscles in both the groups. This result is supported by the recent study of Lee kim et al (2014) which showed NMES combined with traditional dysphagia therapy was a more beneficial treatment for acute/ subacute stroke patients with dysphagia(26).

In a study by Pownall et al (2017), NMES is postulated to improve swallowing function by activating fast twitch muscle fibers ( type 2 motor unit fibers), enhancing the muscle strength of suprahyoid which is responsible for anterior and superior movement of the hyoid (4).

Though the swallowing function was significantly improved in both the groups, our results supported the hypothesis i.e, post interventional results of FOIS was improved significantly in experimental group when compared to control group. Hence, feedback based strategies has a better effect on improving swallowing function than the conventional therapy

### **SEMG of suprahyoid muscles :**

Despite the small sample size, there is a significant difference in post EMG activity of suprahyoid muscles of both right ( $p=0.02$ ) and left side ( $p=0.01$ ) which revealed feedback based therapy had a significant positive effect on increasing the suprahyoid muscle recruitment on dry swallowing.

Zanato et al (2016) reported the highest electrical activity was found in the suprahyoid muscles with values of 9.6 ( $\mu\text{v}$ ) on the left side and 9.5 ( $\mu\text{v}$ ) when compared to temporalis and masseter during the swallow of 10 ml water in normal individuals, since the suprahyoid muscles are the elevators of the hyoid bone which is an essential movement for swallowing (22). Hara et al (2018) found jaw opening exercises resulted in greater activation of the SH muscles and further studies of dysphagia rehabilitation, focusing on jaw-opening movements, may contribute to advancements in dysphagia rehabilitation (27).

A review by Frietas et al (2016) states that EMG biofeedback and other biofeedback methods can be effective in constant self correction. Motor planning and motor control skills are continuously stimulated and beneficial neuronal plasticity is induced. When visual biofeedback is used, changes occurring in cortical activation, highlighting a cortical reorganization (19). A study by Hoon Kim et al (2017) found the combination of the Mendelsohn maneuver with effortful swallowing can produce strong activation of the suprahyoid muscles implies a large recruitment of muscle fibers which directly have a positive influence on the reduction in aspiration. As per the above mentioned studies, suprahyoid muscle activity was improved (28).

## **LIMITATIONS**

The study was done with a small sample size, there was no long-term follow-up with the study patients.

## **CONCLUSION**

We conclude that Feedback based therapy protocol shows better improvement in post stroke patients with dysphagia. Inclusion of this protocol in therapy may influence standards of care and best practices for post stroke dysphagia patients.

## **RECOMMENDATIONS FOR FUTURE STUDY**

1. This study can be replicated on larger samples.
2. A similar study can be conducted on a group with adequate follow-up for 3 months.
3. A similar study can be conducted with assessment of quality of life and clinical indicators at repeated intervals.

## TABLES

### INTERVENTIONAL EXERCISE PROTOCOL

Table 6 : The following exercise are trained along with SEMG biofeedback of each exercises for 5 times once daily for 10 days.

S.no	Exercises	Instructions
1	Hyoid lift maneuver	<ol style="list-style-type: none"><li>1. Multiple small pieces of paper placed on a towel in front of the patient.</li><li>2. Place a straw in mouth and ask to suck the piece of paper on the straw, allowing the paper to get picked up by the tip of the straw.</li><li>3. Keep sucking on the straw ask to carry the straw over to a cup, and stop sucking to release the paper into the cup.</li><li>4. Ask to place all of the pieces of paper into the cup. Gradually increases from 3 to 5 and around 10 pieces.</li></ol>
2	Effortful Swallow	<ol style="list-style-type: none"><li>1. Push tongue up to the roof of the mouth</li><li>2. Swallow "hard" with effort</li><li>3. Pretend as like swallowing a golf ball</li><li>4. Ask to do 3 sessions of this exercise per day</li></ol>
3	Mendelsohn	<ol style="list-style-type: none"><li>1. Begin a regular swallow</li><li>2. Place middle three fingers on Adam's apple. Feel the larynx move in an upward direction and squeeze muscles</li><li>3. Ask to face the screen and instructed the participant to swallow "long and strong" with a squeeze at the peak of the swallow for 3 to 4 seconds</li></ol>
4	Jaw-opening exercise.	Ask to open the jaw to the maximum extent and maintain this position for 10 s.

**Table 7 : CONVENTIONAL THERAPY**

1	Neuromuscular electrical stimulation	Electrical stimulation was given for a period of ten days once daily.  The following are the parameters for stimulation : Current- faradic, Contraction- 90. Placement of electrodes-on either side of midline above the lesser horns of the hyoid bone(anterior neck)
2	Shaker Exercise	Lying on the back on the bed.  Keeping shoulders flat against the bed, and lifting up the head, bringing the chin down to the chest.  Keeping the head lifted for 60 seconds, and then lower the head and rest for 60 seconds.
3	Lip Press	Tightly close upper and lower lip against each other for 5 seconds, repeat 10 times
4	Tongue Push Side to Side	Extend tongue out of mouth as far as possible, patient attempts to move tongue toward the corner of mouth and against the cheek, hold for 5 seconds, repeat 10 times.
5	Tongue Side to Side	Move tongue out of mouth, rotate tongue tip to left side of mouth, hold for 5 seconds, then to right side of mouth, hold for 5 seconds, repeat 10 times.
6	Tongue Push Forward	Open mouth and move tongue such that tongue tip rests against the lower lip, patient attempts to move tongue forward, hold for 5 seconds, repeat 10 times.
7	Tongue Push Up	Extend tongue out of mouth, patient attempts to move tongue upward against the upper lip, hold for 5 seconds, repeat 10 times

## REFERENCES

1. Pandian JD, Sudhan P. Stroke Epidemiology and Stroke Care Services in India. *J Stroke*. 2013;15(3):128.
2. Balamurali K, Sekar D, Thangaraj M, Kumar MA. Dysphagia in Patients with Stroke: A Prospective Study. *Int J Contemp Med Surg Radiol*. 2018 Jul;3(2).
3. Vose A, Nonnenmacher J, Singer ML, González-Fernández M. Dysphagia Management in Acute and Sub-acute Stroke. *Curr Phys Med Rehabil Reports*. 2014 Dec;2(4):197–206.
4. Pownall S, Enderby P, Sproson L. Electrical stimulation for the treatment of dysphagia. In: *Electroceuticals: Advances in Electrostimulation Therapies*. Springer International Publishing; 2017. p. 137–56.
5. Bogaardt HCA (Henri CA. Current aspects of assessment and treatment of dysphagia. [s.n.]; 2009.
6. Huang KL, Liu TY, Huang YC, Leong CP, Lin WC, Pong YP. Functional outcome in acute stroke patients with oropharyngeal dysphagia after swallowing therapy. *J Stroke Cerebrovasc Dis*. 2014;23(10):2547–53.
7. McCullough GH, Kamarunas E, Mann GC, Schmidley JW, Robbins JAA, Crary MA. Effects of Mendelsohn maneuver on measures of swallowing duration post stroke. *Top Stroke Rehabil*. 2012 Jan 1;19(3):234–43.
8. Crary MA, Carnaby GD, Groher ME, Helseth E. Functional benefits of dysphagia therapy using adjunctive sEMG biofeedback. *Dysphagia*. 2004;19(3):160–4.
9. Kim JH, Choi JW, Lee SH, Cho YS. Difference in tongue strength using visual

feedback in healthy adults. *J Phys Ther Sci.* 2017;29(12):2075–6.

10. Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ, Culebras A, et al. An updated definition of stroke for the 21st century: A statement for healthcare professionals from the American heart association/American stroke association. *Stroke.* 2013;44(7):2064–89.
11. Cohen DL, Roffe C, Beavan J, Blackett B, Fairfield CA, Hamdy S, et al. Post-stroke dysphagia: A review and design considerations for future trials. Vol. 11, *International Journal of Stroke.* Blackwell Publishing Ltd; 2016. p. 399–411.
12. M. C, W. J, Chapman-Jay S, Cliffe R, M. S, Oshall M. Electromyography as a Biofeedback Tool for Rehabilitating Swallowing Muscle Function. *Appl EMG Clin Sport Med.* 2012;(March 2014).
13. Langmore SE, Pisegna JM. Efficacy of exercises to rehabilitate dysphagia: A critique of the literature. *Int J Speech Lang Pathol [Internet].* 2015;17(3):222–9. Available from: <http://dx.doi.org/10.3109/17549507.2015.1024171>
14. El-Tamawy MS, Darwish MH, El-Azizi HS, Abdelalim AM, Taha SI. The influence of physical therapy on oropharyngeal dysphagia in acute stroke patients. *Egypt J Neurol Psychiatry Neurosurg.* 2015 Jul 1;52(3):201–5.
15. Lim KB, Lee HJ, Lim SS, Choi YI. Neuromuscular electrical and thermal-tactile stimulation for dysphagia caused by stroke: A randomized controlled trial. *J Rehabil Med.* 2009 Feb;41(3):174–8.

16. Li YC, Wu CY, Hsieh YW, Lin KC, Yao G, Chen CL, et al. The Priming Effects of Mirror Visual Feedback on Bilateral Task Practice: A Randomized Controlled Study. *Occup Ther Int.* 2019;2019.
17. Bogaardt HCA, Grolman W, Fokkens WJ. The use of biofeedback in the treatment of chronic dysphagia in stroke patients. *Folia Phoniatr Logop.* 2009 Sep;61(4):200–5.
18. Gonçalves R. A eficácia da reabilitação em disfagia orofaríngea \*\* Efficacy of rehabilitation in oropharyngeal dysphagia. 2007;123–30.
19. Freitas GS de, Mituuti CT, Furkim AM, Busanello-Stella AR, Stefani FM, Arone MMA da S, et al. Biofeedback eletromiográfico no tratamento das disfunções orofaciais neurogênicas: revisão sistemática de literatura. *Audiol - Commun Res.* 2016;21(0).
20. Konecny P, Elfmark M, Bastlova P, Alacova PG. New Orofacial Physiotherapy of Dysphagia after Stroke. *Int J Phys Med Rehabil.* 2017;05(03).
21. Michael E. Groher P IMichael ACPGDCMPM. Initial Psychometric Assessment of a Functional Oral Intake Scale for Dysphagia in Stroke Patients. 2005;86(August):1516–20.
22. Zanato LE, Chiari BM, Vieira MM, Bommarito S. Study of the electrical activity of muscles : Masseter , temporal and supra-hyoid during swallowing. 2016;3(1):1–4.
23. Belo LR, Menezes DC De. REFERENCE VALUES FOR THE ELECTROMYOGRAPHY OF MUSCLES INVOLVED IN SWALLOWING : 1956;
24. Poorjavad M, Talebian S, Nouredin , Ansari N, Soleymani Z, Phd ; Soleymani Z.

Surface Electromyographic Assessment of Swallowing Function. Vol. 42, Iran J Med Sci March. 2017.

25. Aoki S, Hosomi N, Hirayama J, Nakamori M, Yoshikawa M, Nezu T, et al. The multidisciplinary swallowing team approach decreases pneumonia onset in acute stroke patients. PLoS One. 2016;11(5):1–8.
26. Lee KW, Kim SB, Lee JH, Lee SJ, Ri JW, Park JG. The effect of early neuromuscular electrical stimulation therapy in acute/subacute ischemic stroke patients with dysphagia. Ann Rehabil Med. 2014;38(2):153–9.
27. Hara K, Tohara H, Minakuchi S. Treatment and evaluation of dysphagia rehabilitation especially on suprahyoid muscles as jaw-opening muscles. Jpn Dent Sci Rev [Internet]. 2018;54(4):151–9. Available from: <https://doi.org/10.1016/j.jdsr.2018.06.003>
28. Ji-Hoon Kim<sup>1)</sup>, Young-A Kim<sup>2)</sup>, Hye-Jin Lee<sup>2)</sup>, Keum-Sook Kim<sup>3)</sup>, Seung-Tae Kim<sup>4)</sup>, Tae-Sue Kim<sup>5)</sup> Y-SCh. Effect of the combination of Mendelsohn maneuver and effortful swallowing on aspiration in patients with dysphagia after stroke. 2017;1967–9.

UNDER PEER REVIEW