

Characterization of Native Fluorescent Pseudomonads against Sclerotinia Blight of Brinjal

Abstract

Brinjal (*Solanum melongena* L.) is one of the most common know as one of the rich source of vitamin B1, dietary fibre, manganese, niacin, copper, vitamin B6, folate, potassium, Vitamin K. Brinjal cultivation is affected by various pest and diseases. Sclerotinia Blight is one of the most destructive diseases of brinjal caused by *Sclerotinia sclerotiorum* (Lib) de Bary. Under the present investigation two isolates of Fluorescent Pseudomonads (FLPs) viz., FLP-Brinjal 2020-1 and FLP-Brinjal 2020-2, were evaluated against th *Sclerotinia sclerotiorum* (SCL) under *in vivo* and *in vitro* conditions using dual culture, inverted plate assay. FLP-Brinjal 2020-2 gave maximum inhibition over the control followed by FLP-Brinjal 2020-1 against the pathogen in *in vitro* condition. Experiments conducted in pluck trays indicated that FLP-Brinjal 2020-2 was superior with least sclerotinia blight incidence (20.53%). The identity and diversity among the FLP isolates were examined by sequence analysis of 16s rRNA gene. The gene was amplified by PCR using primers pair, 27 F/1492 R. Amplified DNA products were sequenced, and analysed by BLAST (BLASTn) programme at NCBI database for species identification. Among the 10 FLPs isolates assessed 9 were found to have maximum similarity with *Pseudomonas aeruginosa* including FLP 2020-2 and FLP 2020-1 was found to be *Pseudomonas fluorescence* which is in confirmation with the morphological observations. Significant diversity among the isolates was observed when dendrogram of the sequence was plotted using CLUSTAL W.

Keywords: Antagonist, Brinjal, Biological Control, Fluorescent Pseudomonads, *Sclerotinia sclerotiorum*.

Introduction

The population of India is growing day by day and is expected to reach up to 9.7 billion by 2050 (Godfray et al., 2010; World Population Prospects, United Nation, 2019). Therefore, a significant food supply increase is required to meet the demand of the population. Brinjal is considered as the king of vegetables. It is one of the most common vegetable crops grown in tropical and subtropical type of climatic conditions. Brinjal belongs to the family Solanaceae, also known as nightshades, are cousins of potatoes, tomatoes, etc. It

contributes 9% of the total vegetable production of the country (Sidhu et al., 2006). Eggplant is rich in dietary fiber, Vitamin B1 and copper. Other nutrients like manganese, vitamin B6, niacin, potassium, folate and Vitamin K are also found in good quantities. It also plays an important role in combating malnutrition in undernourished regions. According to the Food and Agriculture Organization of the United Nations (FAO), a minimum raise of 50% in agricultural food production is required in a scenario of modest economic growth. So, we have to focus on increasing the yield per area and decreasing the yield losses (FAO, 2017; Mauser et al., 2015; Pradhan et al., 2015). Pests and diseases cause severe losses to the overall agricultural production. Therefore, reducing the yield losses due to diseases is important for sustainable food production and nutritional security (FAO, 2009, Valin et al., 2014). Various chemicals are used for meeting this objective, they not only control the diseases but also increase crop yield, quality and shelf life (Cooper and Dobson, 2007; Kawasaki and Lichtenberg, 2015). Despite their usefulness in disease management, continued use of chemicals are a cause of concern because of their negative impact on human health and environment including toxicity in soil and water (Carvalho, 2006; Damalas et al., 2011; Geiger et al., 2010; Kim et al., 2017).

Brinjal is affected by several diseases which are a major limiting factor for decreasing total brinjal production. They mainly affect the roots, leaves, stems and fruits. Sclerotinia Blight is one of the most destructive disease of brinjal. Yield losses in susceptible crops vary and can be as high as 100% (Purdy, 1979).

Pseudomonads are soil borne, unicellular, rod shaped, obligate aerobic bacteria, with the long axis straight or curved but not helical, motile by means of one or more flagella, gram negative, do not form spore, stalks, or sheaths. Fluorescent Pseudomonads (FLPs) belong to the genus *Pseudomonas*, family Pseudomonadaceae, order Gamma proteobacteria and phylum Proteobacteria. There are 191 species in this genus and Pseudomonads are the most diverse and ecologically significant group of bacteria which are able to survive harsh conditions, are ubiquitous and form association with both plants and animals (Spiers et al. 2000).

Three species of *Pseudomonas* are Fluorescent, namely *Pseudomonas aeruginosa*, *P. fluorescence* and *P. putida* (Stanier et al. 1966). *P. aeruginosa* is the type species of genus *Pseudomonas* and is more uniform as it has less number of biotypes as compared to other Fluorescent Pseudomonads.

Hence, keeping in mind the exponential need to discover PGPR that are suitable for the crops and climate, following research has been taken up to identify the potential

indigenous FLPs of local areas. It aims to compare and find potential strains that can be used in disease reduction as an alternative to chemicals with the following objectives. *In vitro* and *in vivo* evaluation of bio-control potential of Fluorescent Pseudomonads against sclerotinia blight of brinjal as well as molecular characterization of native Fluorescent Pseudomonads isolates.

Materials and Methods

Antagonistic bacteria

Ten isolates of Fluorescent Pseudomonads were isolated from the rhizosphere of different plants at the farm of Bihar Agricultural University, Sabour. In our present study, we took two best performing isolates among them, named FLP 2020-1 and FLP 2020-2 both of which originated from brinjal rhizosphere.

Isolation and identification of the test pathogens

Parts of brinjal infected with *Sclerotinia sclerotiorum* (SCL) were removed and surface sterilized using 70% ethanol. They were further cut into small pieces (2.0 × 2.0 × 2.0 cm) using a sterilized scalpel and washed with sterilized distilled water. Then the infected pieces were soaked into 1% sodium hypochlorite for 3 min, then rinsed for 1 min in several changes of distilled water three times and dried between two layers of sterilized filter paper to remove excess water. Finally, the surface-sterilised pieces were plated onto potato dextrose agar (PDA) media (Prepared by the procedure of Tuite, 1969) in Petri-plate (70mm) under laminar airflow cabinet and incubated at 25±2 °C for 5 days.

After complete fungal growth it was confirmed by microscopic view. For confirmation through Koch Postulate followed by stem cutting method was performed (Kull et al., 2003) for inoculating plant.

Dual culture technique

Bioagent isolates were streaked in a single straight line 1 cm away from the edge of one side of the Petri-plate (70mm) containing potato dextrose agar (PDA) media. Fungus (4mm agar plug) were placed on opposite side of the Petri-plate, about 1cm away from the rim of the Petri-plate (70mm). A plate without any bioagent inoculation was carried out as negative control (Dennis and Webster, 1971).

Inverted plate technique

A plate (70mm) containing PDA was inoculated with a 4mm agar plug of active fungal test pathogen using a cork borer and a loop full of bioagent by a streaking loop separately in an individual plate. Half plate with the fungal test pathogen was inverted over the half plate containing the bioagent and in case of negative control half plate with the

fungal test pathogen was inverted over a half plate containing only PDA, without any bioagent. Seven replications were maintained for each case (Tapwal et al., 2015). The plate was allowed to incubate until complete growth was reported on control plate and all the experiments were maintained at 25 ± 2 °C.

Preparation of talc-based formulation of Fluorescent Pseudomonads and seed treatment

The talc-based formulation of Fluorescent Pseudomonads was prepared with some modifications to the method described by Vidhyasekaran and Muthamilan (1997). A loopful of Fluorescent Pseudomonads strain was inoculated into the nutrient broth Media and incubated in an incubator cum shaker at 150 rpm for 48 hours at the temperature of 28 ± 2 °C. 1 kg of talc powder was taken in a sterilized tray and its pH was adjusted to 7 by mixing CaCO_3 at the rate of 15 gm kg^{-1} . 10 gm of carboxymethyl cellulose (CMC) was mixed with 1 kg of talc and the mixture was autoclaved for 30 minutes for two consecutive days. 400 ml of 48 hour old bacterial suspension containing $9\times 10^8 \text{ cfu ml}^{-1}$ (adjusted by Spectrophotometer) was added to the carrier-cellulose mixture under aseptic conditions. After drying (approximately to 30% moisture content) overnight, the formulation was packed in a polypropylene bag, sealed and stored at room temperature (25 ± 2 °C). This *Pseudomonas* formulation powder @ 10g/kg of seeds was mixed in 1 lit of water and the brinjal seeds were soaked in this mixture for 30 minutes before sowing.

Artificial inoculation of test pathogens

For inoculation of *Sclerotinia sclerotiorum* the main stems of 5-week-old brinjal plants (fifth to sixth trifoliolate leaf fully expanded) were horizontally severed with a sterile razor blade at 0.5 cm above either the fourth or fifth node. A single mycelial plug (3 mm^2) was placed mycelial-side down on the cut stem (Kull et al., 2003). We took observations of no. of diseased plants and healthy plants.

For this experiment, atrial (with SCL) was conducted in a completely randomized design (CRD) with four replicates (four random blocks) with variety Muktakashi of brinjal. The studies are on six treatments per block as described below (Table 1). All statistical analyses were performed with OPSTAT software.

Table 1: Experimental design

T1	: Only seed
T2	: Bioagent 1 (FLP 2020-1)
T3	: Bioagent 2 (FLP 2020-2)
T4	: Bioagent 1 (FLP 2020-1) + Pathogen
T5	: Bioagent 2 (FLP 2020-2) + Pathogen

DNA extraction and PCR amplification of specific gene of FluorescentPseudomonads strains

Extraction of genomic DNA from all the selected bioagent strains was performed using a DNA isolation kit (Nucleo-pore gDNA Fungal Bacterial Mini Kit (50). After extraction, the quantity, integrity, and quality of the DNA obtained was checked by agarose gel electrophoresis, followed by staining in ethidium bromide, and visualization under UV light. To amplify the 16S rRNA gene, polymerase chain reaction (PCR – Eppendorf Master cycler) was performed from the genomic DNA of strains using a 16S rRNA gene specific primers - 27F (AGAGTTTGATCMTGGCTCAG) and 1492R (TACGGYTACCTTGTTACGACTT) as forward and reverse primer respectively, which was set at the amplification conditions (Initial denaturation at 94 °C for 3 minutes, denaturation at 94 °C for 30 seconds, annealing at 50 °C for 45 seconds, extension at 72 °C for 1:50 minutes and final extension at 72 °C for 10 minutes). PCR Products were checked by agarose gel electrophoresis, followed by staining in ethidium bromide, and visualization under UV light, following this, a ladder will be used as size markers. Then sequencing was done to check the diversity (Vinay et al., 2016).

Identification of FluorescentPseudomonads and Sequence Analysis

The amplified 16s rRNA gene region, Pseudomonas specific gene region of the FluorescentPseudomonads isolates were amplified with same primer pair in same procedure as above mentioned and sequenced directly by primer walking technique. Then the sequences were compared with those extracted from GenBank using BLASTN programme in NCBI (National Centre for Biotechnology Information) website and aligned using CLUSTAL W programme using MEGA 11 software.

Results

Isolation of test pathogen

The infected brinjal plant samples of fusarium wilt, sclerotinia blight and bacterial wilt were collected in polythene bags from vegetable farm of Bihar Agricultural University. The collected samples were processed at PG Lab-3, Department of Plant Pathology. The samples with a fluffy or cottony, white moldy growth (mycelium) around of lower stems were used for *Sclerotinia sclerotiorum* isolation. The surface sterilisation of diseased samples was done with 1% sodium hypochlorite. Standard isolation method was followed for isolation

of fungal pathogens. The pure culture of each pathogen was observed and identified based on their colony characteristics and spore morphology.

***Sclerotinia sclerotiorum* (SCL)**

The colony colour of fungi on petri dish was white colour and layer of mycelium was in irregular pattern, sclerodes were also seen which was rounded and black. The width of hyphae was 2.5-9 μm , microconidia was of 2-3 μm size, width of 2-6.5 μm and length of 2.5-14 μm for sclerotia, width of 4-6 μm and length of 6-12 μm for ascospore which was hyaline was also observed.

In vitro* evaluation of FLPs against *Sclerotinia sclerotiorum

Sclerotinia sclerotiorum showed the full growth within 5 days of inoculation into the petri-dish. The recorded data indicates that two FLPs isolates showed their potentiality to control the growth of *Sclerotinia sclerotiorum* in *in vitro*. The maximum radial growth of fungus was observed in FLP 2020-1 (1.71 mm) and minimum radial growth of fungus was reported in FLP 2020-2 (0.96 mm) in case of dual culture method. Similarly, maximum mycelium inhibition compare to negative control of the fungus was reported under FLP 2020-2 (98.59%) which is at per with FLP 2020-1 (97.56%). In case of inverted plate technique the maximum radial growth of fungus was observed in FLP 2020-1 (3.07 mm) and minimum radial growth of fungus was reported in FLP 2020-2 (1.36 mm). Similarly, maximum mycelium inhibition compare to negative control of the fungus was reported under FLP 2020-2 (98.06%) followed by FLP 2020-1 (95.61%). (Table 2, Fig 1 and 2)

***In vivo* evaluation of FLPs against *Sclerotinia sclerotiorum* (SCL)**

Brinjal seeds (Var. Muktakeshi) were bacterized with FLPs isolates by talc-based formulation containing 1×10^9 CFU/gm in case of treatments with FLPs isolates (FLP 2020-1, FLP 2020-2, FLP 2020-1+SCL, FLP 2020-2+SCL). The seedlings of FLPs treated seeds were challenged with SCL through stem cutting method. One negative control without FLPs treatment was also carried out by same procedure. Disease Incidence (%), biocontrol efficiency and no. of diseased and healthy plants were recorded under these various treatments. No disease incidence was observed under the treatments without any pathogen inoculation. Least disease incidence was reported under FLP 2020-2 (20.53%) treatment (FLP 2020-2+SCL) followed by FLP 2020-1 (23.47%) treatment (FLP 2020-1+SCL) compare to negative control (90.60%) treatment (Treatment with only pathogen). Maximum disease control was reported by FLP 2020-2 (77.33%) followed by FLP 2020-1 (74.09%) compare to negative control treatment (0%). So, FLP 2020-2 has more biocontrol efficiency than FLP 2020-1 against *Sclerotinia sclerotiorum*. No diseased plant was observed under the

treatments which are without pathogen inoculation. Maximum healthy plant was observed with FLP 2020-2 treatment (around 23 plants), which was at par with FLP 2020-1 treatment (22 plants) compare to negative control (around 3 plants). Least no. of diseased plant was reported with FLP 2020-2 (6 plants) followed by FLP 2020-1 (around 7 plants) compare to negative control (around 27 plants). (Table 3; Fig 3)

Molecular characterization of FLP strains

Bacterial genomic DNA was isolated from fifteen different FLP strains and isolated genomic DNA was analysed by electrophoresis on 1 % agarose gel. PCR was carried out for the amplification of 16S rDNA region by using 27F (AGAGTTTGATCMTGGCTCAG) and 1492R (TACGGYTACCTTGTTACGACTT) as forward and reverse primers respectively. Amplified PCR products were also analysed by electrophoresis on 1 % agarose gel. The DNA bands were visualized under UV trans-illuminator. Using this pair of primers, 1500 bp amplified band was obtained which was specific for 16S rDNA gene in *Pseudomonas spp.* isolates. Further PCR products were sequenced and diversity among strains was reported.

Construction and Analysis of Phylogenetic structure deduced from nucleotide sequences of the 16s rRNA

The nucleotide sequences of the 16S rRNA, *Pseudomonas* specific genes from ten Fluorescent *Pseudomonads* isolates were determined and phylogenetic tree based on this data showing evolutionary relatedness of the isolates with known *Pseudomonas spp.*, which were obtained by BLAST of the nucleotide sequences of our investigating isolates in NCBI website, was constructed by the Neighbour-Joining (NJ) method with Bootstrap value 1000.

In the constructed phylogenetic tree (Fig 4), our ten investigating Fluorescent *Pseudomonads* isolates grouped into a single cluster and known *Pseudomonas spp.* (Fig 4) from NCBI (National Centre for Biotechnology Information) website into another cluster. These two clusters showed 82% identity with each other. Among the 10 FLP isolates assessed 9 were found to have maximum similarity with *Pseudomonas aeruginosa* including FLP 2020-2 and FLP 2020-1 was found to be *Pseudomonas fluorescence* which is in confirmation with the morphological observations (Table 4).

Discussion

In vitro* evaluation of Fluorescent Pseudomonads isolates against *Sclerotinia sclerotiorum

In our investigation *Sclerotinia sclerotiorum* showed complete growth within 5 days of inoculation into the Petri-plate. Growth inhibition of the fungus was reported by all the isolates - FLP 2020-1 and FLP 2020-2 in dual culture method as well as inverted plate technique in *in vitro* condition.

The radial growth of the pathogen varies from 1 to 2 mm and 1 to 3 mm respectively in case of dual culture method and inverted plate technique against the bio-agents. The strains of FLPs shows inhibition against the pathogen at the range of 98% to 99% and 96% to 98% respectively in case of dual culture method and inverted plate technique. The maximum radial growth of fungus was observed in FLP 2020-1 (1.71 mm) and minimum radial growth of fungus was reported in FLP 2020-2 (0.96 mm) in case of dual culture method. Similarly, maximum mycelium inhibition compare to negative control of the fungus was reported under FLP 2020-2 (98.59%) which is at par with FLP 2020-1 (97.56%) in case of dual culture method. The maximum radial growth of fungus was observed in FLP 2020-1 (3.07 mm) and minimum radial growth of fungus was reported in FLP 2020-2 (1.36 mm) in case of inverted plate technique. Similarly, maximum mycelium inhibition compare to negative control of the fungus was reported under FLP 2020-2 (98.06%) followed by FLP 2020-1 (95.61%) in case of inverted plate technique.

Three Pseudomonads strains was examined by Mokrani et al., (2018). These Pseudomonads strains showed antagonistic effect against *Fusarium oxysporum* f.sp *phaseoli* and *Sclerotinia sclerotiorum* in dual culture. They showed inhibition ranging from 47.78 % to 100 per cent. The possible reasons for mycelial inhibition in dual culture method are the antimicrobial substances like - Phenazine-1-carboxylic acid, acetamido anthranilic phenol, HCN etc. and in inverted plate technique are the volatile compounds like – dimethyl disulphide, naphthalene, 1-methyl, benzaldehyde etc. which are produced by *fluorescent pseudomonads*. So, from our present study we can confirm that FLP 2020-2 produces more antimicrobial substances and volatile compounds than FLP 2020-1 (Karimi et al., 2012; Abed et al., 2016; Islam et al., 2018; SKM Al-Tameemi et al., 2019)

In vivo* evaluation of Fluorescent Pseudomonads isolates against *Sclerotinia sclerotiorum

Seed bacterization with Fluorescent Pseudomonads enhances the plant growth as well as resistance against plant pathogenic fungi under controlled and field conditions (Kumar and Dubey, 1992). Bacterization of peanut seeds with GRC1 strain of *Pseudomonas spp.* resulted in reduced stem-rot of peanut in *Sclerotium sclerotiorum* infested soil by 87%.

Fernando et al., (2007) worked on biological control of *Sclerotinia sclerotiorum* by *Pseudomonas* and *Bacillus spp.* on canola petals and reported disease incidence of 18.8 to

27.9% in *Pseudomonas spp.* treated plot. Singh et al., (2015) observed *in vivo* efficacy of *Pseudomonas* against *Sclerotinia sclerotiorum* of rajmash by pot experiment and they found 36% disease intensity in *Pseudomonas* treated pot which is less than untreated pot. Biocontrol of *Sclerotinia sclerotiorum* in oilseeds were also reported by Kamal et al., (2016).

In present study brinjal seeds (Var. Muktakeshi) were bacterized with FLPs isolates by FLPs formulation powder in case of treatments with FLPs isolates (FLP 2020-1, FLP 2020-2, FLP 2020-1+SCL, FLP 2020-2+SCL) and stem cutting method was carried out to inoculate the pathogen in the respective treatments with the test pathogen (Pathogen, FLP 2020-1+SCL, FLP 2020-2+SCL). No disease incidence was observed under the treatments without any pathogen inoculation.

The strains of FLPs shows disease incidence at the range of 20% to 24% and disease control percentage from 74% to 78%. Least disease incidence was reported under FLP 2020-2 (20.53%) treatment (FLP 2020-2+SCL) followed by FLP 2020-1 (23.47%) treatment (FLP 2020-1+SCL) compare to negative control (90.60%) treatment (Treatment with only pathogen). Maximum disease control was reported by FLP 2020-2 (77.33%) followed by FLP 2020-1 (74.09%) compare to negative control treatment (0%). So, FLP 2020-2 has more biocontrol efficiency than FLP 2020-1 against *Sclerotinia sclerotiorum*. No diseased plant was observed under the treatments which are without pathogen inoculation. Maximum healthy plant was observed with FLP 2020-2 treatment (around 23 plants), which was at par with FLP 2020-1 treatment (22 plants) compare to negative control (around 3 plants). Least no. of diseased plant was reported with FLP 2020-2 (6 plants) followed by FLP 2020-1 (around 7 plants) compare to negative control (around 27 plants).

Fluorescent *Pseudomonads* produce antimicrobial substances (Phenazine-1-carboxylic acid, acetamido anthranilic phenol etc.) which can inhibit the growth of mycelium and germination of ascospores of the pathogen (Fernando et al., 2007). Bacterial antagonists induce resistance by enhancement of lignification and stimulation of host defence enzymes and synthesis of pathogenesis-related (PR) proteins. The PR proteins like- chitinase and β -1,3-glucanase inhibits fungal pathogens. The enhanced accumulation of PR proteins and oxidative enzymes including chitinase and β -1,3-glucanase may also be responsible for the reduction of *Sclerotinia* infection. As, fungi have chitin and glucan as cell wall components, increased activity of chitinase and β -1,3- glucanase may prevent establishment of pathogen. These activities of the bio-agent can be the reason behind the least disease incidence percentage and more disease control percentage in the bio-agent treated plot. (Ramamoorthy et al., 2002)

Molecular characterization of Fluorescent Pseudomonads isolates

In the present study the nucleotide sequences of the 16s rRNA, *Pseudomonas* specific genes from ten Fluorescent Pseudomonads isolates were determined and a phylogenetic tree based on this data showing evolutionary relatedness of the isolates with known *Pseudomonas* spp., which were obtained from NCBI database, were constructed by the Neighbour-Joining (NJ) method with Bootstrap value 1000. The Phylogenetic analysis of the Fluorescent Pseudomonad isolates grouped into a single cluster and known *Pseudomonas* spp. (Fig 4) from NCBI (National Centre for Biotechnology Information) website were into another cluster. These two clusters showed 82% identity with each other. Among the 10 FLP isolates assessed, 9 were found to have maximum similarity with *Pseudomonas aeruginosa* including FLP 2020-2 and FLP 2020-1 was found to be *Pseudomonas fluorescence*. So, we can say that our investigating isolates are related to *Pseudomonas aeruginosa* strain and *Pseudomonas* spp. strain. All investigating isolates formed a cluster which may be due to geographical characteristics. Similar observation was recorded by Krishnamoorthi et al., (2020) working on molecular characterization of *Pseudomonas* spp. by 16S rRNA primer pair 27F & 1492 R and observed 1500 bp amplified band after PCR and they identified *Pseudomonas aeruginosa* after the analysis of sequence of the strains.

Conclusion

The present experiment was carried out to characterize the two best performing native Fluorescent Pseudomonads isolated and collected from the rhizosphere of Brinjal from agroclimatic zone of Bihar. The isolates were examined against *Sclerotium sclerotiorum* of brinjal under *in vivo* and *in vitro* conditions by dual culture method and inverted plate technique. In both conditions, FLP 2020-2 gave the better result by inhibiting the fungal pathogen. From our study we can suggest the use of the respective strain of Fluorescent Pseudomonads as not only a bio-control measure of diseases but also as plant growth promoting rhizobacteria for achieving sustainable agriculture and our goal should be to make them into reliable, assessable product for farmers at commercial level.

References

- Abed H, Rouag N, Mouatassem D, Rouabhi A(2016). Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of *Fusarium* wilt of chickpea. *Eurasian Journal of Soil Science*, 5(3): 182-191.

- Al-Tameemi, S K, Matloob, A A A H (2019). Effect of some biological control agents in reducing the disease incidence and severity of white mold disease on eggplant caused by *sclerotinia sclerotiorum*. *Plant Archives*, 19(2): 646-652.
- Carvalho F P (2006). Agriculture, pesticides, food security and food safety. *Environmental science and policy*, 9(7-8): 685-692.
- Damalas C A, Eleftherohorinos I G (2011). Pesticide exposure, safety issues, and risk assessment indicators. *International journal of environmental research and public health*, 8(5): 1402-1419.
- Dennis C, Webster J (1971). Antagonistic properties of species-groups of *Trichoderma*: III. Hyphal interaction. *Transactions of the British Mycological Society*, 57(3): 363-IN2.
- Fernando W G D, Nakkeeran S, Zhang Y, Savchuk S (2007). Biological control of *Sclerotinia sclerotiorum* (Lib.) de Bary by *Pseudomonas* and *Bacillus species* on canola petals. *Crop protection*, 26(2): 100-107.
- Geiger F, Bengtsson J, Berendse F, Weisser W W, Emmerson M, Morales M B, Ceryngier P, Liira J, Tschardt T, Winqvist C, Eggers S, Bommarco R, Pärt T, Bretagnolle V, Plantegenest M, Clement L W, Dennis C, Palmer C, Oñate J J, Guerrero I, Hawro V, Aavik T, Thies C, Flohre A, Hänke S, Fischer C, Goedhart P W, Inchausti P, Inchausti P (2010). Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology*, 11(2): 97-105.
- Godfray H C J, Beddington J R, Crute I R, Haddad L, Lawrence D, Muir J F, Pretty J, Robinson S, Thomas S M, Toulmin C (2010). Food security: the challenge of feeding 9 billion people. *science*, 327(5967): 812-818.
- Islam MA, Nain Z, Alam MK, Banu NA, Islam MR (2018). In vitro study of biocontrol potential of *rhizospheric Pseudomonads aeruginosa* against *Fusarium oxysporum* sp. *cucumerinum*. *Egyptian journal of Biological Pest Control*, 28(1): 90.
- Kamal M, Savocchia S, Lindbeck K D, Ash G J (2016). Biology and biocontrol of *Sclerotinia sclerotiorum* (Lib.) de Bary in oilseed Brassicas. *Australasian Plant Pathology*, 45(1): 1-14.

- Karimi K, Amini J, Harighi B, Bahramnejad B (2012). Evaluation of biocontrol potential of '*Pseudomonas*' and '*Bacillus*' spp. against fusarium wilt of chickpea. *Australian Journal of Crop Science*, 6(4): 695-703.
- Kawasaki K, Lichtenberg E (2015). Quality versus quantity effects of pesticides: joint estimation of quality grade and crop yield (No. 330-2016-13915).
- Kim K H, Kabir E, Jahan S A (2017). Exposure to pesticides and the associated human health effects. *Science of the total environment*, 575: 525-535.
- Krishnamoorthi M, Muthamilan M, Sendhilvel V, Angappan K, Anandham R (2020). Isolation and molecular characterization of *Pseudomonas* spp. with antagonistic activity against *Macrophomina phaseolina*. *Journal of Pharmacognosy and Phytochemistry*, 9(3): 2155-2158.
- Kull L S, Vuong T D, Powers K S, Eskridge K M, Steadman J R, Hartman G L (2003). Evaluation of resistance screening methods for *Sclerotinia* stem rot of soybean and dry bean. *Plant disease*, 87(12): 1471-1476.
- Kumar B D, Dube H C (1992). Seed bacterization with a fluorescent *Pseudomonas* for enhanced plant growth, yield and disease control. *Soil Biology and Biochemistry*, 24(6): 539-542.
- Mausser W, Klepper G, Zabel F, Delzeit R, Hank T, Putzenlechner B, Calzadilla A (2015). Global biomass production potentials exceed expected future demand without the need for cropland expansion. *Nature communications*, 6(1): 1-11.
- Mokrani S, Belabid L, Nabti E (2018). Antifungal Activities of *Pseudomonas* Strains Against Indigenous Phytopathogenic fungi of *Phaseolus vulgaris* L. Isolated from a Greenhouse in western Algeria *Res. J. Plant Pathol.*, 2(1): 07.
- Pradhan P, Fischer G, van Velthuis H, Reusser D E, Kropp J P (2015). Closing yield gaps: how sustainable can we be?. *PloS one*, 10(6): e0129487.
- Purdy L (1979). *Sclerotinia sclerotiorum*: history, diseases and symptomatology, host range, geographic distribution, and impact. *Phytopathology*, 69(8): 875-880.
- Ramamoorthy V, Raguchander T, Samiyappan R (2002). Induction of defense-related proteins in tomato roots treated with *Pseudomonas fluorescens* Pf1 and *Fusarium oxysporum* f. sp. *lycopersici*. *Plant and soil*, 239(1): 55-68.

- Sidhu A S, Dhatt, A S (2006, December). Current status of brinjal research in India. In I International Conference on Indigenous Vegetables and Legumes. Prospectus for Fighting Poverty, Hunger and Malnutrition 752: (pp. 243-248).
- Singh R, Singh D K, Singh G. Biological control of stem rot of rajmash (*Phaseolus vulgaris* L.).
- Spiers, Andrew and Buckling, Angus and Rainey, Paul (2000). The cause of *Pseudomonas* diversity. *Microbiology* (Reading, England), 146(10):2345-50.
- Stanier RY, Palleroni NJ, Doudoroff M (1966). The aerobic *pseudomonads*: a taxonomic study. *J Gen Microbiol*, 43(2):159-271.
- Tamura K, Stecher G, Kumar S (2021). MEGA 11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* <https://doi.org/10.1093/molbev/msab120>.
- Tapwal A, Thakur G, Chandra S, Tyagi A (2015). In-vitro evaluation of *Trichoderma* species against seed borne pathogens. *Int J Biol Chem Sci*, 1(10): 14-19.
- The future of food and agriculture Trends and challenges, Food and Agriculture Organization of the United Nations Rome, 2017: <https://www.fao.org/3/i6583e/i6583e.pdf>.
- The state of food and agriculture, Food and Agriculture Organization of the United Nations Rome, 2009: <https://www.fao.org/3/i0680e/i0680e.pdf>.
- Tuite J (1969). *Plant Pathological method. Fungi and Bacteria*.
- Valin H, Sands R D, Van der Mensbrugge D, Nelson G C, Ahammad H, Blanc E, Bodirsky B, Fujimori S, Hasegawa T, Havlik P, Heyhoe E, Kyle P, Mason-D' Croz D, Paltsev S, Rolinski S, Tabeau A, Meijl H V, Lampe M V, Willenbockel D (2014). The future of food demand: understanding differences in global economic models. *Agricultural Economics*, 45(1): 51-67.
- Vidhyasekaran P, Rabindran R, Muthamilan M, Nayar K, Rajappan K, Subramanian N, Vasumathi K, (1997). Development of powder formulation of *Pseudomonas yuorescens* for control of rice blast. *Plant Pathology*, 46: 291-297.

Vinay J U, Naik M K, Rangeshwaran R, Chennappa G, Shaikh S S, Sayyed R Z (2016).

Detection of antimicrobial traits in fluorescent pseudomonads and molecular characterization of an antibiotic pyoluteorin. 3 Biotech, 6(2): 1-11.

World Population Prospects 2019, United Nation:

<https://www.un.org/development/desa/publications/world-population-prospects-2019-highlights.html>.

UNDER PEER REVIEW

Table 2: *In vitro* evaluation of FluorescentPseudomonads isolates against *Sclerotinia sclerotiorum* of brinjal

<i>Sclerotinia sclerotiorum</i>				
	Dual culture method		Inverted plate technique	
Treatments	Radial growth (mm)	Inhibition over control (%)	Radial growth (mm)	Inhibition over control (%)
FLP 2020-1	1.71 ^b	97.56 ^a	3.07 ^b	95.61 ^b
FLP 2020-2	0.96 ^a	98.59 ^a	1.36 ^a	98.06 ^a
Control	70 ^c	0 ^b	70 ^c	0 ^c
SeM (±)	0.24	0.35	0.28	0.39
CD (0.01)	0.71	1.04	0.82	1.17
CV (%)	2.61	1.42	2.94	1.61

Table 3: *In vivo* evaluation of FluorescentPseudomonads isolates against *Sclerotinia sclerotiorum* of brinjal

Treatments	<i>Sclerotinia sclerotiorum</i>			
	Disease incidence (%)	Disease control (%)	No. of healthy plant	No. of diseased plant
Control	0 ^a	-	29 ^a	0 ^a
FLP 2020-1	0 ^a	100 ^a	29 ^a	0 ^a
FLP 2020-2	0 ^a	100 ^a	29 ^a	0 ^a
FLP 2020-1+SCL	23.47 ^c	74.09 ^c	22 ^b	7 ^c
FLP 2020-2+SCL	20.53 ^b	77.33 ^b	23 ^b	6 ^b
SCL	90.60 ^d	0 ^d	3 ^c	27 ^d
SeM (±)	0.75	0.32	0.44	0.28
CD (0.01)	2.23	0.96	1.32	0.84
CV (%)	6.69	0.86	29.25	8.64

Table 4: Fluorescent Pseudomonads isolates, used in phylogenetic analysis

Sl. No.	Fluorescent Pseudomonads isolates	Origin	Code	Similarity	Sequenced product length
1.	<i>Pseudomonas spp.</i>	Brinjal Rhizosphere	PF	<i>Pseudomonas aeruginosa</i> (90-95%)	1215 bp
2.	<i>Pseudomonas spp.</i>	Brinjal Rhizosphere	Br2 (FLP 2020-1)	<i>Pseudomonas fluorescence</i> (98.5%)	1239 bp
3.	<i>Pseudomonas spp.</i>	Brinjal Rhizosphere	Br4 (FLP 2020-2)	<i>Pseudomonas aeruginosa</i> (98.6%)	946 bp
4.	<i>Pseudomonas spp.</i>	Brinjal Rhizosphere	Pt	<i>Pseudomonas aeruginosa</i> (99%)	1274 bp
5.	<i>Pseudomonas spp.</i>	Rice Rhizosphere	Ri	<i>Pseudomonas aeruginosa</i> (97.5%)	1101 bp
6.	<i>Pseudomonas spp.</i>	Wheat Rhizosphere	Wh	<i>Pseudomonas aeruginosa</i> (98.6%)	1224 bp
7.	<i>Pseudomonas spp.</i>	Barley Rhizosphere	B	<i>Pseudomonas aeruginosa</i> (98.3%)	1276 bp
8.	<i>Pseudomonas spp.</i>	Mustard Rhizosphere	Mu2	<i>Pseudomonas aeruginosa</i> (98%)	1285 bp
9.	<i>Pseudomonas spp.</i>	Mango Rhizosphere	Ma	<i>Pseudomonas aeruginosa</i> (98.7%)	1154 bp
10.	<i>Pseudomonas spp.</i>	Turmeric Rhizosphere	Tu	<i>Pseudomonas aeruginosa</i> (98.6%)	1239 bp



Fig 1: *In vitro* antagonistic potential of Fluorescent Pseudomonads isolates against *Sclerotinia sclerotiorum* of brinjal under dual culture technique



Fig 2: *In vitro* antagonistic potential of Fluorescent Pseudomonads isolates against *Sclerotinia sclerotiorum* of brinjal under Inverted plate technique





Fig 3: Healthy and SCL infected plant under challenge inoculation conditions




Fig 4: Phylogenetic tree based on 16S rRNA gene sequences