

Formatted: English (United States)

Crop regulation studies in guava (*Psidium guajava* L.) cv. L 49 under Ultra High Density Planting

Formatted: English (United States)

ABSTRACT: A field experiment was carried out on 8 years old guava cv L 49 trees to assess the effect of pruning level and growth regulators with different combinations on vegetative growth and fruiting attributes in Ultra High Density Planting. The experiment consisted of 2 factors with 36 combinations viz., factor -1 (pruning level) P₁- pruning of 25 cm of the shoot from the tip, P₂- pruning of 50 cm of the shoot from the tip, P₃- pruning of 75 cm of the shoot from the tip, P₄ – control (without pruning). Factor -2(growth regulators) G₁, G₂- Potassium nitrate(1 ,2%) , G₃, G₄- Thiourea (0.1 ,0.2%) , G₅, G₆- Ethephon (250, 500 ppm) , G₇, G₈- Cycocel (250, 500 ppm) and G₉ – Control. Results showed that among different levels of pruning, 50cm pruning from the tip with potassium nitrate spray at 2 different concentrations was found to be the best for promoting early vegetative bud burst, fruit set and fruit yield. The increment in plant height and canopy spread in East-West and North-South were recorded more in 25 cm pruning from the tip. A greater number of shoots are visible in the P₃G₅ combination. When compared to other treatments, the P₃G₂ combination significantly increased the average index number of bud bursts and shoot length under Ultra High Density Planting.

Keywords: guava, factors, pruning, growth regulators, concentrations

Formatted: English (United States)

Formatted: English (United States)

INTRODUCTION

Guava (*Psidium guajava* L.) belongs to the family Myrtaceae, which has more than 80 genera and 3,000 species, distributed in the tropics and subtropics, native to tropical America stretching from Mexico to Peru. Guava is rich in ascorbic acid, calcium, iron, and phosphorus, which has led to it being referred to as the "poor man's apple" or the "apple of the tropics"(Prakash *et al.*, 2002). It contains 2-5 times more vitamin C as compared to oranges. According to an estimate, 100 g of guava fruit contains approximately 260 mg of vitamin C. Guava is the fifth most important fruit pertaining to area, production, and productivity among different fruit crops grown in India. It contributes 3.4 percent of the area and 3.9 percent of production in total fruit crops grown all over India. The other major states for guava cultivation are Uttar Pradesh, Bihar, and Karnataka. Maharashtra's most common variety is guava L-49 (Sardar). The production of 4.92 million MT of guava fruit has recently been regulated to a ~~315 thousand hectare~~ 315-thousand-hectare area in India.

Getting an increase in guava production per unit area can be accomplished by increasing the plant population (Singh *et al.*,1980, Mitra *et al.*, 1984). In the initial years, ~~ultra-high density~~ultra-high-density planting increases yield while simultaneously increasing net economic returns per unit area and allowing for more effective input utilization (Reddy, 2004).

MATERIAL AND METHODS

A field experiment was conducted at Horticultural College and Research Institute, Periyakulam, Theni in March 2023. Eight years old guava trees under Ultra High ~~Density Planting~~Density Planting with a spacing of 3 x 1.5 m were selected for a research trial. The treatment

Guava fruit is produced on the stalk of the current season and is produced by guava flowers, which appear singly or in cymes of two or three in the axil of leaves. Guava pruning is one of the most important practices that influences the vigor, productivity, and quality of the fruits. To build a strong architecture that can support a heavy crop load, early pruning is done (Lian *et al.*, 2019).

To maximize the unit area production and keep in mind its good demand, the production of genuine pruning techniques is a pre-requisite. It is always important for the fruit growers to adopt the best pruning technique. Therefore, the experiment was conducted ~~with a goal to~~ standardize the pruning technique for distribution to fruit growers. For the above purpose, standardize the pruning techniques needed to obtain good quality fruits. To adopt the correct pruning technique under UHDP, the level of pruning is important. Plant growth regulators like Potassium nitrate, Thiourea, Ethephon, and Cycocel play important roles in guava production by direct or indirect influences on the growth and development of guava. Pruning by different levels and application of growth regulators improve the yield and quality of the guava. In the UHDP system, the study's objective is to standardize the pruning level with the optimal growth regulators.

includes nine different concentrations of growth regulators (Potassium nitrate @ 1% & 2%, Thiourea @ 0.1% & 0.2%, Ethephon @ 250 ppm & 500 ppm, Cycocel @ 250 ppm & 500 ppm, and Control) as well as four various levels of pruning (25, 50, 75 cm of the shoot from the tip, and Control). The 36

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

was recorded in the P₂G₁ combination (21.61 m³) (Table 4). Canopy volume was at its highest in 25% pruning of previous season growth as compared to 75% pruning of previous season growth (Singh and Canaan 2005). Since pruning eliminates carbon-starved, fruit-exhausted branches and encourages the growth of new leaves to store carbs for the following flowering, it also permits lateral buds to sprout, which in turn affects the canopy volume and other vegetative characteristics of the plants. This is consistent with research on guava (Pilanin *et al.*, 2010).

1.4. Time taken for first vegetative bud burst (days)

The data on variations in pruning level and growth regulators showed that the P₂G₂ combination had the earliest emergence (5.01 days) and maximal (22.00 days) of vegetative bud burst in the P₄G₈ combination (Table 5). Early pruning caused new branches to sprout early. Similar findings were reported that a rise in pruning severity promotes the early bud burst (Lian *et al.*, 2019, Basu *et al.*, 2007, Dhaliwal *et al.*, 2000). Severe pruning along with potassium nitrate at 4% induced the highest average index number of bud bursts (Bhagawati *et al.*, 2015).

1.5. Number of shoots emerged from the pruned branches

The P₃G₅ combination (7.50) produced more on average number of new shoots followed by the P₃G₁ combination (7.30), according to the data in table (6). The lowest number of new shoots was recorded in the P₄G₃ combination (2.00). 75 cm pruning with ethephon at 250 ppm produce the highest number of new shoots compared to others. Pruning enhanced the guava's vegetative development similar these results (Bhagawati *et al.*, 2015, Pilanin *et al.*, 2010).

1.6. Length of newly emerged shoots (cm)

In terms of substance effect, the P₂G₄ combination had the longest average shoot length (43.50 cm), followed by the P₃G₃ and P₂G₁ combinations (Table 7). ~~With the passage of time~~ With time, moderate pruning has given a favorable effect. Interaction between the concentration of growth regulators and pruning level showed that the P₂G₄ combination gave the greatest value of shoot length. On ~~another side~~ the other hand, the shortest shoot length was recorded in the P₃G₁ combination (23.80 cm). Shoot length in guava plants increased as a result of pruning, which encouraged vegetative development (Shaban and Haseeb 2009). When the shoots were pruned three times a year to half of their original length, the maximum gross return was achieved as well (Bisla *et al.*, 1988).

2. Effect of pruning level and growth regulators on fruiting characters

2.1. Bud burst activity

Flower bud burst activity data in Table (8) showed that the highest significant average index number was recorded in the P₃G₂ combination (4.90) followed by the P₁G₂ combination (3.60). The lowest average index number was recorded in the P₄G₈ combination (1.40). Meanwhile, 75 cm pruning with KNO₃ at 2% shows the highest average index number of the flower bud burst.

2.2. Fruit set (%)

Over the control, all interaction effects showed an increase in the fruit set. The average Fruit set percentage was more in the P₂G₁ combination (96.00 %) followed by the P₂G₂ combination (89.00%). Absolute control (P₄G₀ combination) was recorded lowest fruit set (70.00%) (Table 9). A direct relationship was observed between the concentration of chemical substances and the fruit set was recorded with the highest concentrations. Moderate pruning and the application of 4% potassium nitrate were shown to significantly boost the early fruit set (Bhagawati *et al.*, 2015). Etephon and ethephon, when used at higher concentrations, decreased fruit set (Mehta *et al.*, 2012) [19]. Similar outcomes were also obtained in an increase in fruit set and fruit retention in guava by pruning over control (Curry and Williams 1989, Pilanin *et al.*, 2010).

2.3. Number of fruits/ Plant

The highest number of fruit per plant was recorded in the P₂G₁ combination (192.00) and the least number of fruit per plant was recorded in the P₄G₉ combination (55.80) (Table 10). The interaction between the two factors has a considerable impact on the total amount of fruit. You may control your tree's size and fruit production with the help of shoot pruning. Pruning increases the amount of fruit that a plant produces. Others made similar observations as well. Fruit yield is seen to decline as pruning severity is increased (Dalal *et al.*, 2000, Lopez *et al.*, 1982, Quijada *et al.*, 1999). Pruning, which decreased the fruiting area and on the other hand encouraged vegetative growth at the expense of reproductive growth, is the cause of the drop in fruit production per plant (Kumar and Rattanpal 2010).

2.4. Fruit yield (kg/ Plant)

Maximum yield was achieved with a P₂G₂ combination (27.10 kg) followed by a P₁G₈ combination (25.98 kg). The minimum yield was recorded in the P₄G₉ combination (8.37 kg) (Table 11). According to Dutta, foliar spraying of potassium increased the yield and quality and decreased with the pruning intensity in Sardar guava. Pruning increases the fruit weight in guava (Maji *et al.*, 2015). Using various crop regulation practices like the pruning of shoots, defoliation, or deblossoming, it is necessary to decrease fruit set during the rainy season and then

Formatted: English (United States)

Formatted: English (United States)

increase fruit set during the winter season in order to regulate guava crop, provides support to the present finding (Boora et al., 2016). The findings of

this study are consistent with 45 cm shoot pruning in May would be ideal for good guava off-season output (Meena et al., 2016).

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	0.83	0.63	1.19	0.13	0.70
G ₂	0.89	0.91	1.21	0.34	0.84
G ₃	0.74	1.27	1.05	0.41	0.87
G ₄	1.16	1.19	1.08	0.78	1.05
G ₅	0.38	0.88	0.92	0.70	0.72
G ₆	0.87	0.93	0.87	0.42	0.77
G ₇	0.68	1.00	0.91	0.37	0.74
G ₈	0.72	0.77	0.99	0.56	0.76
G ₉	0.71	0.90	1.24	0.58	0.86
Mean	0.78	0.94	1.05	0.48	
	Pruning		Growth regulators		Interaction (P×G)
SE(d)	0.008		0.012		0.025
CD(0.05)	0.017		0.025		0.051

Table 1. Effect of pruning level and growth regulators on increment in tree height

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	0.55	0.35	0.10	0.75	0.44
G ₂	1.00	0.40	0.25	0.50	0.54
G ₃	0.60	0.05	0.55	0.70	0.48
G ₄	0.70	0.15	0.30	0.10	0.31
G ₅	0.19	0.12	0.30	0.65	0.29
G ₆	0.85	0.20	0.40	0.50	0.49

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

G ₇	0.25	0.45	0.30	0.35	0.34
G ₈	0.65	0.11	0.45	0.85	0.52
G ₉	0.05	0.60	0.70	0.90	0.56
Mean	0.53	0.26	0.37	0.59	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.025		0.038		0.077
CD(0.05)	0.052		0.078		0.157

Table 2. Effect of pruning level and growth regulators on increment canopy spread (E-W)

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	0.20	0.91	0.55	0.55	0.55
G ₂	1.50	0.30	0.15	0.10	0.51
G ₃	0.05	0.60	0.40	0.35	0.35
G ₄	0.15	0.25	0.30	0.95	0.41
G ₅	0.35	0.20	0.50	0.30	0.34
G ₆	0.45	0.15	0.60	0.90	0.53
G ₇	0.15	0.40	0.50	0.85	0.48
G ₈	1.10	0.40	0.60	0.60	0.68
G ₉	0.25	0.65	0.75	1.30	0.74
Mean	0.47	0.43	0.48	0.66	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.024		0.036		0.072
CD(0.05)	0.048		0.073		0.146

Table 3. Effect of pruning level and growth regulators on increment canopy spread (N-S)

Growth regulators	Pruning level	Mean
-------------------	---------------	------

	P ₁	P ₂	P ₃	P ₄	
G ₁	35.41	21.61	29.32	32.18	29.63
G ₂	35.29	29.88	29.23	30.43	31.21
G ₃	32.48	29.07	26.60	29.44	29.40
G ₄	44.22	31.54	28.77	36.34	35.22
G ₅	27.19	31.58	25.56	33.31	29.41
G ₆	31.58	27.16	26.60	24.54	27.47
G ₇	33.31	33.91	25.07	27.16	29.86
G ₈	33.20	29.39	26.10	32.89	30.40
G ₉	33.88	41.18	32.16	43.21	37.61
Mean	34.06	30.59	27.71	32.17	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.217		0.326		0.653
CD(0.05)	0.441		0.662		1.325

Table 4. Effect of pruning level and growth regulators on canopy volume

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	11.22	9.37	8.00	19.65	12.06
G ₂	13.50	5.01	11.48	19.22	12.30
G ₃	13.32	7.46	8.52	23.56	13.22
G ₄	12.10	7.50	9.33	18.00	11.73
G ₅	15.67	8.27	7.58	22.59	13.53
G ₆	15.15	7.00	6.34	19.66	12.04
G ₇	11.00	10.35	12.29	21.63	13.82
G ₈	10.08	9.86	11.67	22.00	13.40
G ₉	9.00	11.07	10.74	20.53	12.84
Mean	12.34	8.43	9.55	20.76	

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

	Pruning	Growth regulators	Interaction (P×G)
SE(d)	0.121	0.181	0.36
CD(0.05)	0.246	0.369	0.738

Table 5. Effect of pruning level and growth regulators Time taken for first vegetative bud burst

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	2.60	5.50	7.30	3.25	4.66
G ₂	3.10	6.50	6.30	4.50	5.10
G ₃	2.60	6.60	6.50	2.00	4.43
G ₄	3.50	6.30	7.00	3.00	4.95
G ₅	2.10	6.83	7.50	3.00	4.86
G ₆	2.50	5.83	7.16	4.25	4.94
G ₇	2.50	6.63	6.16	4.75	5.01
G ₈	2.80	7.00	6.30	3.50	4.90
G ₉	2.50	6.00	6.60	4.25	4.84
Mean	2.69	6.35	6.76	3.61	

	Pruning	Growth regulators	Interaction (P×G)
SE(d)	0.041	0.062	0.125
CD(0.05)	0.085	0.127	0.255

Table 6. Effect of pruning level and growth regulators Number of shoots emerged from the pruned branches

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	29.50	42.10	23.80	27.00	30.60
G ₂	33.30	41.80	36.20	25.00	34.08
G ₃	33.30	42.10	42.40	30.60	37.10
G ₄	29.30	43.50	40.80	32.10	36.43
G ₅	35.50	41.50	30.40	27.50	33.73

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

G ₆	29.60	42.10	32.80	29.50	33.50
G ₇	32.10	35.80	32.80	33.10	33.45
G ₈	34.30	38.00	35.60	39.10	36.75
G ₉	30.80	36.50	32.80	38.80	34.73
Mean	31.97	40.38	34.18	31.41	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.331		0.496		0.993
CD(0.05)	0.672		1.008		2.017

Table 7. Effect of pruning level and growth regulators Length of newly emerged shoots

Growth regulators	Pruning level				Mean
	P ₁	P ₂	P ₃	P ₄	
G ₁	3.03	3.00	3.30	2.90	3.06
G ₂	3.60	3.20	4.90	2.00	3.43
G ₃	2.33	2.42	2.77	2.16	2.42
G ₄	1.93	2.15	2.31	1.49	1.97
G ₅	2.01	2.20	2.43	2.50	2.29
G ₆	1.60	2.28	2.40	1.77	2.01
G ₇	1.73	1.76	2.41	1.57	1.87
G ₈	2.03	1.80	1.82	1.40	1.76
G ₉	1.93	1.95	2.10	1.60	1.90
Mean	2.24	2.31	2.72	1.93	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.019		0.028		0.057
CD(0.05)	0.038		0.058		0.116

Table 8. Effect of pruning level and growth regulators bud burst activity

Growth regulators	Pruning level	Mean
-------------------	---------------	------

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font color: Auto, English (United States)

	P₁	P₂	P₃	P₄	
G₁	88.00	96.00	78.26	86.00	87.07
G₂	73.33	89.00	75.75	83.34	80.36
G₃	83.30	83.63	73.60	75.00	78.88
G₄	84.84	86.63	82.22	83.73	84.36
G₅	82.00	87.50	72.41	73.80	78.93
G₆	81.81	86.50	78.94	86.84	83.52
G₇	85.71	86.50	74.19	73.07	79.87
G₈	83.87	84.61	80.00	73.30	80.45
G₉	76.66	88.70	72.63	70.00	77.00
Mean	82.17	87.67	76.44	78.34	
	Pruning		Growth regulators		Interaction (PxG)
SE(d)	0.658		0.988		1.976
CD(0.05)	1.337		2.005		4.011

Table 9. Effect of pruning level and growth regulators on fruit set

Growth regulators	Pruning level				Mean
	P₁	P₂	P₃	P₄	
G₁	162.10	180.67	75.30	93.50	127.89
G₂	168.40	192.00	82.40	100.30	135.78
G₃	165.10	172.78	69.80	82.30	122.50
G₄	153.20	163.11	70.20	85.50	118.00
G₅	156.50	154.65	63.70	92.20	116.76
G₆	161.70	150.23	65.50	95.30	118.18
G₇	171.40	164.50	59.50	97.80	123.30
G₈	173.20	170.30	63.50	86.30	123.33
G₉	167.20	152.40	64.50	55.80	109.98
Mean	164.31	166.74	68.27	87.67	

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

ultimately impacts fruit quality and yield. This study's focus on crop regulation in guava under UHDP provides insights into managing plant growth, flowering, and fruiting patterns (Olivares et al. 2022). These findings can be extended to other fruit crops, contributing to the development of improved cultivation practices (Hernandez et al. 2020).

As global populations continue to rise, the demand for nutritious and sustainable food sources increases. Guava is known for its nutritional value, and optimizing its cultivation under UHDP can enhance yield while conserving land and resources. This study's findings could aid in meeting food security goals (Hernandez and Olivares, 2020; Pitti et al. 2021; Montenegro et al. 2021).

The study addresses the challenges of maximizing agricultural productivity in limited land areas (Hernandez et al. 2018a; 2018b), a common concern in densely populated countries like India. By investigating guava cultivation in UHDP, the research offers potential solutions to the problem of land scarcity and showcases a strategy to enhance productivity without expanding agricultural land (Hernandez and Olivares, 2019).

Climate change poses significant threats to crop production (Parra et al. 2012; Vilorio et al. 2023).

REFERENCES

Arava-Alman, M., Olivares, B., Acevedo-Opazo, C. et al. (2020). Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. *J Soil Sci Plant Nutr*, 20 (3): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>

Basu, J., Das, B., Sarkar, S., Mandal, K. K., Banik, B. C., Kundu, S., & Ray, S. K. (2007). Studies on the response of pruning for rejuvenation of old guava orchard. *Acta horticulturae*.

Bhagawati, R., Bhagawati, K., Choudhary, V. K., Rajkova, D. J., & Sharma, R. (2015). Effect of pruning Intensities on the performance of fruit plants under mid-hill condition of Eastern Himalayas: case study on Guava. *International Letters of Natural Sciences*, 46.

Bisla, S. S., Dhiman, B. K., & Daulta, B. S. (1988). Studies on pruning and spacing in ber (*Ziziphus mausitana* Lamk). Effect on vegetative growth. *Haryana J. Hort. Sci*, 17(3-4), 177-182.

Boora, R. S., Dhaliwal, H. S., & Arora, N. K. (2016). Crop regulation in guava—A review. *Agric Rev* 37 (1): 1-9.

Curry, E. A., & Williams, M. A. (1989). Fruit thinning with ethephon. *Good Fruit Grower, Wenatchee, Washington*, 90(8), 8-10.

Campos, B. O. (2023). Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health.

The study's findings on guava cultivation under UHDP could provide insights into how this planting technique influences the crop's resilience to changing climate conditions. This aspect of the study contributes to the broader understanding of crop adaptation to environmental challenges (Zingaretti et al. 2016; Olivares et al. 2020).

Conclusion

In conclusion the typical flowering and fruiting behavior of the plant required to be controlled in order to produce an enormous crop load and make guava cultivation very profitable under Ultra High density planting. In terms of growth and flowering yield, all crop regulating techniques were determined to be superior to the untreated control. Unpruned guava trees have a tendency to prolong vegetative growth and diminish the bearing area, which reduces fruit size, yield, and quality. Pruning is therefore necessary to achieve a suitable balance between vegetative and reproductive growth. According to this study's findings, 50 cm of pruning from the tip using potassium nitrate at 1% and 2% is superior for enhancing vegetative growth and production of guava trees.

Formatted: Font: Not Bold, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Springer Nature. <https://doi.org/10.1007/978-3-031-34475-6>

Dalal, S. R., Golliwari, V. J., Patil, S. R., Khobragade, R. I., & Dalal, N. R. (2000). Effect of severity of pruning on growth, yield and quality of fruits of 25 year old guava cv Sardar. *Journal of Soils and Crops*, 10(2), 298-300.

Dhaliwal, G. S., Nanra, N. K., & Rattanpal, H. S. (2002). Effect of chemicals on flower drop, fruit set and yield on rainy and winter season crops of guava. *Indian Journal of Horticulture*, 59(1), 31-33.

Dhaliwal, G. S., Rattanpal, H. S., & Gill, H. S. (2000). Effect of time and severity of pruning on cropping and physico-chemical properties of Sardar guava. *Haryana Journal of Horticultural Sciences*, 29(1/2), 17-20.

Hariom, S., & Shant, L. (2015). Effect of shoot pruning on growth, flowering and yield in meadow orchard of guava cv Pant Prabhat. *International Journal of Basic and Applied Agricultural Research*, 13(3), 395-399.

Hernandez, R., Olivares, B., Arias, A., Molina, J.C., Pereira, Y. (2020). Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*, 38(2):95-102. <http://dx.doi.org/10.4067/S071834292020000200095>

Hernández, R., Olivares, B. (2020). Application of multivariate techniques in the agricultural land's

aptitude in Carabobo, Venezuela. <i>Tropical and Subtropical Agroecosystems</i>, 23(2):1-12. https://n9.cl/zeedh	Olivares, B. (2022). Machine learning and the new sustainable agriculture: Applications in banana production systems of Venezuela. <i>Agric. Res. Updates</i>, 42, 133-157.	Formatted
Hernández, R; Olivares, B. Arias, A; Molina, J.C., Pereira, Y. (2018a). Agroclimatic zoning of corn crop for sustainable agricultural production in Carabobo, Venezuela. <i>Revista Universitaria de Geografía</i>, 27 (2): 139-159. https://n9.cl/12m83	Olivares, B., Paredes, F., Rev, J., Lobo, D., Galvis-Causil, S. (2021). The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. <i>SAINS TANAH - Journal of Soil Science and Agroclimatology</i>, 18(1), 58-64. http://dx.doi.org/10.20961/stjssa.v18i1.50379	Formatted Formatted: English (United States)
Hernández, R; Olivares, B., Arias, A; Molina, J.C., Pereira, Y. (2018b). Identification of potential agroclimatic zones for the production of onion (<i>Allium cepa</i> L.) in Carabobo, Venezuela. <i>Journal of the Selva Andina Biosphere</i>, 6 (2): 70-82. http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf	Olivares, B., Pitti, J., Montenegro, E. (2020). Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. <i>Revista Brasileira de Gestao e Desenvolvimento Regional</i>, 16(3):59-71. https://doi.org/10.54399/rbdr.v16i3.5871	Formatted Formatted: English (United States)
Hernández, R. Olivares, B., (2019). Ecoterritorial sectorization for the sustainable agricultural production of potato (<i>Solanum tuberosum</i> L.) in Carabobo, Venezuela. <i>Agricultural Science and Technology</i>, 20(2): 339-354. https://doi.org/10.21930/rcta.vol20_num2_art:1462	Parra, R., Olivares, B., Cortez, A. v Rodríguez, M.F. Pluviometric homogeneity patterns in climatic stations of Anzoátegui state, Venezuela. <i>Revista Multidisciplinaria</i>, 2012, 12 (Extraordinario): 11-17. https://n9.cl/xbslq	Formatted Formatted
Kumar, Y., & Rattanpal, H. S. (2010). Effect of pruning in guava planted at different spacings under Punjab conditions. <i>Indian Journal of Horticulture</i>, 67(4), 115-119.	Pillania, S., Shukla, A. K., Mahawer, L. N., Sharma, R., & Bairwa, H. L. (2010). Standardization of pruning intensity and integrated nutrient management in meadow orcharding of guava (<i>Psidium guajava</i>). <i>Indian Journal of Agricultural Sciences</i>, 80(8), 673.	Formatted Formatted
Lian, H. N., Singh, B., Senjam, B. D., & Ramjan, M. (2019). Effect of Shoot Pruning on Growth and Yield of Guava (<i>Psidium guajava</i> L.) cv. L-49 under Foothills of Arunachal Pradesh. <i>Int. J. Curr. Microbiol. App. Sci</i>, 8(3), 2020-2027.	Pitti, J; Olivares, B; Montenegro, E. (2021). The role of agriculture in the Changuinola District: a case of applied economics in Panama. <i>Tropical and Subtropical Agroecosystems</i>, 25 - 1, 1 - 11. http://dx.doi.org/10.56369/tsaes.3815	Formatted Formatted
Lopez, J. G. V., Manica, I., Koller, O. C., & Ribold, I. J. (1982). Effect of six pruning periods on the yield of guava in Novo Hamburgo, Rio Grande do Sul, Brazil. In <i>Proc. Trop. Reg. Am. Soc. Hort. Sci</i> (Vol. 25, pp. 259-62).	Prakash, D. P., Narayanaswamy, P., & Sundur, S. N. (2002). Analysis of molecular diversity in guava using RAPD markers. <i>The Journal of Horticultural Science and Biotechnology</i>, 77(3), 287-293.	Formatted Formatted
Maji, S., Das, B. C., & Sarkar, S. K. (2015). Efficiency of some chemicals on crop regulation of Sardar guava. <i>Scientia Horticulturae</i>, 188, 66-70.	Quijada, O., Araujo, F., Corzo, P.(1999). Effect of pruning and hydrogen cyanamide on bud break, flowering, fruit yield and quality of guava (<i>Psidium guajava</i> L.) in the municipality of mara, state of zulía. <i>Revi.de la Facultad de Agro. Universidad del Zulía</i>, 16, 276-291.	Formatted Formatted Formatted
Meena, K. R., Maji, S., Kumar, S., & Verma, S. (2016). Influence Of Shoot Pruning For Crop Regulation And Improving Fruit Yield Of Guava. <i>The Bio scan</i>, 11,1355-1359.	Reddy, B. M. C. (2004, November). High density planting in horticultural crops. In <i>First Indian Horticulture Congress</i> (Vol. 6, p. 36).	Formatted Formatted
Mehta, S., Singh, S. K., Das, B., Jana, B. R., & Mali, S. (2012). Effect of pruning on guava cv. Sardar under ultra high density orcharding system.	Shaban, A. E. A., & Haseeb, G. M. M. (2009). Effect of pruning severity and spraying some chemical substances on growth and fruiting of guava trees. <i>American-Eurasian Journal of Agricultural and Environmental Science</i>, 5(6), 825-831.	Formatted Formatted Formatted
Mitra, S. K., Sen, S. K., Maiti, S. C., & Bose, T. K. (1984). Effect of plant density on growth, yield and fruit quality in guava. <i>Bangladesh Hort</i>, 12, 7-9.	Singh, G., & Chanana, Y. R. (2005). Influence of pruning intensity and pruning frequency on vegetative and reproductive attributes in guava 'L-49'. In <i>L-49 Abstract: 1st Inter Guava Symposium, CASH, Lucknow Pp</i> (Vol. 52).	Formatted Formatted
Montenegro, E., Pitti-Rodríguez, J., Olivares-Campos, B. (2021). Identification of the main subsistence crops of Teribe: a case study based on multivariate techniques. <i>Idesia (Arica)</i>, 39(3), 83-94. https://dx.doi.org/10.4067/S0718-34292021000300083	Singh, G., Singh, A. K., & Rajan, S. (2001). Influence of pruning date on fruit yield of guava (<i>Psidium guajava</i> L.) under subtropics. <i>J. Appl. Hort</i>, 3(1), 37-40.	Formatted Formatted
Norton, M. (2001, July). Fruit wood rejuvenation by reducing tree height and shoot removal year 2 progress report. In <i>V International Peach Symposium</i> 592 (pp. 401-403).		Formatted Formatted

Singh, I. S., Singh, H. K., & Chauhan, K. S. (1980). Effect of high and low density plantation on yield and quality of guava under semi-arid conditions. *Journal of research-Haryana Agricultural University*.

Viloria, J.A.; Olivares, B.O.; García, P.; Paredes-Trejo, F.; Rosales, A. (2023). Mapping Projected Variations of Temperature and Precipitation Due to Climate Change in Venezuela. *Hydrology*, 10, 96. <https://doi.org/10.3390/hydrology10040096>

Vega A, Olivares B, Calderón MAR, Rev JC, Lobo D, Gómez JA, Landa BB. (2022). Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. *Plants*, 11(15):2070. <https://doi.org/10.3390/plants11152070>

Zingaretti, M.L., Olivares, B., Demey Zambrano, J.A. y Demey, J.R. (2016). Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. *Revista FAVE -Ciencias Agrarias*, 15 (2): 39-50. <https://doi.org/10.14409/fa.v15i2.6587>

Formatted: No underline, English (United States)

Formatted: Font: Times New Roman, 9 pt, No underline, Font color: Custom Color(RGB(34,34,34)), English (United States), Pattern: Clear (White)

Formatted: No underline, English (United States)

Formatted: Font: Times New Roman, 9 pt, No underline, Font color: Custom Color(RGB(34,34,34)), English (United States), Pattern: Clear (White)

Formatted: English (United States)

Formatted: English (United States)

Formatted: No underline, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: Font: Italic, No underline, English (United States)

Formatted: No underline, English (United States)

Formatted: English (United States)

Formatted: English (United States)

UNDER PEER REVIEW