

## Original Research Article

### **Studies on heterosis for fruit yield and quality attributing characters in yardlong bean (*Vigna unguiculata* (L.) Walp. ssp. *Sesquipedalis* Verdc.)**

#### **Abstract**

A set of 29 genotypes including 7 parents, their 21 resultant hybrids and one commercial check variety (Arka Mangala) were sown during summer, 2019 at the College Instructional Farm, College of Horticulture, Venkataramannagudem to study the magnitude of heterosis using half diallel analysis for eleven characters. Significant differences were observed among the parents and hybrids indicating considerable genetic variation among these genotypes. Significant standard heterosis and high *per se* performance with regards to fruit yield per plant were recorded by the crosses *viz.*, Babli x Lola , Babli x Bobbili Local, Babli x Trivendrum Local, Bobbili Local x Lola, Bobbili Local x Trivendrum Local and Lola x Trivendrum Local.

**Keywords:** Yardlong bean, Heterosis, Breeding programmes, Yield, Quality

#### **Introduction**

Yardlong bean (*Vigna unguiculata* (L.) Walp. ssp. *sesquipedalis* Verdc.) belongs to the family Fabaceae with chromosome number  $2n=2x=22$ . Yardlong bean has a typical leguminous flower and is predominantly a self pollinated crop. However, cross pollination up to 10 percent has been reported [4]. It is a distinct form of cowpea grown as a vegetable crop throughout Asia especially in South and South East Asian countries including India. It is also known as asparagus bean, chinese long bean, pea bean, string bean, snake bean, snake pea, snap pea, borbatu *etc.*, in different parts of the world.

Area under vegetable crops is mostly covered by the F<sub>1</sub> hybrids which are in great demand because of manifold advantages such as high yield, quality, uniformity and biotic/abiotic stress tolerance and yardlong bean is no exception. However, there are no F<sub>1</sub> hybrids of yardlong bean released from either public sector or private organizations and only open pollinated varieties are being cultivated commercially. Thus, there is a need to test the scope for developing such hybrids in yardlong bean which can stand test of time for yield and quality.

### **Materials and Methods**

The experimental material consisted of seven parental lines *viz.*, Geethika, Babli, Vizianagaram Local, Bobbili Local, Lola, Trivendram Local and Bhuvanesarwar Local which were crossed in diallel fashion excluding reciprocals during *Rabi*, 2018. The resultant 21 F<sub>1</sub> hybrids along with seven parents and one check were evaluated in randomized block design with three replications with spacing of 1.0 x 1.0 m during *Summer*, 2019. Observations were recorded on five randomly selected plants from each plot for yield and quality attributing traits *viz.*, number of clusters per plant, number of pods per cluster, number of pods per plant, pod length (cm), pod girth (cm), pod weight (g), number of seeds per pod, 100 seed weight (g), pod yield per plant (kg), TSS (<sup>o</sup>Brix) and protein (%).

### **Results and Discussion**

The exploitation of heterosis in crop plant is regarded as one of the major breakthrough in the field of plant breeding. The application of heterosis is considered to an outstanding application of principles of genetics to agriculture. The scope of exploitation of hybrid vigour depends on direction and magnitude of heterosis and type of gene action involved.

In the analysis of mean squares the differences due to the treatments were significant for all the characters studied. The treatment means were further divided into parents, hybrids

and parents versus hybrids. The parents and hybrids were significant for all the characters studied while parents versus hybrids were significant for all the characters studied except for number of seeds per pod (Table-1).

For the character number of clusters per plant, significant positive heterosis was observed in 3 crosses over the mid parent and 4 hybrids were significantly superior over better parent. Four hybrids were significantly superior by exhibiting significant positive heterosis over check Arka Mangala. These results are in conformity with the findings of [8] and [7] in mung bean. Seven hybrids expressed significant positive relative heterosis. The heterobeltiosis was positively significant for only one cross and fourteen hybrids recorded significant positive standard heterosis and proved superior to the check Arka Mangala for the character number of pods per cluster. These results are in conformity with the findings of [17] and [7] in mung bean. For the character number of pods per plant, significant positive heterosis was observed in 16 hybrids over the mid parent and in 13 hybrids over better parent. Fourteen hybrids were significantly superior by exhibiting significant positive heterosis over check Arka Mangala. These results are in conformity with the findings of [16] in cowpea, [2] in mung bean and [13] in chick pea (Table-2).

For the character pod length, significant positive heterosis was observed in 14 hybrids over the mid parent and 4 hybrids exhibited significant superiority over better parent. Three hybrids were significantly superior to the check Arka Mangala by exhibiting significant positive heterosis. These results are in conformity with the findings of [11] and [7] in mung bean. For pod girth, six hybrids and three hybrids showed significant positive relative heterosis and heterobeltiosis respectively. Eight hybrids were significantly superior by exhibiting significant positive heterosis over check Arka Mangala. Similar findings were reported by [5] in velvet bean

and [9] in mung bean. For pod weight, twenty hybrids exhibited significant positive relative heterosis and twelve hybrids exhibited significant and positive heterosis over better parent. All 21 hybrids registered significant and positive standard heterosis and were found superior over Arka Mangala. Superiority of this trait was earlier observed in cowpea and other vegetables by [15] in cowpea, [3] and [14] in Indian bean which support the findings of the investigation (Table-3).

Seven, one and eight hybrids were significantly superior by exhibiting significant positive heterosis over mid, better parents and Arka Mangala respectively for the character number of seeds per pod. These results are in conformity with the findings of [8] and [7] in mung bean. Seventeen, nine and seven hybrids exhibited significant positive heterosis for 100 seed weight over mid parent, better parent and Arka Mangala (check) respectively. These findings are in accordance with earlier reports of [6] in cowpea, [12] in mung bean. For pod yield per plant, significant positive heterosis was observed in all 21 hybrids over the mid parent and over check Arka Mangala and in 20 hybrids over better parent. These results are in conformity with findings of [7] in mung bean and [14] in Indian bean (Table-4).

For character TSS, Three, one and three hybrids recorded significant positive heterosis mid, better parents and Arka Mangala respectively. The results are in conformity with earlier reports of [10] in pea. For protein content, Three, one and two hybrids significantly recorded high heterosis over mid, better parents and Arka Mangala respectively. The superiority of this trait also was observed by [1] in cowpea, [8] and [7] in mung bean (Table-5).

## **Conclusion**

The cross combinations, Babli x Lola, Babli x Bobbili Local, Babli x Trivendrum Local, Bobbili Local x Lola, Bobbili Local x Trivendrum Local and Lola x Trivendrum Local recorded highest

pod length, pod girth, pod weight, seeds per pod, 100 seed weight, TSS and protein content which contributed towards highest pod yield per plant.

## References

- [1] K. R. Anitha, K. Thiyagarajan, S. P. Bharathi and R. Rajendran (2016). Heterosis for yield and its components in fodder cowpea (*Vigna unguiculate* (L.) Walp.). *Electronic Journal of Plant Breeding.*, **7(4)**: 1208-1215.
- [2] R. T. Bhavani, R. M. Chauhan, J. R. Patel and K. Dixita (2017). Heterosis studies for grain yield and its related characters in mungbean (*Vigna radiata* (L.). *International Journal of Genetics.*, **9(2)**: 232-236.
- [3] D. Borwal, B. H. Kale, A. P. Bhuriya, K. G. Modha, K. S. Mungra and D. A. Chauhan (2018). Heterosis, inbreeding depression for seed yield and its component traits in Indian bean (*Lablab purpureus* (L.) Sweet). *International Journal of Chemical Studies.*, **6(5)**: 1955-1958.
- [4] G. Carlos, D. D. Oliveria and O. S. Ohashi (1998). Pollinator contribution to the production of cowpea in the Amazon. *Hort Science.*, **33(7)**: 1157-1159.
- [5] A. Chinapolaiah, A. H. Bindu, G. N. Manjesh, N. H. Rao, S. S. Kumar and T. V. Kumar (2019). Heterosis and combining ability analysis for yield and yield contributing traits in velvet bean (*Mucuna pruriens* L.). *Legume Research.*, **42(1)**: 10-17.
- [6] P. Hiral, J. B. Patel, S. C. Sharma and S. Acharya (2013). Heterosis and inbreeding depression study in cowpea (*Vigna unguiculate* (L.) Walp.). *An international e-Journal.*, **2(2)**: 165-172.
- [7] S. S. Kakde, A. B. Gawate and S. V. Mandge (2019). Heterosis for yield component traits and seed yield in mungbean (*Vigna radiata* (L.) Wilczek). *Journal of Pharmacognosy*

- and Phytochemistry., **8(2)**: 296-300.
- [8] S. Kalpana, N. V. Naidu and D. M. Reddy (2018). Heterosis for yield, yield components and nutritional traits in mungbean (*Vigna radiata* (L.). International Journal of Current Microbiology and Applied Sciences., **7(11)**: 1780-1791.
- [9] E. B. Khaimichho, L. Hijam, K. K. Sarkar and S. Mukherjee (2016). Estimation of combining ability and heterosis for yield related characters with protein content in seed of mungbean. Electronic Journal of Plant Breeding., **7(4)**: 849-856.
- [10] S. Kumar, V. Katoch, A. Bharti, S. Sharma, A. Sharma and V. Kumari (2019). Heterosis, inbreeding depression and combining ability studies in garden pea (*Pisum sativum* L.). Legume Research., 1-7.
- [11] P. Mehta and G. M. Lal (2018). Heterosis and genetic variability for yield related attributes in blackgram (*Vigna mungo* (L.) Hepper). Global Journal of Bio-Science and Biotechnology., **7(3)**: 383-391.
- [12] A. Nath and S. R. Maloo (2017). Heterosis studies for yield and yield component in greengram (*Vigna radiata* (L.) Wilczek). The Bioscan., **12(2)**: 1247-1252.
- [13] P. R. Sasane, A. W. Thorat, M. P. Meshram, A. R. Rajane and A. N. Patil (2018). Heterosis for seed yield and yield contributing characters on Kabuli x Kabuli crosses of chickpea (*Cicer arietinum* L.). International Journal of Chemical Sciences., **6(5)**: 1443-1450.
- [14] S. Sen, R. Patel, K. G. Modha and K. Gautam (2019). Analysis of Heterosis in Indian bean (*Lablab purpureus* (L.) Sweet). Journal of Pharmacognosy and Phytochemistry., **2**: 968-973.
- [15] D. Sharma, N. Mehta, J. Singh and C. R. Gupta (2013). Genetic study for some pod

- characters in vegetable cowpea (*Vigna unguiculate* (L.) Walp.). *Vegetable Science.*, **40(1)**: 73-76.
- [16] N. Sharma, R. K. Mittal, V. K. Sood and S. Sharma (2019). Heterosis for earliness, yield and quality traits in uradbean (*Vigna mungo* (L.) Hepper). *International Journal of Pure Applied Biosciences.*, **7(2)**: 190-197.
- [17] R. L. Srivastava and G. Singh (2013). Heterosis for yield and its contributing characters in mungbean (*Vigna radiata* (L.) Wilczek). *Indian Journal of Science and Research.*, **4(1)**: 131-134.

UNDER PEER REVIEW

**Table 1.** Analysis of variance for yield and quality traits 7 x 7 half diallel in yardlong bean

Source	Df	Number of clusters per plant	Number of pods per cluster	Number of pods per plant	Pod length (cm)	Pod girth (cm)	Pod weight (gm)	Number of seeds per pod	100 seed weight (gm)	Pod yield per plant (kg)	TSS ( <sup>o</sup> brix)	Protein (%)
<b>Treatments</b>	27	10.931**	0.512**	867.170**	204.942**	0.169**	24.500**	7.671**	16.116**	1.091**	0.396**	0.309**
<b>Parents</b>	6	3.795**	0.827**	342.725**	279.403**	0.076**	22.599**	8.477**	17.538**	0.279**	0.112**	0.289**
<b>Hybrids</b>	20	13.284**	0.399**	847.359**	186.168**	0.199**	11.860**	7.799**	15.010**	0.690**	0.346**	0.299**
<b>Parent Vs.Hybrids</b>	1	6.671**	0.869**	4410.070**	133.664**	0.114**	288.707**	0.267	29.726**	13.968**	3.085**	0.620**
<b>Error</b>	54	0.709	0.081	16.869	2.056	0.005	1.400	0.281	0.204	0.022	0.006	0.008

\*\* 1% level of significance, \* 5% level of significance

**Table 2.** Estimates of relative heterosis (RH), heterobeltiosis (Hb) and standard heterosis (SH) for number of clusters per plant, number of pods per cluster and number of pods per plant in yardlong bean

Cross combinations	Number of clusters per plant			Number of pods per cluster			Number of pods per plant		
	RH	Hb	SH	RH	Hb	SH	RH	Hb	SH
Geethika x Babli	0.73	-1.28	-1.43	-0.68	-12.05 **	17.74 **	10.78 **	-1.14	17.13 **
Geethika x Vizianagaram Local	-1.85	-3.32	-7.32 *	12.70 *	10.94	14.52 *	12.53 **	10.73 **	2.97
Geethika x Bobbili Local	-8.58 **	-13.82 **	-6.69	3.03	0.00	9.68	-0.13	-7.54 *	0.94
Geethika x Lola	-11.54 **	-14.55 **	-12.10 **	0.00	-4.29	8.06	2.80	-1.78	0.27
Geethika x Trivendrum Local	-8.82 **	-10.29 **	-11.15 **	-2.16	-9.33	9.68	0.81	-6.18	1.28
Geethika x Bhuvanewar Local	-9.40 **	-10.30 **	-14.01 **	2.40	0.00	3.23	6.52	5.44	-1.96
Babli x Vizianagaram Local	-2.89	-6.22	-6.37	-13.10 **	-24.10 **	1.61	-3.10	-14.74 **	1.01
Babli x Bobbili Local	12.01 **	7.65 *	16.56 **	7.28	-2.41	30.65 **	31.64 **	26.47 **	49.83 **
Babli x Lola	15.95 **	14.24 **	17.52 **	9.80 *	1.20	35.48 **	34.64 **	25.33 **	48.48 **
Babli x Trivendrum Local	9.53 **	9.09 **	8.92 **	1.27	-3.61	29.03 **	26.27 **	20.66 **	42.95 **
Babli x Bhuvanewar Local	-1.07	-3.99	-4.14	-5.56	-18.07 **	9.68	15.64 **	2.28	21.17 **
Vizianagaram Local x Bobbili Local	-7.91 **	-14.41 **	-7.32 *	10.77 *	5.88	16.13 **	20.38 **	9.82 **	19.89 **
Vizianagaram Local x Lola	-6.67 *	-11.15 **	-8.60 *	4.55	-1.43	11.29	16.39 **	9.51 **	11.80 **

Vizianagaram Local x Trivendrum Local	-13.10 **	-15.76 **	-16.56 **	6.57	-2.67	17.74 **	8.72 **	-0.31	7.62 *
Vizianagaram Local x Bhuvanewar Local	-11.41 **	-11.86 **	-17.20 **	17.07 **	16.13 **	16.13 **	11.24 **	10.58 **	0.74
Bobbili local x Lola	1.66	-0.88	7.32 *	11.59 *	10.00	24.19 **	30.10 **	25.88 **	37.42 **
Bobbili local x Trivendrum Local	-1.08	-5.29	2.55	6.29	1.33	22.58 **	21.68 **	21.00 **	32.10 **
Bobbili local x Bhuvanewar Local	-14.33 **	-20.00 **	-13.38 **	11.63 *	5.88	16.13 **	21.21 **	11.18 **	21.38 **
Lola x Trivendrum Local	2.21	0.31	3.18	7.59	4.00	25.81 **	30.79 **	27.23 **	37.36**
Lola x Bhuvanewar Local	4.53	8.67**	-6.05	14.50**	7.14	20.97**	24.12**	17.44**	19.89**
Trivendrum Local x Bhuvanewar Local	-7.26*	-9.65**	-10.51**	5.88	-4.00	16.13**	19.11**	9.81**	18.54**
<b>Range</b>	<b>-14.33 to 15.95</b>	<b>-20.00 to 14.24</b>	<b>-17.20 to 17.52</b>	<b>-13.10 to 17.07</b>	<b>-24.10 to 16.13</b>	<b>1.61 to 35.48</b>	<b>-3.10 to 34.64</b>	<b>-14.74 to 27.23</b>	<b>-1.96 to 49.83</b>

\*\* 1% level of significance, \* 5% level of significance

**Table 3.** Estimates of relative heterosis (RH), heterobeltiosis (Hb) and standard heterosis (SH) for pod length (cm), pod girth (cm) and pod weight (g) in yardlong bean

Cross combinations	Pod length (cm)			Pod girth (cm)			Pod weight (g)		
	RH	Hb	SH	RH	Hb	SH	RH	Hb	SH
Geethika x Babli	-0.03	-16.44 **	-11.35 **	-15.89 **	-20.19 **	-4.75	15.15 **	7.11 *	27.07 **
Geethika x Vizianagaram Local	44.93 **	33.31 **	-5.00 *	7.89 **	3.11	10.46 **	29.16 **	18.73 **	21.18 **
Geethika x Bobbili Local	-3.27	-17.15 **	-17.20 **	-15.51 **	-18.26 **	-6.34 *	8.80 **	2.89	17.81 **
Geethika x Lola	18.31 **	11.67 **	-10.36 **	-5.58 *	-7.52 **	3.33	10.55 **	5.66	18.31 **
Geethika x Trivendrum Local	12.80 **	4.75	-12.92 **	-12.88 **	-14.73 **	-4.60	10.14 **	7.48	15.27 **
Geethika x Bhuvanewar Local	8.21 **	2.96	-18.75 **	-14.74 **	-17.01 **	-11.09 **	13.90 **	13.42 **	15.76 **
Babli x Vizianagaram Local	0.91	-21.09 **	-16.28 **	-7.67 **	-16.07 **	0.16	12.79 **	-2.93	15.16 **
Babli x Bobbili Local	4.46 **	1.43	7.61 **	6.10 **	3.98	24.09 **	11.57 **	9.62 **	30.05 **
Babli x Lola	16.72 **	2.52	8.76 **	7.82 **	4.38	24.56 **	19.04 **	15.70 **	37.25 **

Babli x Trivendrum Local	11.13 **	-0.90	5.14 *	6.79 **	3.45	23.45 **	21.08 **	15.27 **	36.75 **
Babli x Bhuvaneswar Local	-13.80 **	-24.84 **	-20.26 **	-8.26 **	-15.14 **	1.27	11.56 **	3.36	22.62 **
Vizianagaram Local x Bobbili Local	-17.65 **	-34.17 **	-34.21 **	-17.70 **	-23.79 **	-12.68 **	19.57 **	4.46	19.61 **
Vizianagaram Local x Lola	9.23 **	-4.68	-23.48 **	-2.35	-8.51 **	2.22	19.55 **	5.46	18.09 **
Vizianagaram Local x Trivendrum Local	-6.98 **	-20.01 **	-33.50 **	-15.43 **	-20.82 **	-11.41 **	22.35 **	9.98 **	17.95 **
Vizianagaram Local x Bhuvaneswar Local	6.42 *	-6.44 *	-26.17 **	4.14	2.19	3.65	30.61 **	20.52 **	21.97 **
Bobbili local x Lola	13.93 **	2.72	2.66	7.14 **	5.81 *	21.24 **	19.14 **	17.82 **	34.90 **
Bobbili local x Trivendrum Local	6.77 **	-2.21	-2.26	4.13	2.90	17.91 **	21.10 **	17.27 **	34.26 **
Bobbili local x Bhuvaneswar Local	5.86 **	-5.28 **	-5.33 **	-1.54	-7.19 **	6.34 *	13.87 **	7.26 *	22.81 **
Lola x Trivendrum Local	21.37 **	19.28 **	-0.84	5.88 **	5.81 *	18.38 **	19.11 **	16.59 **	30.55 **
Lola x Bhuvaneswar Local	8.11**	8.88**	-26.86**	3.05	7.52**	3.33	5.99	0.89	12.97**
Trivendrum Local x Bhuvaneswar Local	-1.49	-3.99	-20.18**	-9.21**	-13.46**	-3.17	8.65**	5.59	13.24**
<b>Range</b>	<b>-17.65 to 44.93</b>	<b>-34.17 to 33.31</b>	<b>-33.50 to 8.76</b>	<b>-17.70 to 7.89</b>	<b>-23.79 to 5.81</b>	<b>-12.68 to 24.56</b>	<b>8.80 to 30.61</b>	<b>-2.93 to 20.52</b>	<b>12.97 to 37.25</b>

\*\* 1% level of significance, \* 5% level of significance

**Table 4.** Estimates of relative heterosis (RH), heterobeltiosis (Hb) and standard heterosis (SH) for number of seeds per pod, 100 seed weight (g) and pod yield per plant (kg) in yardlong bean

Cross combinations	Number of seeds per pod			100 seed weight (g)			Pod yield per plant (kg)		
	RH	Hb	SH	RH	Hb	SH	RH	Hb	SH
Geethika x Babli	-6.34**	-7.72**	-0.40	13.33**	-2.99	-1.45	20.06 **	6.95 *	34.20 **
Geethika x Vizianagaram Local	8.94**	-3.03	1.59	31.28**	21.34**	3.41	32.18 **	32.10 **	29.56 **
Geethika x Bobbili Local	-12.52**	-14.02**	-9.92**	10.36**	-1.78	-8.93**	22.41 **	12.87 **	31.14 **
Geethika x Lola	-3.95	-4.49	1.19	26.63**	13.16**	3.96*	25.81 **	20.19 **	29.45 **
Geethika x Trivendrum Local	-7.54*	-9.47**	-5.16	-1.85	-17.45**	-12.48**	29.26 **	25.60 **	30.58 **
Geethika x Bhuvaneswar Local	-12.40**	-19.70**	-15.87**	11.25**	3.33	-12.86**	30.17 **	28.85 **	28.99 **
Babli x Vizianagaram Local	-5.86*	-17.28**	-10.71**	9.83**	0.98	2.59	16.47 **	3.70	30.12 **
Babli x Bobbili Local	6.26**	2.94	11.11**	9.66**	4.88**	6.55**	39.64 **	34.48 **	68.74 **
Babli x Lola	5.38*	4.41	12.70**	10.73**	5.43**	7.11**	50.36 **	39.71 **	75.31 **
Babli x Trivendrum Local	4.38	0.74	8.73**	6.20**	3.98**	10.24**	43.44 **	31.14 **	64.55 **

Babli x Bhuvanesar Local	0.81	-8.82**	-1.59	9.46**	0.17	1.76	19.08 **	7.04 *	34.31 **
Vizianagaram Local x Bobbili Local	-14.10**	-22.35**	-21.43**	5.46**	1.19	-6.17**	25.75 **	15.89 **	34.65 **
Vizianagaram Local x Lola	-3.17	-14.23**	-9.13**	8.85**	4.92*	-3.62*	22.36 **	16.82 **	25.82 **
Vizianagaram Local x Trivendrum Local	-12.85**	-20.95**	-20.63**	-5.93**	-15.16**	-10.05**	23.39 **	19.83 **	24.58 **
Vizianagaram Local x Bhuvanesar Local	10.80**	7.27*	-6.35*	18.17**	17.55**	0.19	28.87 **	27.49 **	27.63 **
Bobbili local x Lola	3.45	1.12	7.14**	17.57**	17.03**	8.51**	44.26 **	38.99 **	61.49 **
Bobbili local x Trivendrum Local	2.76	2.35	3.57	8.26**	1.47	7.57**	43.21 **	35.67 **	57.64 **
Bobbili local x Bhuvanesar Local	6.95**	-0.39	0.79	-10.67**	-14.71**	-20.91**	16.54 **	8.48 *	26.05 **
Lola x Trivendrum Local	1.15	-1.50	4.37	7.02**	-0.12	5.89**	39.43 **	37.01 **	47.57 **
Lola x Bhuvanesar Local	5.13*	4.12	1.59	9.42**	13.13**	-20.20**	30.38**	26.08**	35.79**
Trivendrum Local x Bhuvanesar Local	6.55**	-0.40	0.00	-20.13**	-28.30**	-23.98**	19.76**	17.54**	22.20**
<b>Range</b>	<b>-14.10 to 10.80</b>	<b>-22.35 to 7.27</b>	<b>-21.43 to 12.70</b>	<b>-20.13 to 31.28</b>	<b>-28.30 to 21.34</b>	<b>-23.98 to 10.24</b>	<b>16.47 to 50.36</b>	<b>3.70 to 39.71</b>	<b>24.58 to 68.74</b>

\*\* 1% level of significance, \* 5% level of significance

**Table 5.** Estimates of relative heterosis (RH), heterobeltiosis (Hb) and standard heterosis (SH) for TSS (<sup>0</sup>brix) and protein content (%) in yardlong bean

Cross combinations	TSS ( <sup>0</sup> brix)			Protein content (%)		
	RH	Hb	SH	RH	Hb	SH
Geethika x Babli	-17.30 **	-18.92 **	-8.36 **	-16.90 **	-21.63 **	-16.11 **
Geethika x Vizianagaram Local	-16.76 **	-19.30 **	-12.38 **	0.51	-1.01	-6.09 *
Geethika x Bobbili Local	-17.89 **	-22.90 **	-16.29 **	-21.49 **	-23.72 **	-23.27 **
Geethika x Lola	-19.66 **	-20.80 **	-11.51 **	-6.45 **	-8.53 **	-9.19 **
Geethika x Trivendrum Local	-26.34 **	-27.06 **	-19.22 **	-18.62 **	-23.01 **	-18.14 **
Geethika x Bhuvanesar Local	-17.99 **	-18.40 **	-11.40 **	-8.67 **	-18.49 **	-22.67 **
Babli x Vizianagaram Local	-20.40 **	-24.30 **	-14.44 **	-12.23 **	-18.39 **	-12.65 **
Babli x Bobbili Local	-7.76 **	-14.99 **	-3.91	-1.84	-4.79	1.91
Babli x Lola	-12.75 **	-13.26 **	-1.95	0.29	-3.34	3.46
Babli x Trivendrum Local	2.18	1.15	14.33 **	0.56	0.22	7.28 **
Babli x Bhuvanesar Local	-16.69 **	-18.73 **	-8.14 **	-6.77 **	-20.96 **	-15.39 **
Vizianagaram Local x Bobbili Local	-7.87 **	-10.86 **	-9.12 **	-24.16 **	-27.40 **	-26.97 **

Vizianagaram Local x Lola	-35.37 **	-38.19 **	-30.94 **	-19.15 **	-22.12 **	-22.67 **
Vizianagaram Local x Trivendrum Local	-24.35 **	-27.35 **	-19.54 **	-29.00 **	-33.78 **	-29.59 **
Vizianagaram Local x Bhuvanewar Local	-14.26 **	-16.46 **	-10.21 **	-4.95	-14.01 **	-20.88 **
Bobbili local x Lola	9.39 **	1.36	13.25 **	5.91 *	5.22	5.85 *
Bobbili local x Trivendrum Local	7.48 **	0.00	10.75 **	-5.77 *	-8.31 **	-2.51
Bobbili local x Bhuvanewar Local	1.71	-4.04 *	3.15	9.48 **	-4.74	-4.18
Lola x Trivendrum Local	-15.37 **	-15.74 **	-5.86 **	-1.57	-4.83	1.19
Lola x Bhuvanewar Local	18.57**	20.12**	-10.75**	2.75	10.10**	-10.74**
Trivendrum Local x Bhuvanewar Local	-11.64**	-12.94**	-3.58	6.14*	-9.76**	-4.06
<b>Range</b>	<b>-35.37 to 18.57</b>	<b>-38.19 to 20.12</b>	<b>-30.94 to 14.33</b>	<b>-29.00 to 9.48</b>	<b>-33.78 to 10.10</b>	<b>-29.59 to 7.28</b>

\*\* 1% level of significance, \* 5% level of significance

UNDER PEER REVIEW