

Characterization of soil properties and their relationship with various sulphur fractions in Groundnut Farming Regions of Bikaner District

Abstract:

An exploratory survey was conducted in Bikaner district during the 2015-16 kharif season to study the sulfur fractions and other soil properties in groundnut growing areas. 40 farmers from four tehsils (Lunkaransar, Shri Dungargarh, Nokha, and Kolayat) were selected for the study. Soil samples were collected and analyzed for pH, EC, CaCO₃, organic carbon, available nutrients (N, P₂O₅, K₂O, S), and sulfur fractions. The results showed that the soils were sandy, loamy sand, and sandy loam in texture and alkaline in reaction. The majority of soils were calcareous with low organic carbon content. The available nitrogen was found to be low while available P₂O₅ and K₂O were medium. The distribution of different forms of sulfur in soil was found to be strongly dependent on soil characteristics such as clay content, silt and organic carbon content.

Keywords: Sulphur, Fraction, pH, Survey, soil Quality

Introduction

Sulphur is considered the fourth most important plant nutrient after nitrogen, phosphorus, and potassium. It plays a crucial role in quality crop production. The intensification of agriculture, including the use of high-yielding varieties, multiple cropping, and increased irrigation, along with the increased use of chemical fertilizers low in sulphur and reduced sulphur dioxide emissions from industries (Shah *et al.*, 2022; Abdallah *et al.*, 2010), has created a significant gap between the amount of sulphur added to soils and the demand for sulphur by crops. This situation is further exacerbated by the loss of sulphur or its adsorption in soils. Coarse-textured soils, which are highly permeable, are particularly susceptible to sulphur deficiency (Carlson and Ramaswami, 1992).

The positive response of crops to sulphur application is a clear indication of its deficiency in soils (Raza *et al.*, 2018). To maintain high levels of soil fertility and crop productivity, the amount of sulphur removed by crops and lost through leaching must be replenished through sulphur application (Ramdevputra *et al.*, 2010). Otherwise, sulphur deficiencies may pose an immediate threat to targeted food production. The increasing prevalence of sulphur deficiency in crops across different soils in the country, and the

positive response to sulphur application from various sources, highlights the importance of this nutrient in crop production (Joshi *et al.*, 2021).

Sulphur plays a crucial role in the formation of amino acids such as cystine, cysteine, and methionine, as well as in the oil content of oilseeds and the nutritive quality of forages. It is involved in the synthesis of certain vitamins (B1, biotin, and thiamine), the metabolism of carbohydrates, proteins, and oils, and the formation of flavour compounds in crucifers. Sulphur also contributes to the market quality of produce from several crops (Pagare *et al.*, 2015; Hawkesford and De Kok 2006; Abrol and Ahmed, 2003).

Sulphur is also a constituent of glutathione, a compound that plays a crucial role in plant respiration. Additionally, sulphur is essential for chlorophyll formation and the synthesis of protein building blocks (Juszczuk and Ostaszewska, 2011). Research has shown that sulphur promotes nodulation in legumes and is responsible for the development of large grains in oilseeds. In the absence of sufficient sulphur, essential enzymatic activities and physiological functions are inhibited, leading to reduced crop quantity and quality. Sulphur deficiency in soil results in weak, stunted plants with pale green to yellow coloration, weak stems, and delayed maturity, leading to significant economic losses (Gates, 1974). Sulphur deficiency has been shown to reduce crop yields by up to 35% (Pal and Singh, 1992).

In India, it is estimated that growing crops remove 1.8 million tonnes of sulphur per year, while only 0.8 million tonnes are added through fertilizers, resulting in an annual deficit of 1.0 million tonnes (Gupta, 2006). In intensive cropping systems, sulphur removal ranges from 30 to 70 kg ha⁻¹ per annum. Therefore, maintaining optimal levels of sulphur in soil relative to other nutrients is crucial for achieving maximum crop production and quality (Scherer, 2001).

Sulphur exists in soil in various forms, including free and adsorbed sulphate, as well as diverse organic and inorganic compounds. In humid regions, sulphur is predominantly present in organic form, while in arid soils, sulphate salts of calcium, magnesium, sodium, and potassium predominate (Kanwar 1976). Organically bound sulphur can be divided into two groups: carbon-bonded sulphur, which includes the sulphur of amino acids, and non-carbon-bonded sulphur, which includes phenolic and choline sulphates as well as lipids. The inorganic forms of sulphur in soil consist mainly of SO₄-S.

In soil, sulphur can be broadly grouped into five forms: total-S, organic-S, non-sulphate-S, available-S, and water-soluble-S. The term “available sulphur” refers to water-soluble sulphur, adsorbed sulphur, and easily hydrolyzable organic sulphur compounds. The amount and availability of various forms of sulphur vary depending on soil physicochemical

characteristics such as texture, pH, calcium carbonate content, and organic matter content. The distribution of different forms of sulphur and their relationship with important soil characteristics determine the sulphur-supplying power of soil by influencing its release and dynamics (Balanagoudar and Satyanarayana, 1990b; Das et al., 2011). Therefore, assessing the status of different forms of sulphur in soil is essential for improving crop sulphur nutrition.

In arable soils, sulphur occurs mostly in organic forms and partially in inorganic forms. Organic sulphur must be mineralized to sulphate-S to become available to plants (Tabatabai, 1982). Thus, the supply of sulphur to plants from a given soil depends on the inorganic sulphate content of the soil and the rate of mineralization of organic sulphur. The fraction of inorganic sulphur present in soils depends on several soil properties and climatic conditions. Sulphur availability is influenced by various soil factors, resulting in wide variation in the status of different forms of sulphur across different soils (Balanagoudar and Satyanarayana, 1990b).

The amount and forms of sulphur in soils are largely determined by the mineralogical composition of the parent material, the degree of weathering, and the mechanical composition of the soil. In the studied area, the northwestern part of Rajasthan, soils are coarse-textured and low in organic matter content. Since most of the sulphur in soil is present in organic form, the total reserve of sulphur in such soils is deficient. This problem is further exacerbated by high temperatures, which result in rapid mineralization and subsequent leaching losses of sulphate due to the low retention capacity of alkaline, coarse-textured sandy soils. This current study was thus planned to evaluate the distribution of soil sulphur in the *Torripsamments* (Sehgal, 1996) of groundnut growing areas of Bikaner.

Materials and Methods:

An exploratory survey of groundnut growing areas in Bikaner district was conducted during the 2015-16 kharif season. 40 farmers, 10 from each tehsil (Lunkaransar, Shri Dungargarh, Nokha, and Kolayat), growing groundnut for 3+ years were selected. Initial composite soil samples were collected for analysis of sulfur fractions and other soil properties. Data on nutrient application, management practices, and average yield were also collected. Soil samples were analyzed for pH, EC, CaCO₃, organic carbon, available nutrients (N, P₂O₅, K₂O, S), and sulfur fractions using methods described in Table 1.

Table 1: The details of methods and the procedure for these standard methods.

S. No.	Properties	Procedure	Reference
1	pH (1:2.5 Soil water)	Using glass electrode pH meter	Richards (1953)
2	EC (1:2.5 Soil water)	Using standard precision conductivity bridge	Richards (1953)
3	Organic carbon (%)	Wet digestion using normal solution of chromic acid and titration with 0.5 N FAS in the presence of ferroin indicator	Walkley and Black (1934)
4	Particle size analysis (mm)	Hydrometer method	Bouyoucos (1962)
5	CaCO ₃	Rapid titration method using EDTA	Bascomb, (1964)
6	Available N	Alkaline KMnO ₄ Method, in micro kjeldahl in the presence of NaOH (2.5%)	Subbaih and Asija (1956)
7	Available P ₂ O ₅	Extractant: 0.5 M NaHCO ₃ (pH 8.5), Estimation: Colorimetric	Olsen et al. (1954)
8	Available K ₂ O	Using neutral normal ammonium acetate as extractant and measuring K using flame photometer	Jackson (1973)
9	Available S	Extracted by 0.15% CaCl ₂ and analysed using spectrometric method in the presence of BaCl ₂	Chesnin and Yien (1950)
10	Organic sulfur	Extracted by NaH ₂ PO ₄ and 2N Acetic Acid Solution	Bardsley and Lancaster (1960)
11	Water Soluble S	Extracted by shaking with distilled water	Freney (1957)
12	Heat Soluble sulfur	Extracted with 1% NaCl	Williams and Steinbergs (1959)
13	SO ₄ Sulfur	By 0.15% CaCl ₂ , by phosphate extraction	Williams and Steinbergs

		(500 ppm P KH_2PO_4), Sulfur will be estimated by turbidimetric method.	(1959), Ensminger (1954), Chesnin and Yien (1950)
14	Total sulfur	By Acid digestion method	Tabatabai (1982)
15	Non SO_4 sulfur	Computed by subtracting the sum of organic S and SO_4 -S from Total-S	

Correlation-regression analyses were conducted to determine the relationships between different soil sulfur fractions and important soil properties. The critical limit of sulfur in soil for groundnut in the studied area was determined using the Cate and Nelson technique (1972).

Results:

a. Textural classification:

Particle size analysis data (Table 1) showed that sand content in Kolayat tehsil soils ranged from 77.19% to 88.42%, with a mean of 83.77%. Silt content ranged from 4.22% to 9.66%, with a mean of 6.51%, and clay content ranged from 7.00% to 13.75%, with a mean of 9.56%. In Lunkaransar tehsil, sand, silt, and clay content varied from 83.65% to 91.70%, 2.25% to 7.10%, and 6.05% to 9.25%, with means of 87.96%, 4.41%, and 7.65%, respectively. In Nokha tehsil, sand, silt, and clay content varied from 86.95% to 91.81%, 2.86% to 5.83%, and 7.42% to 9.02%, with means of 88.54%, 4.03%, and 7.42%, respectively. In Shri Dungargarh tehsil, sand, silt, and clay content varied from 86.90% to 91.42%, 2.53% to 5.05%, and 5.85% to 9.28%, with means of 89.01%, 3.65%, and 7.34%, respectively. The data shows that there is variation in sand, silt, and clay content across the different tehsils of Bikaner district. Sand content is generally high, ranging from 77.19% to 91.81%, with mean values ranging from 83.77% to 89.01% across the four tehsils. Silt content is generally lower, ranging from 2.25% to 9.66%, with mean values ranging from 3.65% to 6.51%. Clay content is also relatively low, ranging from 5.85% to 13.75%, with mean values ranging from 7.34% to 9.56%. Overall, the soils in the studied area are predominantly sandy, with lower levels of silt and clay.

The table 3 shows that silt and clay content were positively and significantly correlated with all sulfur fractions, while sand content was negatively and significantly correlated with all sulfur fractions. The correlation coefficients between silt, clay, and sand content and different sulfur fractions vary across the different tehsils of Bikaner. In Kolayat tehsil, silt and clay content were positively and significantly correlated with all sulfur fractions, while sand content was negatively and significantly correlated with all sulfur fractions. In Lunkaransar tehsil, silt and clay content were also positively and significantly correlated with all sulfur fractions, while sand content was negatively and significantly correlated with all sulfur fractions. In Nokha tehsil, silt content showed a positive but non-significant correlation with all sulfur fractions, while clay content was positively and significantly correlated with all sulfur fractions. Sand content was negatively and significantly correlated with all sulfur fractions.

Overall, the data suggests that in all three tehsils, as the silt and clay content of the soil increases, the levels of different sulfur fractions also increase, while as sand content increases, the levels of different sulfur fractions decrease. However, the strength of these relationships varies across the different tehsils.

- b. Calcium Carbonate: The Calcium carbonate content in overall sampling area (Table 6) is around 4.87 meq/lt. and it varies from 0.5 to 10 meq/lt. in terms of tehsil wise variation the data is presented in table. In tehsil Kolayat (Table 2), the calcium carbonate in soil was found to be in the range of 0.5 to 7 % with average values around 3.2%. the lowest value of CaCO_3 was observed in *Amaepura k-2* (0.5%) and highest was observed in *Akkasar K-7* (7%). As compared to *Lunkaransar* (Table 3) where the data varies from 3 to 8.5% and averaging around 5.6% this shows the higher concentration of salts in terms of lower and upper limits of CaCO_3 is concerned. The highest value was recorded in *Chak 277* and lowest was observed in *Chak 269* village. In *Nokha* (Table 4) the values of CaCO_3 were in the range of 0.5 and 10% with average values of 6.25% this tehsil saw the most variation in CaCO_3 concentration. The highest reading was observed in *Kakada* (10%), and least value was seen in *Berasar* (10%). Also, CaCO_3 in *Shri Dungargarh* (Table 5) varied from

1.5 to 8% averaging around 4.45%. the lowest value of CaCO_3 was seen in *Seruna D-10* while in *Derajsar D-6* the value was lowest.

The correlation coefficient of the CaCO_3 doesn't show significant correlation among the Sulphur fractions with calcium carbonate concentration. The values of correlation coefficient for different tehsils are presented in Table (7-10)

- c. pH: The pH of the sampling area (Table 6) averages out at 8.895 meaning the region falls under very high salinity, the range for the entire sampling area was observed to be 8.0 to 9.44 in *Chak 269* and *Akkasar k-10*, respectively. Tehsil wise data shows that highest pH was overserved in Nokha (Table 2) where the lowest value was observed in *Zhadeli* (8.05) and highest in *Sindhu* (9.34). In other tehsils the variation as depicted in table. Shows *Kolayat* (Table3) having highest level of salinity with data ranging from 8.68 to 9.44 in *Amarpura K-2* and *Akkasar K-10*. *Lunkaransar* (Table 4) having pH ranging from 8.0 to 9.29 in *Chak 269* and *Nathyana 9LKD*, respectively.

The correlation among pH and different fraction of Sulphur shows very high negative correlation coefficient ranging from -0.594 between pH and Sulphate sulphur of Nokha, whereas highest negative correlation was observed among pH and total sulphur fraction in soils of Kolayat. Shows that as the pH of a soil increases the total sulphur as well as other fractions of sulphur also show a decreasing trend. Table (7-10)

- d. Electrical Conductivity: The data shows that average values of EC ranges (Table 6) from 0.07 dSm^{-1} in *Goplsar D-4* to 0.20 dSm^{-1} in *Akkasar K-10* with mean values of 0.14 dSm^{-1} . EC values show no constraint of salinity in sampling area. The highest variation was observed in *Kolayat* (Table 2) where EC fluctuated between 0.09 to 0.20 dSm^{-1} . Whereas least variation in EC was seen in *Lunkaransar* (Table 3) with range between 0.12 to 0.18 dSm^{-1} . In *Nokha* (Table 4) EC ranges from 0.10 to 0.19 dSm^{-1} . In *Shri Dungargarh* (Table 5) EC varied between 0.007 to 0.16 dSm^{-1} .

Correlation Coefficient between EC and Different fractions of Sulphur in surveyed area shows very high negative values with ranging from $r=-0.712$ with heat soluble fraction in *Shri Dungargarh* area samples to $r= -0.989$

among Sulphate sulphur and EC of Kolayat samples. The Data of EC with different fraction is given in Table (7-10).

e. Organic Carbon:

The soil in four tehsils of Rajasthan – Kolayat (Table 2), Lunkaransar (Table 3), Nokha (Table 4), and Shri Dungargarh (Table5) - was analyzed for its organic carbon content. In Kolayat, the organic carbon content ranged from 0.08% to 0.21%, with an average of 0.15%. Amarpura K-2 village had the highest content while Akkasar K-6 village had the lowest. In Lunkaransar, the range was 0.10% to 0.24%, with an average of 0.17%. Chak 269 village had the highest content and Chak 277 village had the lowest. In Nokha, the range was 0.09% to 0.25%, with an average of 0.15%. Zhadeli village had the highest content while Sindhu village had the lowest. In Shri Dungargarh, the range was 0.10% to 0.34%, with an average of 0.23%. Seruna D-9 village had the highest content while Gopalsar D-5 village had the lowest.

In four tehsils of Rajasthan - Shri Dungargarh, Nokha, Lunkaransar, and Kolayat - a strong positive correlation was observed between the organic carbon content of soil and various forms of sulfur. This suggests that the presence of sulfur in the soil is closely related to its organic carbon content. Table (7-10)

f. Available Nitrogen: The available nitrogen content (Table 6) in soil was analyzed for villages in four tehsils of Rajasthan – Kolayat (Table 2), Lunkaransar (Table 3), Nokha (Table 4), and Shri Dungargarh (Table 5). In Kolayat, the available nitrogen content ranged from 100.10 to 221.10 kg/ha, with an average of 192.02 kg/ha. All villages in this tehsil belonged to the low category of soil available nitrogen. Amarpura K-2 village had the highest content while Angnaeu K-4 village had the lowest. In Lunkaransar, the range was 170.58 to 233.60 kg/ha, with an average of 205.35 kg/ha. All villages in this tehsil also belonged to the low category of soil available nitrogen. Chak 269 village had the highest content and Chak 277 village had the lowest. In Nokha, the range was 166.25 to 235.20 kg/ha, with an average of 202.03 kg/ha. All villages in this tehsil also belonged to the low category of soil available nitrogen. Zhadeli village had the highest content while Sindhu village had the lowest. In Shri Dungargarh, the range was 172.50 to 299.84 kg/ha, with an

average of 223.04 kg/ha. Seruna D-9 village had the highest content while Gopalsar D-5 village had the lowest.

The correlation between available nitrogen and various forms of sulfur was analyzed for four different sets of data. In all four sets, a positive correlation was observed between available nitrogen and total sulfur, organic sulfur, available sulfur, non-sulfate sulfur, sulfate sulfur, water-soluble sulfur, and heat-soluble sulfur. The strength of the correlation varied between different sets of data (Table 7-10).

This data suggests that there is a positive relationship between the available nitrogen content of soil and the presence of various forms of sulfur, with the strength of the correlation varying between different sets of data.

- g. Available Phosphorus: The available phosphorus content in soil was analyzed for villages in four tehsils of Rajasthan – Kolayat (Table 2), Lunkaransar (Table 3), Nokha (Table 4), and Shri Dungargarh (Table 5). In all four tehsils, the villages belonged to the medium category of soil available phosphorus. In Kolayat, the available phosphorus content ranged from 12.45 to 27.17 kg/ha, with an average of 18.39 kg/ha. Amarpura K-2 village had the highest content while Akkasar K-10 village had the lowest. In Lunkaransar, the range was 13.07 to 27.49 kg/ha, with an average of 17.56 kg/ha. Chak 269 village had the highest content and Nathvana 9LKD village had the lowest. In Nokha, the range was 14.06 to 20.93 kg/ha, with an average of 17.38 kg/ha. Zhadeli village had the highest content while Sindhu village had the lowest. In Shri Dungargarh, the range was 11.21 to 20.62 kg/ha, with an average of 14.58 kg/ha. Seruna D-9 village had the highest content while Gopalsar D-5 village had the lowest.

The correlation between available phosphorus and various forms of sulfur was analyzed for four different sets of data. In all four sets, a significant positive correlation was observed between available phosphorus and total sulfur, organic sulfur, available sulfur, non-sulfate sulfur, sulfate sulfur, water-soluble sulfur, and heat-soluble sulfur. The strength of the correlation varied between different sets of data (Table 7-10).

This data suggests that there is a strong positive relationship between the available phosphorus content of soil and the presence of various forms of sulfur.

- h. Available potassium: The available potassium content in soil was analyzed for villages in four tehsils of Rajasthan (Table 6) – Kolayat (Table 2), Lunkaransar (Table 3), Nokha (Table 4), and Shri Dungargarh (Table 5). In Kolayat, the available potassium content ranged from 124.90 to 214.20 kg/ha, with an average of 180.99 kg/ha. All villages in this tehsil belonged to the medium category of soil available potassium. Amarpura K-2 village had the highest content while Akkasar K-10 village had the lowest. In Lunkaransar, the range was 155.90 to 322.60 kg/ha, with an average of 201.35 kg/ha. Chak 269 village had the highest content and Nathvana 9LKD village had the lowest. In Nokha, the range was 142.57 to 409.44 kg/ha, with an average of 216.46 kg/ha. Zhadeli village had the highest content while Sindhu village had the lowest. In Shri Dungargarh, the range was 113.37 to 571.50 kg/ha, with an average of 187.70 kg/ha. Seruna D-9 village had the highest content while Gopalsar D-5 village had the lowest.

The correlation between available potassium and various forms of sulfur was analyzed for four different sets of data. In all four sets, a significant positive correlation was observed between available potassium and total sulfur, organic sulfur, available sulfur, non-sulfate sulfur, sulfate sulfur, water-soluble sulfur, and heat-soluble sulfur. The strength of the correlation varied between different sets of data (Table 7-10)

This data suggests that there is a strong positive relationship between the available potassium content of soil and the presence of various forms of sulfur.

Discussion:

Soil Textural Classes:

The data on particle size analysis of soils revealed that the majority of soils belonged to the loamy sand category, indicating that they were formed mostly from alluvial material and characterized by a coarse texture. The variation in soil texture might be due to factors such as topographic position, nature of parent material, in-situ weathering, and age of the soils.

Silt and clay content were found to be positively and significantly correlated with various forms of sulfur, indicating that a significant quantity of sulfur is adsorbed on the finer fractions of soil and that the availability of sulfur may increase with an increase in fineness of texture. Sand, on the other hand, showed a significant negative relationship with all forms of sulfur, suggesting that the value of all forms of sulfur decreases with an increase in sand

particles in the soil. This may be attributed to less organic carbon accumulation and high leaching.

These findings are consistent with previous research by Trivedi et al. (1998), Singh et al. (2006), Athokpam et al. (2007), and Singh et al. (2009).

CaCO₃:

The calcium carbonate (CaCO₃) content of soil samples from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed as a useful parameter to assess nutrient availability and release behavior. The CaCO₃ content varied between 0.50% to 7.00% in Kolayat, 3.00% to 8.50% in Lunkaransar, 0.50% to 10.00% in Nokha, and 1.50% to 8.00% in Shri Dungargarh. The majority of soils in the area were found to be calcareous (CaCO₃ content > 5%) according to the classification given by F.A.O. (1977).

The accumulation of CaCO₃ in these soils might be due to the semi-arid climatic conditions, where rainfall is less than annual evapo-transpiration, resulting in less water available for leaching of insoluble carbonates and bicarbonates of calcium. The correlation coefficients indicate that CaCO₃ content was negatively correlated with all forms of sulfur fractions, suggesting that non-calcareousness in soil enriches sulfur content.

These findings are consistent with previous research by Trivedi et al. (1998), Chaudhary and Shukla (2002), Deshmukh et al. (2004), Kara and Ceylan, (1988), and Sharma and Gangwar (1997).

pH:

The pH of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed. The minimum pH value was recorded in Lunkaransar while the maximum was observed in Kolayat. The pH of soils in all four tehsils indicated that they were alkaline in reaction, likely due to medium to high base saturation, high amounts of carbonate and bicarbonate, and insufficient rainfall or irrigation to leach down bases.

The pH was found to be negatively and significantly correlated with different forms of sulfur. This may be due to the presence of H⁺ and OH⁻ ions on the soil complex, where H⁺ ions attract SO₄²⁻ ions. These findings are consistent with previous research by Singh et

al. (2000), Singh et al. (2009), Lal and Singh (1974), Bhargava and Sharma (1982), More et al. (1988), Sharma and Gangwar (1997), Singh et al. (2006), Jat and Yadav (2006), and Chaudhary and Shukla (2002).

EC:

The electrical conductivity (EC) of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed as an indicator of total soluble salts. The minimum EC value was recorded in Kolayat while the maximum was observed in Shri Dungargarh. The EC values indicated that the soils had low to moderate salinity, with all soils being non-saline. The lower EC values may be due to high leaching of soils due to their light texture and high permeability, while higher EC values may be due to irrigation with poor quality water and low-lying areas.

The EC of soil was found to be significantly and negatively related to all forms of sulfur fractions, suggesting that under high salinity conditions, SO_4^{2-} may be leached down due to the presence of salts in soluble forms. These findings are consistent with previous research by Vyas et al. (1973), Kaushik and Shukla (1977), Tiwari et al. (1983), Sharma and Gangwar (1997), Gupta et al. (2005), Roy et al. (2002), Tiwari and Pandey (1990), Sadrasania (1992), Sharma and Gangwar (1997), and Chaudhary and Shukla (2002).

OC:

The organic carbon content of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed. The minimum organic carbon value was recorded in Kolayat while the maximum was observed in Shri Dungargarh. The low organic carbon content in sandy soils might be due to factors such as the absence of stable aggregates, severe wind erosion, high microbial decay, high temperature, and good aeration.

A positive and significant relationship was observed between different forms of sulfur and organic carbon, suggesting that the levels of sulfur forms in the soil are dependent on the amount of organic carbon present. These findings are consistent with previous research by Roy et al. (2002), Babu et al. (2006), Mishra and Singh (2008), Srinivas et al. (2011), Singh et al. (1981), Reddy and Mehta (1970), Marsonia et al. (1986), Bhan and Tripathi (1973),

Ruhal and Paliwal (1980), Singh and Sharma (1983), Sharma and Gangwar (1997), Athokpam et al. (2007), and Singh et al. (2009).

Available N:

The available nitrogen content of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed. The minimum available nitrogen value was recorded in Kolayat while the maximum was observed in Shri Dungargarh. The data suggested that the majority of soil samples were low in available nitrogen content, likely due to factors such as hot and dry climate, low organic matter and total nitrogen reserve, and low or no application of organic manures and crop residues.

These findings are consistent with previous research by Revathi et al. (2005), Srinivas et al. (2011), Prasad et al. (1997), and Polara and Kabaria (2006).

A positive and significant relationship was observed between available nitrogen and different forms of sulfur. These findings are consistent with previous research by Desmukh et al. (2004), Singh et al. (2006), and Athokpam et al. (2007).

Available P:

The available phosphorus content of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed. The highest available phosphorus value was recorded in Kolayat while the lowest was observed in Shri Dungargarh. The soils in the study area were found to be medium in available P_2O_5 , with a large range that might be due to variation in soil properties and management practices. The medium to high values of available phosphorus might be attributed to regular application of inorganic phosphatic fertilizers.

A positive significant relationship was observed between different forms of sulfur and available phosphorus, suggesting that all forms of sulfur increased with an increase in the availability of phosphorus in soil. These findings are consistent with previous research by Swarnkar and Verma (1978), Deshmukh et al. (2004), Babu et al. (2006), Jamuna et al. (2008), Srinivas et al. (2011), and Tisdale et al. (2003).

Available K:

The available potassium content of soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - was analyzed. The highest and lowest available

potassium values were both recorded in Shri Dungargarh. The soils in all villages of the four tehsils were found to be medium in available K₂O content. Since potassium fertilizer application is not common in semi-arid and arid regions of Rajasthan, crops are dependent on the native stock of potassium. Depletion of soil potassium due to inadequate fertilization and intensive cultivation in sandy soil has been reported, while medium to high available potassium status could be due to factors such as weathering of potassium-bearing minerals and release of potassium from decomposing organic matter.

A positive correlation was observed between available potash and all forms of sulfur, consistent with previous research by Ramesh and Rao (2005), Mishra and Singh (2008), Surendra et al. (2003), and Revathi et al. (2005).

Conclusion:

In conclusion, an analysis of the soils from four tehsils in Bikaner district - Kolayat, Lunkaransar, Nokha, and Shri Dungargarh - revealed that they were sandy, loamy sand, and sandy loam in texture and alkaline in reaction. The majority of soils were calcareous, with pH values ranging from 8.00 to 9.44 and electrical conductivity varying from 0.07 to 0.20 dSm⁻¹. From the point of view of soil fertility status, the organic carbon content of soil was generally low, ranging from 0.08 to 0.34%, while the available nitrogen, phosphorus, and potassium ranged from 100.10 to 299.84 kg/ha, 11.21 to 27.49 kg/ha, and 113.37 to 571.50 kg/ha, respectively. Thus, the soils of the study area were found to be low in available nitrogen and medium in available P₂O₅ and K₂O content.

Furthermore, the distribution of different forms of sulfur in soil was found to be strongly dependent on soil characteristics and their relationships with each other. Among different soil properties, clay content was found to be the dominant factor explaining variation in sulfur forms in soil, followed by silt and organic carbon content.

References:

Abdallah, M., Dubousset, L., Meuriot, F., Etienne, P., Avice, J. C., & Ourry, A. (2010).

Effect of mineral sulphur availability on nitrogen and sulphur uptake and remobilization during the vegetative growth of *Brassica napus* L. *Journal of Experimental Botany*, 61(10), 2635-2646.

- Abrol, Y. P., & Ahmad, A. (Eds.). (2003). *Sulphur in plants*. Springer Science & Business Media.
- Athokpam, H.S., Singh, R.K.K., Singh, L.N., Singh, N.G., Nandini Chonghtam and Singh, A.K.K. (2007). Distribution of different forms of sulphur in rice growing soils of Nadia District of West Bengal. *Indian J of Agric. Res.* **41**: 205-209.
- Babu, Vijaya Shankar. M., Adinarayana, G.K., Rama subbaiah, Balaguravaiah, D and Yellamanda Reddy, T. 2006. Evaluation of nutrient status of rainfed chickpea growing Vertisols of Anantapur District, Andhra Pradesh. *Indian Journal of Dryland Agriculture Research and Development.* **21**(1): 24-26.
- Balanagoudar, S.R. and Satyanarayana, T. (1990b). Correlation of different forms of sulphur with soil properties and with organic carbon and nitrogen in Vertisols and Alfisols. *J. Indian Soc. Soil Sci.* **38** (4): 641-645.
- Bardsley, C.F. and Lancaster, J. D. (1960). Determination of reserve sulphur and soluble sulphates in soils. *Soil Sci. Soc. Amer. Proc.* **24**: 265-268
- Bascomb, C. L. (1964). Rapid method for the determination of cation- exchange capacity of calcareous and non- calcareous soils. *Journal of the Science of Food and Agriculture*, 15(12), 821-823.
- Bouyoucos, G. J. (1962). Hydrometer method improved for making particle size analyses of soils 1. *Agronomy journal*, 54(5), 464-465.
- Cate Jr, R. B., & Nelson, L. A. (1971). A simple statistical procedure for partitioning soil test correlation data into two classes. *Soil Science Society of America Journal*, 35(4), 658-660.
- Chaudhary, D.R. and Shukla, L.M. (2002). Sulphur status of arid soils of Western Rajasthan. *Annals Agric. Res.* **23**(3): 371-376.
- Chesnin, L. and Yien, C.H. (1950). Turbidimetric determination of available sulphates. *Proc. Soil Sci. Ame.* **15**: 149-151.
- Clarson, D., & Ramaswami, P. P. (1992). Sulphur balance in soils as influenced by sulphur carriers and crops. *Journal of the Indian Society of Soil Science*, 40(2), 313-316.

- Das, K.N., Basumatary, A. and Bikram, Borokotoki. 2011. Interrelationships of forms of sulphur with its availability indices and soil properties in Entisols of Assam. *Journal of the Indian Society of Soil Science*. 59 (2): 134-140.
- Deshmukh, A.H., Ravankar, H.N., Hadole, S.S., Sarap, P.A. and Nagone, A.H. (2004). Status of available sulphur in soils of western Vidarbha. *Ann. of Pl. Physio*. **18** (1): 37-41.
- Deshmukh, A.H., Ravankar, H.N., Hadole, S.S., Sarap, P.A. and Nagone, A.H. (2004). Status of available sulphur in soils of western Vidarbha. *Ann. of Pl. Physio*. **18** (1): 37-41.
- Ensminger, L.E. (1954). Some factors affecting the adsorption of sulphate by Alabama soils. *Soil Sci. Soc. Amer. Proc.* **18** (3): 259-263.
- F.A.O. (1977). Guidelines: For Soil Profile Description 2nd edition. FAO, Rome.
- Freney, J.R. (1967). "Sulphur" Containing Organics in A.D. Mcharen and H.G. Peterson Eds., *Soil Biochemistry*, Dekker, New York. pp. 229-259.
- Gates, C. T. (1974). Nodule and plant development in *Stylosanthes humilis* HBK: symbiotic response to phosphorus and sulphur. *Australian Journal of Botany*, 22(1), 45-55.
- Gupta, G.D., Gupta, R.D., Deepak, Kher., Gupta, J.P and Sumeria, N.M. 2005. Studies on the fertility status in relation to soil properties in Kandi belt of Jammu district. *Journal of Research SKU*. **4**(1): 102-110.
- Gupta, R.C. (2006). Proceedings of TSI-FAI-IFA Symposium-cum- Workshop on Sulphur in Balanced Fertilization. New Delhi. pp. 9-14.
- Hawkesford, M. J., & De Kok, L. J. (2006). Managing sulphur metabolism in plants. *Plant, Cell & Environment*, 29(3), 382-395.
- Jackson, M.L. (1973). *Soil Chemical Analysis*. Prentice Hall of India Pvt Ltd. New Delhi. pp. 327-350.
- Jamuna, P., Nooka Raju, Y and Ramalinga Swamy, K. 2008. Fertility status of soils in Vizianagaram District. *Journal of Research ANGRAU*. 36 (1): 36-41.

- Jat, J.R. and Yadav, B.L. (2006). Different forms of sulphur and their relationship with properties of Entisols of Jaipur District (Rajasthan) under mustard cultivation. *J. Indian Soc. Soil Sci.* **54** (2): 208-212.
- Joshi, N., Gothalwal, R., Singh, M., & Dave, K. (2021). Novel sulphur-oxidizing bacteria consummate sulphur deficiency in oil seed crop. *Archives of Microbiology*, *203*, 1-6.
- Juszczuk, I. M., & Ostaszewska, M. (2011). Respiratory activity, energy and redox status in sulphur-deficient bean plants. *Environmental and Experimental Botany*, *74*, 245-254.
- Kanwar, J.S. (1967). Sulphur deficiency—a key factor in Indian agriculture. *Ferti. News.* **12**:1-4.
- Kara, H., & Ceylan, R. (1988). Removal of sulphur from four Central Anatolian lignites by NaOH. *Fuel*, *67*(2), 170-172.
- Kaushik, R. N., & Shukla, U. C. (1977). A Study on the Development of Salt-Affected Soils in a Typical Area. *Journal of the Indian Society of Soil Science*, *25*(3), 276-283.
- Lal, P., & Singh, K. S. (1974). A comparative study of the effect of qualities of irrigation water on different soils. *J INDIAN SOC SOIL SCI.* *22* (1), P 19-25, 1974.
- Mishra, S.K and Singh, R.P. 2008. Physico-chemical properties and major nutrients (N, P, K and S) status in soils of Arazilline block of Varanasi, Uttar Pradesh. *Crop Research.* *36* (1, 2 &3): 133-136.
- Olsen, S.R., Cole, C.V., Watanable, F.S. and Dean, L.A. (1954). Estimation of available phosphorus in soil by extraction with sodium bicarbonate. USDA, Circular No. 939
- Pagare, S., Bhatia, M., Tripathi, N., Pagare, S., & Bansal, Y. K. (2015). Secondary metabolites of plants and their role: Overview. *Current Trends in Biotechnology and Pharmacy*, *9*(3), 293-304.
- Pal, Y. and Singh, R. (1992). *Farmers and Parliament.* **27** (7):13.
- Polara, J.V and Kabaria, B.D. 2006. Fertility status of irrigated soils of coastal Amreli district of Gujarat. *Journal of Indian Society of Coastal Agricultural Research.* *24* (1): 50-51.

- Prasad, P.R.K., Subbaiah, G.V and Venkateswarulu, B. 1997. Nutrient status of predominant soil types in Krishna, Godavari and Sarada river command areas of Andhra Pradesh. *Journal of Indian Society of Coastal Agricultural Research*. 15 (21): 9-17.
- Ramdevputra, M. V., Akbari, K. N., Sutaria, G. S., Vora, V. D., & Padmani, D. R. (2010). Effect of sulphur application on yield of groundnut and soil fertility under rainfed conditions. *Legume Research-An International Journal*, 33(2), 143-145.
- Ramesh, G., and Rao, P.K. 2005. Nutrient status of groundnut growing soils under rainfed conditions. *Indian Journal of Dryland Agriculture Research and Development*. **20** (1): 127-130.
- Raza, M. A., Feng, L. Y., Manaf, A., Wasaya, A., Ansar, M., Hussain, A., ... & Yang, W. (2018). Sulphur application increases seed yield and oil content in sesame seeds under rainfed conditions. *Field Crops Research*, 218, 51-58.
- Reddy, C.S. and Mehta, B.V. (1970). Fraction of sulphur in some soils of Gujarat to evolve a suitable method for assessing available sulphur status. *Indian J. Agric. Sci.* **40** (1): 5-12.
- Revathi, D., Venkaiah, K., Naidu, M.V.S and Ramavatharam, N. 2005. Nutrient status of tomato (*Lycopersicon esculentum* Mill.) growing soils in Madanapalle division of Chittoor district of Andhra Pradesh. *The Andhra Agricultural Journal*. 52 (1&2): 141-146.
- Richard, L.A. (1954). Diagnosis and improvement of saline and alkali soils. USDA Hand Book No. 60 Oxford and IBH Pub. Co. Culcutta-16. pp.160.
- Roy, H.K., Ajay Kumar and Sarkar, A.K. 2002. Potassium availability in four soil series of Dumka and Palamau regions of Jharkhand. *Journal of research*.14 (2): 153-158.
- Ruhal, D. S. and Paliwal, K. V., 1980, Soil properties affecting the availability of sulphur in soils of Rajasthan. *J. Indian Soc. Soil Sci.*, **28** : 529-531.
- Sadrasania, R. M. (1992). Studies on sulphur availability in soils of North and North-west Gujarat. M.Sc. (Agri.).Thesis. Gujarat Agricultural University, Sardarkrushinagar.
- Scherer, H. W. (2001). Sulphur in crop production. *European Journal of agronomy*, 14(2), 81-111.

- Sehgal, J. (1996). A Textbook of Pedology: concepts and application. Pp: 245-250. Kalyani Publications, New Delhi, India
- Shah, S. H., Islam, S., & Mohammad, F. (2022). Sulphur as a dynamic mineral element for plants: A review. *Journal of Soil Science and Plant Nutrition*, 22(2), 2118-2143.
- Sharma, A. K. and Gangawar, M. S. 1997, Distribution of different forms of sulphur and their relationship with some soil properties in Alfisols, Inceptisols and Mollisols of Moradabad district, Uttar Pradesh. *J. Indian Soc. Soil Sci.*, **45**(3) : 480-485.
- Singh, A.H., Singh, R.K.K., Singh, L.N. Singh, N.G., Chongtham, N. and Singh, A.K.K. (2006). Status and forms of sulphur in acidic soils of Manipur. *J. Indian Soc. Soil Sci.* **54** (3): 351-353
- Singh, R. S. Gupta, M. D. and Yadav, A. D., 1981, Distribution of different forms of sulphur in the soils of Bundhelkhand region. *Indian J. Agric. Res.*, **15** : 201-207.
- Singh, S. P., Ram, J. and Singh, N., 2000, Forms of sulphur in relation to soil characteristics in some soil series of Nagaland. *J. Maharashtra Agric. Univ.*, **25**(1) : 3-8.
- Singh, S.P, Singh, R., Srivastava, P.C. and Singh, P. (2009). Different forms of sulphur in soils of Udham Singh Nagar district, Uttarakhand and their relationship with soil properties. *Agropedology* **19** (1) : 68-74.
- Subbaih, B.V. and Asija, G. L. (1956). A rapid procedure for estimation of available nitrogen in soil. *Curr. Sci.* **25**: 259-263.
- Surendra, S., Prasad, J., Kumar, B and Singh, R.N. 2002. Status of available nutrients in soils of east and west Singhbhum district of Jharkhand. *Journal of Research*. 14 (2): 225-228.
- Swarnakar, K. C. and Verma, M. M., 1978, Distribution of forms of sulphur in soil profile of Bundhelkhand region. *Indian J. Agric. Chem.*, **11** : 49-57.
- Tabatabai, M. A. (1982). Methods of soil analysis, part 2. *Soil enzymes*, 903-947.
- Tisdale, S.L., Nelson, W.L., Beaton, D.J and Havlin, J.L. 2003. *Soil fertility and fertilizers*. Prentice hall of India private limited, New Delhi 180-182.

- Tiwari, K. N., Vandana Nigam and Pathak, A. N., 1983, Effect of sulphur fertilization on yield response and sulphur and nitrogen composition of rice grown in the soils of Kanpur district. *Indian J. Agric. Sci.*, **53**(9) : 812-819.
- Tiwari, R.C. and Pandey, D.K. (1990). Status of different forms and deficiency of sulphur in some soils of Varanasi region of Eastern Uttar Pradesh. *Fert. New.* **35** (3): 35-40.
- Trivedi, S.K., Bansal, K.N. and Singh, V.B. (1998). Important forms of sulphur in profiles of some soil series of northern M.P. *J. Indian Soc. Soil Sci.* **46** (4): 579-583.
- Trivedi, S.K., Bansal, K.N. and Singh, V.B. (1998). Important forms of sulphur in profiles of some soil series of northern M.P. *J. Indian Soc. Soil Sci.* **46** (4): 579-583.
- Vyas, K. K., Marwaha, P. S., & Didal, M. L. (1973). Studies on saline-alkali soils of Jaipur District (Rajasthan). II. Analysis of saturation extract. *Agriculture and agro-industries journal.*
- Walkely, A. and Black, C.A. (1934). An examination of Degtjarey method for determining soil organic matter and a proposed modification of the chromic acid and titration method. *Soil. Sci.* **37**: 29.
- Williams, C.H. and Steinbergs, A. (1959). Soil sulphur fractions as chemical indices of available sulphur in some Australian Soils. *Aust. J. Agric. Res.* **10**: 340-352.

Table 2 Physio-chemical properties of soils of Kolayat tehsil of groundnut growing areas of Bikaner district

Name of Village	Particle size analysis (%)			CaCO ₃ (%)	EC (dSm ⁻¹)	pH	O.C. (%)	Available nutrients (Kg ha ⁻¹)		
	Sand	Silt	Clay					N	P ₂ O ₅	K ₂ O
Amarpura	86.93	6.02	7.05	2.00	0.16	9.11	0.11	190.10	17.18	169.00
Amarpura	77.19	8.06	13.75	0.50	0.09	8.68	0.21	221.10	27.17	214.20
Angnaeu	87.36	4.60	8.04	1.00	0.17	9.19	0.13	199.20	15.56	168.16
Angnaeu	81.99	7.56	10.45	1.50	0.14	8.92	0.15	100.10	20.62	204.72
Akkasar	79.22	7.27	13.51	6.00	0.10	8.75	0.19	217.20	23.03	211.00
Akkasar	88.42	4.53	7.05	2.50	0.18	9.24	0.08	170.85	15.47	166.60
Akkasar	77.24	9.66	12.12	7.00	0.11	8.82	0.18	216.70	20.62	206.90
Hadla	86.22	5.13	8.65	2.00	0.15	9.01	0.16	212.10	18.05	190.58
Hadla	87.78	4.22	8.00	4.00	0.19	9.33	0.16	210.90	13.70	153.80
Akkasar	84.95	8.05	7.00	5.50	0.20	9.44	0.10	181.90	12.45	124.90
Mean	83.73	6.51	9.56	3.20	0.15	9.05	0.15	192.02	18.39	180.99
Min	77.19	4.22	7.00	0.50	0.09	8.68	0.08	100.10	12.45	124.90
Max	88.42	9.66	13.75	7.00	0.20	9.44	0.21	221.10	27.17	214.20
SD	4.44	1.87	2.69	2.28	0.04	0.26	0.04	36.37	4.52	29.31
CV%	5.31	28.66	28.15	71.11	25.79	2.82	28.33	18.94	24.59	16.19

Table 3 Physio-chemical properties of soils of Lunkaransar tehsil of groundnut growing areas of Bikaner district

Name of Village	Particle size analysis (%)			CaCO ₃ (%)	EC (dSm ⁻¹)	pH	O.C. (%)	Available nutrients (Kg ha ⁻¹)		
	Sand	Silt	Clay					N	P ₂ O ₅	K ₂ O
Kasturiya	90.41	2.86	6.73	7.50	0.16	8.85	0.14	192.50	13.70	164.51
Chak 277,600 RD	86.43	4.75	8.82	8.50	0.13	8.17	0.21	218.90	18.90	228.50
Chak 277,600 RD	91.70	2.25	6.05	7.50	0.18	8.98	0.10	170.58	13.70	161.30
Chak 273	88.22	4.92	6.86	5.00	0.15	8.68	0.15	200.70	16.19	180.16
Chak 269	83.65	7.10	9.25	3.00	0.12	8.00	0.24	233.60	27.49	322.60
Bhadera	88.43	3.75	7.82	5.50	0.15	8.60	0.18	210.50	16.81	180.16
Nathvana, 9LKD	86.61	4.93	8.55	3.50	0.14	8.25	0.20	215.71	18.68	194.90
Nathvana 9LKD	89.96	3.84	6.20	5.00	0.18	9.29	0.13	180.82	13.07	155.90
10 LKD	89.46	3.52	7.02	6.00	0.15	8.48	0.17	209.10	17.43	186.30
10 LKD	84.70	6.15	9.15	4.50	0.12	8.15	0.22	221.10	19.61	239.20
Mean	87.96	4.41	7.65	5.60	0.15	8.55	0.17	205.35	17.56	201.35
Min	83.65	2.25	6.05	3.00	0.12	8.00	0.10	170.58	13.07	155.90
Max	91.70	7.10	9.25	8.50	0.18	9.29	0.24	233.60	27.49	322.60
SD	2.58	1.47	1.23	1.79	0.02	0.41	0.04	19.37	4.19	50.62
CV%	2.93	33.43	16.05	32.00	14.53	4.85	25.44	9.43	23.88	25.14

Table 4 Physio-chemical properties of soils of Nokha tehsil of groundnut growing areas of Bikaner district

Name of Village	Particle size analysis (%)			CaCo ₃ (%)	EC (dSm ⁻¹)	pH	O.C.(%)	Available nutrients (Kg ha ⁻¹)		
	Sand	Silt	Clay					N	P ₂ O ₅	K ₂ O
Sindhu	91.8 1	2.86	5.33	8.00	0.19	9.3 4	0.09	166.2 5	14.0 6	142.5 7
Berasar	86.9 5	5.83	7.22	0.50	0.16	9.1 5	0.11	172.5 0	17.1 8	164.5 1
Kakada	87.1 0	5.04	7.86	5.00	0.15	9.1 4	0.15	200.7 0	17.9 4	192.2 0
Kakada	89.3 2	3.25	7.43	10.00	0.16	9.1 4	0.13	199.1 0	16.1 9	160.8 5
Uadsar	87.2 0	4.85	7.95	4.50	0.13	8.9 8	0.15	214.8 0	18.0 5	208.3 9
Uadsar	88.1 2	3.83	8.05	9.50	0.12	8.9 5	0.17	220.3 0	18.8 1	270.5 2
Zhadeli	87.1 4	3.84	9.02	6.00	0.10	8.0 6	0.25	235.2 0	20.9 3	409.4 4
Zhadeli	88.6 0	4.32	7.08	4.50	0.17	9.1 5	0.10	203.8 0	16.1 9	160.5 0
Jasrasar	88.0 5	3.00	8.95	4.50	0.11	8.6 6	0.23	226.8 0	18.9 0	298.4 0
Sadhasar	91.1 3	3.52	5.35	10.00	0.18	9.1 9	0.10	180.8 0	15.5 6	157.1 9
Mean	88.5 4	4.03	7.42	6.25	0.15	8.9 8	0.15	202.0 3	17.3 8	216.4 6
Min	86.9 5	2.86	5.33	0.50	0.10	8.0 6	0.09	166.2 5	14.0 6	142.5 7
Max	91.8 1	5.83	9.02	10.00	0.19	9.3 4	0.25	235.2 0	20.9 3	409.4 4
SD	1.73	0.97	1.27	3.08	0.03	0.3 7	0.06	23.16	1.97	85.26
CV%	1.95	23.9 4	17.1 4	49.35	20.80	4.1 2	37.25	11.46	11.3 3	39.39

Table 5 Physio-chemical properties of soils of Shri Dungargarh tehsil of groundnut growing areas of Bikaner district

Name of Village	Particle size analysis (%)			CaCo ₃ (%)	EC (dSm ⁻¹)	pH	O.C.(%)	Available nutrients (Kg ha ⁻¹)		
	Sand	Silt	Clay					N	P ₂ O ₅	K ₂ O
Benisar	91.42	2.53	6.05	2.50	0.12	9.11	0.21	200.10	13.07	120.64
Lakhasar	89.56	3.30	7.14	6.00	0.08	8.99	0.24	222.70	14.32	149.88
Gajpura	88.42	2.76	8.82	3.50	0.08	8.92	0.29	235.26	15.56	197.41
Gopalsar	88.94	4.21	6.85	5.50	0.10	9.10	0.22	202.27	13.70	131.60
Gopalsar	91.11	3.04	5.85	4.50	0.16	9.14	0.10	172.50	11.21	113.37
Derajsar	88.95	4.12	6.93	8.00	0.09	9.05	0.23	207.35	13.78	142.57
Derajsar	89.02	5.05	5.93	6.00	0.15	9.12	0.13	199.70	13.07	116.98
Seruna	88.18	3.82	8.00	3.50	0.08	8.95	0.26	230.80	14.94	150.27
Seruna	86.90	3.82	9.28	3.50	0.07	8.65	0.34	299.84	20.62	571.50
Seruna	87.61	3.83	8.56	1.50	0.08	8.93	0.29	259.91	15.56	182.78
Mean	89.01	3.65	7.34	4.45	0.10	9.00	0.23	223.04	14.58	187.70
Min	86.90	2.53	5.85	1.50	0.07	8.65	0.10	172.50	11.21	113.37
Max	91.42	5.05	9.28	8.00	0.16	9.14	0.34	299.84	20.62	571.50
SD	1.41	0.75	1.26	1.94	0.03	0.15	0.07	36.16	2.49	137.62
CV%	1.59	20.69	17.11	43.50	31.47	1.64	31.51	16.21	17.09	73.32

Table 6 Range and mean values of physio-chemical properties of groundnut growing areas of Bikaner district

Properties	Min.	Max.	Mean	SD	CV%
Sand	77.19	91.81	87.31	3.43	3.93
Silt	2.25	9.66	4.65	1.71	3.93
Clay	5.33	13.75	7.99	1.90	23.78
CaCo ₃	0.50	10.00	4.88	2.53	51.82
EC	0.07	0.20	0.14	0.04	26.68
pH	8.00	9.44	8.89	0.36	4.10
Organic carbon	0.08	0.34	0.18	0.06	36.00
Available N	100.10	299.84	205.61	30.76	14.96
Available P ₂ O ₅	11.21	27.49	16.98	3.63	21.41
Available K ₂ O	113.37	571.50	196.62	83.83	42.64

Table 7 Coefficient of correlation (r) between different forms of sulphur and physico-chemical properties of groundnut growing areas of Kolayat tehsil of Bikaner district

Kolayat tehsil	Sand	Silt	Clay	CaCO ₃	EC	pH	OC	Available N	Available P ₂ O ₅	Available K ₂ O
Total Sulphur	- 0.808*	0.528*	0.881*	- 0.065	- 0.970*	- 0.989*	0.794*	0.160	0.945*	0.961*
Organic S	- 0.818*	0.522*	0.904*	- 0.056	- 0.978*	- 0.986*	0.795*	0.202	0.961*	0.939*
Available S	- 0.819*	0.498	0.921*	- 0.045	- 0.980*	- 0.987*	0.841*	0.231	0.963*	0.944*
Non Sulphate S	- 0.785*	0.515*	0.853*	- 0.080	- 0.952*	- 0.979*	0.784*	0.137	0.922*	0.963*
Sulphate S	- 0.877*	0.597*	0.941*	0.036	- 0.989*	- 0.985*	0.790*	0.178	0.976*	0.927*
Water SS	- 0.726*	0.455	0.797*	- 0.157	- 0.933*	- 0.960*	0.629*	0.088	0.917*	0.953*
Heat SS	- 0.570*	0.271	0.681*	- 0.308	- 0.858*	- 0.908*	0.569*	0.074	0.861*	0.932*

* Significant at 5 % level of significance

** Significant at 1 % level of significance

Table 8 Coefficient of correlation (r) between different forms of sulphur and physico-chemical properties of groundnut growing areas of Lunkaransar tehsil of Bikaner district

Lunkaransar tehsil	Sand	Silt	Clay	CaCO ₃	EC	pH	OC	Available N	Available P ₂ O ₅	Available K ₂ O
Total Sulphur	- 0.937*	0.880*	0.916*	- 0.521*	- 0.963*	- 0.896*	0.967*	0.974*	0.872*	0.850*
Organic S	- 0.792*	0.766*	0.749*	- 0.494	- 0.812*	- 0.686*	0.850*	0.877*	0.648*	0.613*
Available S	- 0.843*	0.730*	0.900*	- 0.296	- 0.965*	- 0.976*	0.900*	0.937*	0.828*	0.788*
Non Sulphate S	- 0.922*	0.852*	0.918*	- 0.479	- 0.943*	- 0.931*	0.922*	0.908*	0.931*	0.926*
Sulphate S	- 0.844*	0.732*	0.899*	- 0.297	- 0.965*	- 0.976*	0.899*	0.936*	0.828*	0.788*
Water SS	- 0.840*	0.721*	0.905*	- 0.293	- 0.962*	- 0.976*	0.904*	0.938*	0.829*	0.789*
Heat SS	- 0.838*	0.725*	0.890*	- 0.127	- 0.941*	- 0.933*	0.861*	0.852*	0.808*	0.845*

* Significant at 5 % level of significance

** Significant at 1 % level of significance

Table 9 Coefficient of correlation (r) between different forms of sulphur and physico-chemical properties of groundnut growing areas of Nokha tehsil of Bikaner district

Nokha tehsil	Sand	Silt	Clay	CaC ₀₃	EC	pH	OC	Availa ble N	Availa ble P ₂ O ₅	Availa ble K ₂ O
Total Sulphur	- 0.832*	0.206	0.971*	- 0.253	- 0.945*	- 0.758*	0.852*	0.891*	0.947*	0.787*
Organic S	- 0.855*	0.255	0.966*	- 0.308	- 0.934*	- 0.753*	0.836*	0.860*	0.947*	0.777*
Availab le S	- 0.805*	0.175	0.958*	- 0.239	- 0.959*	- 0.764*	0.876*	0.841*	0.940*	0.806*
Non Sulphate S	- 0.811*	0.171	0.970*	- 0.224	- 0.950*	- 0.771*	0.863*	0.914*	0.949*	0.802*
Sulphat e S	- 0.858*	0.316	0.922*	- 0.289	- 0.854*	- 0.594*	0.746*	0.727*	0.858*	0.629*
Water SS	- 0.637*	- 0.061	0.911*	- 0.075	- 0.988*	- 0.857*	0.950*	0.917*	0.941*	0.926*
Heat SS	- 0.421	- 0.045	0.605*	- 0.071	- 0.712*	- 0.953*	0.789*	0.659*	0.785*	0.903*

* Significant at 5 % level of significance

** Significant at 1 % level of significance

Table 10 Coefficient of correlation (r) between different forms of sulphur and physico-chemical properties of groundnut growing areas of Shri Dungargarh tehsil of Bikaner district

Shri Dungargarh tehsil	Sand	Silt	Clay	CaCO ₃	EC	pH	OC	Available N	Available P ₂ O ₅	Available K ₂ O
Total Sulphur	0.771	0.164	0.966*	0.385	0.854*	0.914*	0.949*	0.901*	0.905*	0.744*
Organic S	0.783	0.087	0.933*	0.376	0.956*	0.800*	0.974*	0.847*	0.800*	0.540*
Available S	0.771	0.084	0.919*	0.332	0.848*	0.977*	0.936*	0.959*	0.966*	0.860*
Non Sulphate S	0.662*	0.230	0.883*	0.357	0.659*	0.882*	0.811*	0.819*	0.866*	0.800*
Sulphate S	0.774*	0.082	0.921*	0.336	0.852*	0.976*	0.937*	0.957*	0.964*	0.853*
Water SS	0.778*	0.080	0.924*	0.345	0.832*	0.979*	0.934*	0.962*	0.972*	0.872*
Heat SS	0.819*	0.107	0.985*	0.437	0.845*	0.871*	0.913*	0.886*	0.827*	0.621*

* Significant at 5 % level of significance

** Significant at 1 % level of significance

Table 11 Coefficient of correlation (r) between different forms of sulphur and physico-chemical properties of groundnut growing areas of all four tehsil of Bikaner district

All tehsils of Bikaner district	Sand	Silt	Clay	CaC o ₃	EC	pH	OC	Availa ble N	Availa ble P ₂ O ₅	Availa ble K ₂ O
Total Sulphu r	- 0.387 **	0.1 49	0.534 **	- 0.03 7	- 0.257	- 0.164	0.166	0.353*	0.691* *	0.604* *
Organi c S	- 0.467 **	0.2 05	0.619 **	- 0.09 2	- 0.327 **	- 0.246	0.241	0.441* *	0.734* *	0.663* *
Availa ble S	- 0.299 *	0.0 95	0.417 **	- 0.05 4	- 0.219	0.069	0.097	0.248	0.493* *	0.395* *
Non Sulphat e S	- 0.407 **	0.1 68	0.551 **	- 0.04 7	- 0.277 **	- 0.192	0.178	0.361* *	0.717* *	0.619* *
Sulpha te S	- 0.241	0.0 27	0.393 **	- 0.03 1	- 0.295 *	0.049	0.165	0.298* *	0.458* *	0.368* *
Water SS	- 0.389 **	0.1 20	0.536 **	0.05 9	- 0.202	- 0.572 **	0.227	0.385* *	0.758* *	0.815* *
Heat SS	- 0.471 **	0.0 70	0.715 **	- 0.07 9	- 0.597 **	- 0.719 **	0.595 **	0.714* *	0.787* *	0.860* *

* Significant at 5 % level of significance

** Significant at 1 % level of significance