

**Original Research Article**  
**IMPACT OF FOLIAR NUTRITION APPLIED AT  
DIFFERENT GROWTH STAGES ON NIGER  
(*Guizotia abyssinica* L.)**

**ABSTRACT**

A field experiment was conducted during *Rabi* season of 2021-22 at Niger Research Station, NAU, Vanarasi, Gujarat to study the effect of foliar spray of nutrients applied at different growth stages on *rabi* niger (*Guizotia abyssinica* L.). The experiment was laid out in randomized block design with three replications. The treatments consisted of seven treatments including foliar spray of 19:19:19 and Novel organic liquid (A Patented NAU product) applied at different growth stages on niger crop. The experimental results revealed that foliar application of 1% 19:19:19 applied at flowering and capitula formation stage recorded significantly higher number of branches per plant. The number of seeds per capitulum and seed yield also remained significantly higher with the application of 1% 19:19:19 applied at flowering and capitula formation stage which was found at par with treatment receiving foliar spray of 1% 19:19:19 at flowering stage and foliar spray of 1 % novel organic liquid at flowering stage.

*Keywords: Niger crop, Foliar spray, 19:19:19, Novel organic liquid*

**1. INTRODUCTION**

Oilseeds occupy a key place in agricultural economy of India, constituting an important group of crops next to cereals. Niger (*Guizotia abyssinica* L.) is believed to have originated from Ethiopia where it was domesticated from about 2000 BC, and then spread to India. In India, Niger is grown on an area of 2.61 lakh ha mainly during *kharif*. Andhra Pradesh, Assam, Chhattisgarh, Gujarat, Jharkhand, Karnataka, Madhya Pradesh, Maharashtra, Odisha and West Bengal are the other states where niger is grown. The average yield in India is 3.21 q/ ha. Niger seeds contains edible oil (38 to 43%), protein (18 to 24%), sugar (12%) and minerals essential for human and animal meals. The main aim of foliar fertilization is to eliminate "hidden hunger", which is a very important consideration in high-input production systems and during critical stages of growth when roots are unable to assimilate the required amounts of nutrients, even in fertile soils. One of the best foliar fertilizers 19:19:19 for crop like pulses, vegetables, paddy, for green house cultivation, nurseries, kitchen garden and for all types of field crops. Use of 19:19:19 result in cost saving and at the same time significantly increases the yield. Economic benefit ultimately matters for farmers and novel organic liquid nutrient plays an important role to improve it, (Singhal *et al.*, 2015). Novel organic liquid nutrient is a product of Navsari Agricultural University, Navsari, Gujarat which was patented in the year of 2012. Since foliar nutrient usually penetrate the leaf cuticle or stomata and enter the cell facilitating easy and rapid utilization of nutrients, so foliar nutrition on niger helps in achieving the optimum grain yield of niger.

**Comment [D1]:** The use

**Comment [D2]:** resulted

**Comment [D3]:** Reference pls

**Comment [D4]:** Give in past tense form

## 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The field experiment entitled "effect of foliar nutrition applied at different growth stages on *rabi niger* (*Guizotia abyssinica* L.)" was conducted during *Rabi* season of the year 2021-22 at Niger Research Station, Navsari Agricultural University, Vanarasi. The soil of the experimental field was clayey in texture with medium in available nitrogen, low in available phosphorus and high in available potassium and moderately alkaline in reaction. The experiment was laid out in randomized block design with three replications. The treatments consisted of seven level of foliar nutrition applied at different growth stages of niger crop viz., F<sub>1</sub>: foliar spray of water (Control), F<sub>2</sub>:1% 19:19:19 at flowering stage, F<sub>3</sub>:1% 19:19:19 at capitula formation stage, F<sub>4</sub>:1% 19:19:19 at flowering and capitula formation stage, F<sub>5</sub>:1% Novel organic liquid at flowering stage, F<sub>6</sub>:1% Novel organic liquid at capitula formation stage, F<sub>7</sub>:1% Novel organic liquid at flowering and capitula formation stage. The niger cultivar GNNIG-3 was fertilized with recommended dose 20 kg N and 20 kg P<sub>2</sub>O<sub>5</sub>. At the time of sowing Nitrogen was supplied through urea while phosphorus supplied through single super phosphate.

**Comment [D5]:** Some more details may included like black soil or red soil, whether irrigated or rained. Rainfall received during the period ,

## 3. RESULTS AND DISCUSSION

### 3.1 Growth parameters in niger as influenced by foliar nutrition applied at different growth stages

Data presented in Table 1 indicated that plant height was not affected significantly due to different foliar nutrition treatment at 30, 60 DAS and at harvest. The plant height at 30 DAS and at harvest were recorded higher with foliar application of 1% novel organic liquid at flowering stage (13.40, 79.20 cm). While, at 60 DAS it was recorded higher with foliar spray of 1% 19:19:19 at flowering stage (F<sub>2</sub>) (51.07cm). Although plant height did not differ statistically, but numerical values showed higher plant height in all the foliar nutrition treatments at harvest stage compared to control (F<sub>1</sub>). Foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) showed significantly higher number of branches per plant at harvest stage (6.13) across all the treatments. Treatment F<sub>4</sub> was found at par with treatment 1 % 19:19:19 at flowering stage (F<sub>2</sub>) and treatment 1% novel organic liquid at flowering stage (F<sub>5</sub>).

**Comment [D6]:** If the treatments are non significant, there is no use of giving the individual plant height data.

It might be due to hastening various metabolic process viz., photosynthesis, symbiotic biological N<sub>2</sub> fixation process due to nutrient availability at the initial stage of the crop. Further, phosphorus present in 19:19:19 fertilizer absorbed directly by the plant might have increased cell division and cell development leading to more number of branches. Similar results were obtained by Shwetha *et al.* (2018) in groundnut crop Banasode and Math (2018) and Deshmukh *et al.* (2023) in soybean.

### 3.2 Yield attributes in niger as influenced by foliar nutrition applied at different growth stages

Data furnished in Table 2 indicated that significantly higher number of seeds per capitulum (27.76) was recorded with foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) and was found at par with rest of foliar nutrition treatments except 1% novel organic liquid at capitula formation stage (F<sub>6</sub>) and foliar spray of water (control) (F<sub>1</sub>). The lowest number of seeds per capitula (20.03) were recorded with foliar spray of water

**Comment [D7]:** The study revealed that

(control) (F<sub>1</sub>). significantly higher seed yield (514kg/ha) was recorded with foliar spray of 1% 19:19:19 at flowering and capitula formation stage (F<sub>4</sub>) and was found at par with 1% 19:19:19 at flowering stage (F<sub>2</sub>), 1% Novel organic liquid at flowering and capitula formation stage (F<sub>7</sub>) and 1% Novel organic liquid at flowering stage (F<sub>5</sub>). While, significantly the lower seed yield (379kg/ha) was recorded with foliar spray of water (F<sub>1</sub>). The increase in yield due to foliar nutrition treatment in F<sub>4</sub>, F<sub>2</sub>, F<sub>7</sub> and F<sub>5</sub> as compared to control (F<sub>1</sub>) was 35%, 27%, 23% and 21% respectively.

It might be due to the foliar application of 19:19:19 that increased number of branches, capitula per plant and seeds per capitulum which directly relate to seed yield per plant and production per unit area. Further, foliar spray may have increased the photosynthetic activity and delays the senescence of leaves, which enhance the supply of photosynthate available for grain filling, resulting in more number of capitula per plant and ultimately seed yield. The results are in confirmation with finding of Krishnaveni *et al.* (2021) and Monika *et al.* (2021) in sesame crop; Bodke *et al.* (2022) and Deshmukh *et al.* (2023) in soybean crop. Novel organic liquid nutrients which provide nutrient supply as well as growth promoting hormones. Application of novel at early growth stages thus might have improved growth and development of the crop resembling in number of branches, capitula per plant and seeds per capitula, which is further reflected in increase of seed yield. Similar finding was reported by Singhal *et al.* (2015) and Champaneri *et al.* (2021).

**Comment [D8]:** Change the sentence to past tense form

**Comment [D9]:** Given full treatment details

**Table 1:** Growth parameters in niger as influenced by foliar nutrition applied at different growth stages

**Comment [D10]:** Add Effect of or study of

Treatments	Plant height (cm)			Number of branches
	30 DAS	60 DAS	At harvest	
F <sub>1</sub> : Foliar spray of water (control)	13.02	50.33	73.17	4.63
F <sub>2</sub> : 1% 19:19:19 at flowering stage	12.73	51.07	78.77	5.87
F <sub>3</sub> : 1% 19:19:19 at capitula formation stage	12.38	47.90	76.73	5.27
F <sub>4</sub> : 1% 19:19:19 at flowering and capitula formation stage	13.05	50.48	97.07	6.13
F <sub>5</sub> : 1% Novel organic liquid at flowering stage	13.40	50.89	79.20	5.93
F <sub>6</sub> : 1% Novel organic liquid at capitula formation stage	11.93	46.20	75.10	5.20
F <sub>7</sub> : 1% Novel organic liquid at flowering and capitula formation stage	12.98	50.20	78.93	5.80
SEm±	0.38	1.88	2.91	0.23
CD at 5%	NS	NS	NS	0.73
CV%	5.3	6.6	6.5	7.5

**Table 2:** Yield attributes in niger as influenced by foliar nutrition applied at different growth stages

Treatments	Number of seed per capitulum	Seed yield (kg/ha)
F <sub>1</sub> : Foliar spray of water (control)	20.03	379
F <sub>2</sub> : 1% 19:19:19 at flowering stage	25.87	482
F <sub>3</sub> : 1% 19:19:19 at capitula formation stage	22.72	426
F <sub>4</sub> : 1% 19:19:19 at flowering and capitula formation stage	27.76	514
F <sub>5</sub> : 1% Novel organic liquid at flowering stage	25.13	462
F <sub>6</sub> : 1% Novel organic liquid at capitula formation stage	21.74	418
F <sub>7</sub> : 1% Novel organic liquid at flowering and capitula formation stage	25.65	468
SEm±	1.05	24
CD at 5%	3.25	74
CV%	7.6	9.2

#### 4. CONCLUSION

On the basis of one year field experiment, it can be concluded that spraying of 1% 19:19:19 or 1% novel organic liquid at flowering stage may be done for getting profitable niger production.

**Comment [D11]:** Give the economics to report as profitable or non profitable

#### REFERENCES

1. Champaneri DD, Patel NK, Desai CS, Desai DH. Efficacy of novel organic liquid nutrient and novel plus organic liquid nutrient on quantitative traits of Indian bean [*Lablab purpureus* (L.) Sweet]. *International Journal of Plant and Soil Science*. 2021; 33 (17): 105-115.
2. Bodke VS, Hanwate GR, Gorde NB, Jaybhaye BB, Pidurkar PK. Studies on influence of foliar application of water-soluble fertilizers on growth and yield of soybean (*Glycine max* L.). *The Pharma Innovation Journal*. 2022; 11 (12): 1682-1686.
3. Monika P, Sumathi V, Reddy PM, Naidu M, Sagar GK. Impact of fertilizers and foliar sprays on productivity of summer sesame (*Sesamum indicum* L.) *Andhra Pradesh Journal of Agricultural Sciences*. 2021; 7 (3): 173-177.
4. Krishnaveni SA, Supriya C, Sridhar SM. Impact of foliar nutrition on the yield and economics of green gram (*Vigna radiata*). *International Journal of Chemical Studies*. 2021; 9 (2): 11-13.
5. Deshmukh SP, Rathva RS, Bambhaneeya SM, Jagtap PK, Mistry PM. Effect of foliar nutrition applied at different growth stages soybean (*Glycine Max.*) under rainfed condition. *Journal of Oilseed Research*. 2023; 39 (special issue): 467-469.

6. Banasode C, Math KK. Effect of foliar fertilization of water-soluble fertilizers on growth and economics of soybean in a vertisol. *Journal of Pharmacognosy and Phytochemistry*. 2018; 7 (2): 2391-2393.
7. Shwetha BN, Anupama C, Sowmya TM, Yaligar R. Effect of foliar nutrition on productivity of groundnut crop. *Journal of Pharmacognosy and Phytochemistry*. 2018;7 (1S): 2357-2360.
8. Singhal VK, Patel GG, Patel DH, Kumar U, Saini LK. Effect of foliar application of water soluble fertilizers on growth, yield and economics of vegetable cowpea production. *Ecosan*. 2015; 7: 79-83.

UNDER PEER REVIEW