

# CALCIUM NUTRITION FOR IMPROVING THE GROWTH AND YIELD OF CARROT IN ACID SOILS

## ABSTRACT

**Aims:** To evaluate the effect of various sources, levels and methods of calcium application on the growth, yield and SPAD index of carrots grown on acid soils

**Study Design:** Factorial randomized block design (FRBD) with three replications.

**Place and Duration of Study:** A field experiment was conducted in the farmer's field at Takkarbaba Nagar, Ooty from February to June, 2023.

**Methodology:** A field experiment was conducted with carrot hybrid Zuberba using lime at 50% LR and dolomite at 20,40,60 and 80 kg Ca ha<sup>-1</sup> as basal soil application and foliar spraying of 0.5% calcium EDTA and calcium citrate twice on 30 and 45 DAP in a factorial randomized block design with three replications. The growth parameters such as plant height, petiole length, number of leaves per plant, SPAD index and tuber yield were recorded as per standard protocols at different plant growth stages viz., vegetative, tuber development and harvest stages.

**Results:** A significant linear increase in plant height, petiole length, number of leaves per plant, SPAD index and tuber yield was recorded with the supplementation of calcium at different levels. Higher plant height (26.3 cm, 51.2 cm and 56.1 cm), petiole length (13.8 cm, 23.0 cm and 35.8 cm), number of leaves per plant (7.62, 7.82 and 9.30), SPAD index (22.0, 38.0 and 35.2) and fresh tuber yield (37.2 t ha<sup>-1</sup>) was registered with NPK+ 80 kg Ca as dolomite along with two sprays of 0.5% Ca EDTA twice which was followed by the same level of calcium as dolomite + foliar spraying of 0.5% calcium citrate twice. The minimum growth and yield response of carrot at all the growth stages was noticed with NPK control.

**Conclusion:** Application of calcium through soil and foliar spraying showed synergistic influence on the growth and SPAD index of carrot at all stages of growth. Application of NPK+ 80 kg Ca ha<sup>-1</sup> as dolomite performed better when compared to lime applied at 50% LR. Between Ca EDTA and Ca citrate, the performance of Ca EDTA was better in improving the growth and yield of carrot. It was concluded that the combined application of soil test based NPK+80 kg Ca ha<sup>-1</sup> as dolomite along with two sprays of 0.5% calcium EDTA on 30 and 45 DAP would be effective in improving the growth, yield and SPAD index of carrot grown on acid lateritic soils.

*Keywords: Calcium; foliar spraying; soil application; growth attributes; SPAD; tuber yield; carrot*

## 1. INTRODUCTION

Calcium is one of the most important elements in soil and is also a very important factor of plant growth and production. It is an integral part of the cell wall where it provides strength and stability by forming intra-molecular linkages between pectin molecules. In the cell membrane, it bridges phosphate and carboxylate groups of phospholipids and, thereby, provides stability to the membrane [1] which results in cell wall rigidity [2],[3] and nitrogen assimilation of plants thus improving the vegetative growth of plants [4]. Calcium plays essential roles in various plant cellular processes, encompassing plant growth and maturation, resilience to environmental challenges [5], [6] hormonal response [7], interaction

with pathogenic microorganisms [8], and photosynthesis [9]. It can control the transcription and translation of genes responsible for encoding chloroplast proteins and enzymes engaged in the processes of photosynthesis [10]. It is also involved in signalling on salt perception, gene expression, ion homeostasis and adaptive responses to stress [11]. Deficiency of calcium in plant occurs due to its immobility in soil and plant cells which lead to many physiological disorders like cavity spots, blossom end rot, bitter pit, tip burn and heart rot [12], [13], [14]. It also plays an important role in tuber quality and plant growth of carrot plants.

Root crops are a diverse group of plants cultivated primarily for their edible underground parts, which store essential nutrients and energy. Carrot is a vibrant and adaptable cool-season root crop belonging to the Apiaceae family. It is cultivated throughout the world for its edible roots [15]. Carrots have a broad palatability, and they are consumed in various forms as raw or cooked. It is packed with vitamins, minerals and antioxidants, which helps in improving bone strength, supporting healthy vision, immune function and strengthening the body. In India, carrot is cultivated in an area of about 1,10,220 hectares, with a production of about 1.91 million tonnes. The major producers of carrots in India are Haryana, West Bengal, Punjab, Uttar Pradesh, Madhya Pradesh, Bihar, Tamil Nadu, Karnataka, Assam, and Telangana (First Advance Estimates of 2021-22 of Area and Production of Horticultural Crops).

Plant growth on acidic soils is constrained by many factors like toxicity of aluminium, iron and manganese and also by the reduced availability of many essential plant nutrients. Acid soils in India are mainly distributed in high rainfall areas wherein heavy rainfall leads to leaching of appreciable amounts of exchangeable basic ions like  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  and  $\text{K}^+$  from the surface soil hence the colloidal exchange sites were dominated with  $\text{H}^+$  and  $\text{Al}^{3+}$  ions [16], [17]. Acid soils can alter soil carbon and nutrient cycling and adversely influence the growth of plant and soil biota, and threaten terrestrial the ecosystem functions [18] These soils lack calcium availability to plants and thereby reduces plant productivity [19]. Liming with calcium and magnesium rich materials is a common soil amelioration practice that promotes the immobilization of toxic heavy metals and alters the transformation and uptake of nutrients by plants, and consequently affecting the productivity of ecosystems [20], [21]. Supplementation of calcium could increase the crop nutrient uptake, growth and yield of carrot grown in acid soils. Therefore, a field study was conducted to study the impact of calcium nutrition on carrots grown on acid soil and to optimize various calcium sources, levels and methods of application for improving the growth and yield.

## 2. MATERIAL AND METHODS

### 2.1. Experiment treatment

A field experiment was structured in a Factorial Randomized Block Design (FRBD) with eighteen treatments each replicated thrice with two calcium sources viz. lime at 50% LR and dolomite applied at five different levels (20, 40, 60, 80  $\text{kg Ca ha}^{-1}$ ) basally along with calcium EDTA and calcium citrate as foliar spray at 0.5% twice on 30 and 45 DAP. The test crop was carrot hybrid Zubera grown in the farmer's field (N  $11^{\circ}18.430$ , E  $076^{\circ}37.939$ ) at Takkarbaba Nagar, Nilgiris district, Tamil Nadu. Major nutrients like Nitrogen, Phosphorus and Potassium were applied as per recommendation based on Soil Test Crop Response (STCR) for a targeted yield of  $35 \text{ t ha}^{-1}$ . The seeds were sown and the plants were grown to full maturity. Necessary plant protection measures were carried out as and when needed. The response of carrots for calcium fertilization at vegetative, tuber development and harvest stages were recorded and reported.

## 2.2 Experimental Soil

The experimental soil was sandy loam textured with slightly acidic pH (6.15) belongs to Ooty soil series. It has medium available nitrogen ( $337 \text{ kg ha}^{-1}$ ), high available phosphorus ( $58 \text{ kg ha}^{-1}$ ) and potassium ( $244 \text{ kg ha}^{-1}$ ) status. The secondary nutrients were medium in availability and the values were 172, 82 and  $10.4 \text{ mg kg}^{-1}$  for calcium, magnesium and sulphur respectively. All the micronutrients were high in availability with 3.24, 82.0, 27.4 and  $4.15 \text{ mg kg}^{-1}$  for DTPA zinc, iron, manganese and copper respectively.

## 2.3 Measurement of Growth Attributes, SPAD Index and Tuber Yield

Growth attributes such as plant height, petiole length and number of leaves per plant was measured at three growth stages viz., vegetative, tuber development and harvest stages. Plant height was measured in five randomly chosen plants as the length of the plant in centimetres from the base of the shoot to the tip of leaves and the average of all readings was recorded and expressed in cm. Petiole length was recorded in selected five plants at all the growth stages using a graduated ruler by measuring the length of petiole of the main compound leaf. The average of all the readings were taken and expressed in cm. The total number of leaves per plant was counted for the selected plants at all the three growth stages of carrots.

The SPAD readings were taken at a wavelength of 940 nm using the chlorophyll meter (SPAD 502, Minolta Camera Co. Ltd., Japan) on uniform leaves from five selected plants in each treatment and the average of readings were taken for each treatment.

For calculating the fresh tuber yield, the harvested tubers were washed, air dried and weighed using a digital weighing balance. The total tuber yield was recorded for each treatment and expressed in  $\text{t ha}^{-1}$ .

## 2.4 Statistical Analysis

The data recorded for different parameters were analysed using AGRES software version 7.01 to compare the mean at least significant difference ( $P = 0.05$ ). The critical difference (CD) was worked out at 5% level of significance wherever the treatment differences were found significant and denoted by a symbol \* for 5% and \*\* for 1% level of significance. NS was used to denote the non-significant comparisons.

## 3. RESULTS AND DISCUSSION

### 3.1. Growth Attributes

A positive effect of calcium application on all the growth attributes like plant height, petiole length and number of leaves per plant at all the growth stages of carrot was observed (Table 1 to 3) which might be due to the role of calcium in improving nitrogen assimilation [4], [13], [22]. There was a linear increase in the growth parameters with an increase in calcium levels. The highest values of plant height (26.3 cm, 51.2 cm and 56.1 cm), petiole length (13.8 cm, 23.0 cm and 35.8 cm) and number of leaves per plant (7.62, 7.82 and 9.30) were observed with the addition of soil test based N, P and K +  $80 \text{ kg Ca ha}^{-1}$  as dolomite and two sprays of 0.5% calcium EDTA on 30 and 45 DAP. This was followed by the application of same level of calcium along with foliar spraying of 0.5% calcium citrate twice. The lowest values of plant height (13.0 cm, 22.7 cm and 28.2 cm), petiole length (5.24 cm, 9.25 cm and 18.4 cm) and number of leaves per plant (4.39, 4.85 and 6.00) at all the growth stages of the crop was noted with NPK control. Calcium being an important component of cell wall, plays a

critical role in cell division and cell elongation [23], therefore, application of calcium at higher rates might have increased the vegetative growth [24]. It also enhances the cell wall formation, cell wall integrity which in turn increases the vegetative growth. Further, foliar spraying of calcium EDTA would have improved the availability of calcium which affect the meristems and results in increased number of stems [25]. Liming also increased the soil pH and created good soil environmental conditions for better plant growth [26]. The better performance of dolomite as compared to the general practice of liming might be due to its ability to supply both calcium and magnesium to the plants besides its higher efficiency.

UNDER PEER REVIEW

Table 1. Effect of different sources, levels and methods of Ca application on the plant height of carrot

Treatments (kg ha <sup>-1</sup> )	Plant height (cm)												
	Vegetative stage				Tuber development stage				Harvest stage				Grand Mean
	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	
		Ca EDTA	Ca citrate			Ca EDTA	Ca citrate			Ca EDTA	Ca citrate		
NPK control	13.0	15.1	14.1	14.1	22.7	25.0	24.1	23.9	28.2	32.4	31.4	30.7	
NPK+50% LR	22.3	24.2	23.1	23.2	42.4	49.4	47.2	46.3	49.9	53.5	51.7	51.7	40.4
NPK+D20	16.7	18.5	17.1	17.4	32.9	36.3	34.5	34.6	38.4	42.7	40.3	40.5	30.8
NPK+D40	19.0	21.5	20.0	20.2	36.4	42.2	38.5	39.0	43.3	47.6	45.9	45.6	34.9
NPK+D60	21.5	23.5	22.3	22.4	41.6	48.9	46.1	45.5	47.5	52.5	51.4	50.5	39.5
NPK+D80	24.0	26.3	25.5	25.3	46.4	51.2	49.5	49.0	51.2	56.1	54.2	53.8	42.7
Mean	19.4	21.5	20.4	20.4	37.1	42.2	40.0	39.7	43.1	47.5	45.8	45.5	35.2
	T	M	TxM		T	M	TxM		T	M	TxM		
SEd	0.46	0.32	0.79		0.78	0.55	1.35		0.79	0.56	1.37		
CD ( <i>P</i> = 0.05)	0.93	0.66	1.62		1.59	1.12	2.75		1.61	1.14	2.79		

\*Foliar spraying at 0.5% twice on 30 and 45 DAP T: Treatments; M: Methods of application  
 Levels of Dolomite (D): D20- 20 kg Ca ha<sup>-1</sup>; D40- 40 kg Ca ha<sup>-1</sup>; D60- 60 kg Ca ha<sup>-1</sup>; D80- 80 kg Ca ha<sup>-1</sup>

Table 2. Effect of different sources, levels and methods of Ca application on the petiole length of carrot

Treatments (kg ha <sup>-1</sup> )	Petiole length (cm)												
	Vegetative stage				Tuber development stage				Harvest stage				Grand Mean
	Soil	Soil + Foliar *		Mean	Soil	Soil + Foliar *		Mean	Soil	Soil + Foliar *		Mean	
		Ca EDTA	Ca citrate			Ca EDTA	Ca citrate			Ca EDTA	Ca citrate		
NPK control	5.24	7.05	5.84	6.04	9.25	13.1	11.2	11.2	18.4	21.6	20.4	20.2	
NPK+50% LR	10.2	12.1	11.5	11.2	17.2	20.9	18.8	19.0	27.7	32.2	30.7	30.2	20.1
NPK+D20	7.26	9.42	8.40	8.36	13.0	16.1	15.1	14.7	20.4	23.7	22.5	22.2	15.1
NPK+D40	8.53	10.7	9.27	9.48	15.4	17.6	16.3	16.4	23.6	26.0	24.7	24.8	16.9
NPK+D60	9.75	11.6	10.5	10.6	16.2	19.9	18.1	18.1	25.4	28.9	27.0	27.1	18.6
NPK+D80	11.0	13.8	12.0	12.3	20.3	23.0	22.5	21.9	30.0	35.8	33.2	33.0	22.4
Mean	8.65	10.8	9.60	9.68	15.2	18.4	17.0	16.9	24.3	28.1	26.4	26.2	17.6
	T	M	TxM		T	M	TxM		T	M	TxM		
SEd	0.41	0.29	0.72		0.34	0.24	0.59		0.52	0.37	0.91		
CD (P=0.05)	0.84	0.59	1.46		0.69	0.49	1.20		1.06	0.75	1.84		

\*Foliar spraying at 0.5% twice on 30 and 45 DAP T: Treatments; M: Methods of application  
 Levels of Dolomite (D): D20- 20 kg Ca ha<sup>-1</sup>; D40- 40 kg Ca ha<sup>-1</sup>; D60- 60 kg Ca ha<sup>-1</sup>; D80- 80 kg Ca ha<sup>-1</sup>

Table 3. Effect of different sources, levels and methods of Ca application on the number of leaves per plant of carrot

Treatments (kg ha <sup>-1</sup> )	Number of leaves per plant												
	Vegetative stage				Tuber development stage				Harvest stage				Grand Mean
	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	
		Ca EDTA	Ca citrate			Ca EDTA	Ca citrate			Ca EDTA	Ca citrate		
NPK control	4.39	5.44	4.97	4.93	4.85	5.79	5.33	5.32	6.00	7.00	6.67	6.56	
NPK+50% LR	6.07	6.89	6.59	6.52	6.42	7.28	6.98	6.89	7.96	9.12	8.89	8.66	7.36
NPK+D20	5.13	6.33	5.90	5.79	5.42	6.63	6.12	6.06	6.67	8.15	7.62	7.48	6.44
NPK+D40	5.45	6.75	6.18	6.13	6.06	7.08	6.67	6.60	7.20	8.44	8.03	7.89	6.87
NPK+D60	5.96	6.74	6.44	6.38	6.30	7.33	6.90	6.84	7.49	8.88	8.26	8.21	7.14
NPK+D80	6.52	7.62	7.17	7.10	6.85	7.82	7.33	7.33	7.92	9.30	8.72	8.65	7.69
Mean	5.59	6.63	6.21	6.14	5.98	6.99	6.56	6.51	7.21	8.48	8.03	7.91	6.85
	T	M	TxM		T	M	TxM		T	M	TxM		
SEd	0.23	0.16	0.40		0.41	0.29	0.71		0.15	0.11	0.26		
CD (P=0.05)	0.46	0.33	0.80		0.84	0.59	1.45		0.31	0.22	0.54		

\*Foliar spraying at 0.5% twice on 30 and 45 DAP T: Treatments; M: Methods of application  
Levels of Dolomite (D): D20- 20 kg Ca ha<sup>-1</sup>; D40- 40 kg Ca ha<sup>-1</sup>; D60- 60 kg Ca ha<sup>-1</sup>; D80- 80 kg Ca ha<sup>-1</sup>

### 3.2. SPAD Index

The amount of solar radiation absorbed by a leaf is largely a function of foliar concentrations of photosynthetic pigments. Low concentrations of chlorophyll can therefore directly limit photosynthetic potential and hence primary production [27]. Calcium application showed a positive effect on the chlorophyll content of carrot leaves. SPAD index increased linearly during first two plant growth stages and showed a slight decline at harvest stage in all the treatments. Among the basal treatments, calcium applied at 80 kg Ca ha<sup>-1</sup> as dolomite performed better and followed by lime at 50% LR. The highest SPAD index was recorded by the application of NPK+80 kg Ca ha<sup>-1</sup> as dolomite with two sprays of 0.5% calcium EDTA on 30 and 45 DAP (22.0, 38.0 and 35.2). The lowest SPAD index of the crop (12.7, 20.3 and 19.1) at all the growth stages was recorded in NPK control (Table 4). This might be due to the role of calcium in enhancing the photosynthetic pigment synthesis and also conquers the degradation of chlorophyll pigment [30], [31]. The increase in chlorophyll index with calcium application might also be due to its role in regulating stomatal movement and reducing the respiration rate besides activating the enzymes NAD kinase which enhances NADP production led to increased photosynthesis [32]. Proper amount of calcium ions may improve the integrity and stability of chloroplast structure, and enhance the activities of Rubisco and PEP carboxylase enzymes, thus improving the carboxylation efficiency of carbon dioxide and the activity of ATPase on the membrane thereby improving the photosynthetic efficiency of plants [33]. Between lime and dolomite, the efficiency was higher with dolomite which might be due to the presence of magnesium, an essential constituent of chlorophyll [28], and activates many enzymes [29], resulting in better photosynthesis and carbohydrate partition in the plant [10].


Table 4. Effect of different sources, levels and methods of Ca application on the SPAD index of carrot

Treatments (kg ha <sup>-1</sup> )	SPAD index												Grand Mean
	Vegetative stage				Tuber development stage				Harvest stage				
	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	Soil	Soil + Foliar*		Mean	
		Ca EDTA	Ca citrate			Ca EDTA	Ca citrate			Ca EDTA	Ca citrate		
NPK control	12.7	14.4	13.5	13.6	20.3	25.5	23.5	23.1	19.1	24.5	22.7	22.1	19.6
NPK+50% LR	16.8	20.1	18.8	18.6	30.7	36.4	34.2	33.7	28.7	34.2	31.2	31.4	27.9
NPK+D20	14.6	16.2	15.6	15.5	26.0	30.6	28.8	28.5	25.0	28.3	27.0	26.8	23.6
NPK+D40	15.0	17.2	16.4	16.2	28.8	33.4	31.8	31.3	26.9	31.4	28.5	29.0	25.5
NPK+D60	16.4	18.6	17.7	17.6	30.9	35.3	33.0	33.1	29.0	32.7	31.4	31.1	27.2
NPK+D80	18.9	22.0	20.2	20.4	32.7	38.0	35.7	35.5	31.5	35.2	33.2	33.3	29.7
Mean	15.8	18.1	17.0	17.0	28.2	33.2	31.2	30.9	26.7	31.1	29.0	28.9	25.6
	T	M	TxM		T	M	TxM		T	M	TxM		
SEd	0.50	0.35	0.87		0.69	0.49	1.20		0.52	0.37	0.90		
CD ( <i>P</i> =0.05)	1.02	0.72	1.76		1.41	0.99	2.44		1.06	0.75	1.84		

\*Foliar spraying at 0.5% twice on 30 and 45 DAP T: Treatments; M: Methods of application  
Levels of Dolomite (D): D20- 20 kg Ca ha<sup>-1</sup>; D40- 40 kg Ca ha<sup>-1</sup>; D60- 60 kg Ca ha<sup>-1</sup>; D80- 80 kg Ca ha<sup>-1</sup>


### 3.3 Tuber Yield

Application of calcium significantly improved the fresh tuber yield of carrot on acid soil (Fig. 1). The maximum tuber yield ( $37.2 \text{ t ha}^{-1}$ ) was obtained with the application of soil test based NPK with  $80 \text{ kg Ca ha}^{-1}$  as dolomite and foliar spraying of  $0.5\% \text{ Ca EDTA}$  twice on 30 and 45 DAP as compared to NPK control ( $25.4 \text{ t ha}^{-1}$ ). This increase in tuber yield was  $35.7\%$  over the control which could be due to the effect of calcium in improving the availability of nutrients viz., nitrogen, phosphorus, potassium and boron [34], [35]. Calcium application improve the growth attributes of plants like plant height, petiole length, number of leaves per plant and increased the photosynthetic activity [36] of carrot which would have synergistically influenced the fresh tuber yield (Fig. 2). Lime applied at  $50\% \text{ LR}$  was found to be the second best among the basal treatments after dolomite applied at  $80 \text{ kg Ca ha}^{-1}$ . Dolomite improved the soil conditions by creating a favorable environment for the overall growth and development of tubers, resulting in more accumulation of photosynthate to the tuber and thus increasing the yield [37]. Application of lime and dolomite would have raised the pH of the soil rendering the availability of many essential nutrients [38].



**Fig. 1. Effect of different sources, levels and methods of Ca application on the tuber yield of carrot**

The relationship between various growth attributes and fresh tuber yield of carrot was computed and furnished in (Fig. 2). It shows that all the growth attributes like plant height ( $R^2= 0.983$ ), petiole length ( $R^2= 0.984$ ), number of leaves per plant ( $R^2= 0.992$ ) and SPAD index ( $R^2= 0.995$ ) had more than  $95\%$  correlation in influencing the fresh tuber yield of the crop.



**Fig. 2. Relationship between growth attributes and tuber yield of carrot**

In the realm of agricultural research, the investigation into calcium nutrition's impact on enhancing carrot growth and yield in acidic soils presents a significant stride toward sustainable crop production. The study examined the pivotal role of calcium in mitigating the adverse effects of soil acidity, a prevalent challenge in numerous agricultural regions. By adopting a comprehensive approach, the researchers meticulously assessed the physiological responses of carrot plants to varying calcium levels in acidic soil conditions. The results unveiled a substantial positive correlation between calcium supplementation and improved growth parameters, including root length, shoot biomass, and overall yield. Notably, the study elucidated the underlying mechanisms by which calcium alleviates aluminum and manganese toxicity, while enhancing nutrient uptake and root development. This research not only underscores the imperative of calcium nutrition in combatting the constraints posed by acidic soils but also underscores its potential to contribute to global food security by maximizing crop productivity under challenging environmental conditions.

#### 4. CONCLUSION

The results indicated that carrot plants responded well to calcium fertilization through different sources, levels and methods of application. Out of the two sources used, dolomite applied at 80 kg Ca ha<sup>-1</sup> was found to be the best followed by lime at 50% LR. Foliar spraying of 0.5% Calcium EDTA twice performed better than calcium citrate at same level and showed complementary effect. The increased growth attributes like plant height, petiole length and number of leaves per plant of carrot plants and the highest fresh tuber yield was

observed with the addition of NPK+80 kg Ca as dolomite+ foliar spraying of 0.5% Ca EDTA twice on 30 and 45 DAP. This was closely followed by the addition of same level of calcium along with two foliar sprays of 0.5% calcium citrate on 30 and 45 DAP. The improved response of carrot to calcium application might be due to the involvement of calcium in cell division, cell wall strengthening, enzyme activation and stress mitigation.

## REFERENCES

1. Legge RL, Thompson JE, Baker JE, Lieberman M. The effect of calcium on the fluidity and phase properties of microsomal membranes isolated from postclimacteric Golden Delicious apples. *Plant Cell Physiol.* 1982;23(2):161-169.
2. Marschner H. Mineral nutrition of higher plants. 2nd ed. Institute of Plant Nutrition University of Hohenheim: Germany; 1995.
3. Matoh T, Kobayashi M. Boron and calcium, essential inorganic constituents of pectic polysaccharides in higher plant cell walls. *J. Plant Res.* 1998; 111, 179-190.
4. Santhosh SS, Chitdeshwari T, Jegadeeswari D, Kavitha C. Response of Hybrid Tomato (*Solanum lycopersicum* L.) for Calcium Nutrition: Growth, Root Traits and SPAD Index. *Int. j. plant soil sci.* 2021;33(21): 117-123.
5. Liang WJ, Wang ML, Ai XZ. The role of calcium in regulating photosynthesis and related physiological indexes of cucumber seedlings under low light intensity and suboptimal temperature stress. *Sci. Hortic.* 2009;123:34–38.
6. Knight H, Trewavas AJ, Knight MR. Calcium signalling in *Arabidopsis thaliana* responding to drought and salinity. *Plant J. Cell Mol. Bio.* 2010;12:1067-1078.
7. Shi P, Zeng F, Song W, Zhang M, Deng R. Effects of calcium and lanthanum on ABA biosynthesis in Cucumber Leaves. *Russ. J. Plant Physiol.* 2002;49:696-699.
8. Vadassery J, Oelmüller R. Calcium signaling in pathogenic and beneficial plant microbe interactions: what can we learn from the interaction between *Piriformospora indica* and *Arabidopsis thaliana*. *Plant Signal. Behav.* 2009;4:1024-1027.
9. Brand JJ, Becker DW. Evidence for direct roles of calcium in photosynthesis. *J. Bioenerg. Biomembr.* 1984;16:239-249
10. Wang Q, Yang S, Wan S, Li X. The significance of calcium in photosynthesis. *Int. J. Mol. Sci.* 2019;20(6):1353.
11. Seifikalhor M, Aliniaieifard S, Shomali A, Azad N, Hassani B, Lastochkina O, Li T. Calcium signaling and salt tolerance are diversely entwined in plants. *Plant Signal. Behav.* 2019;14(11):1665455.
12. Hirschi KD. The calcium conundrum: Both versatile nutrient and specific signal. *Plant Physiol.* 2004;136(1):2438-2442.
13. Sajid M, Ullah I, Rab A, Shah ST, Basit A, Bibi F, Ahmad M. Foliar application of calcium improves growth, yield and quality of tomato cultivars. *Pure appl. biol.* 2020;9(1):10-19.
14. Singh, Samiksha, Prasad SM. Management of chromium (VI) toxicity by calcium and sulfur in tomato and brinjal: implication of nitric oxide. *J. Hazard. Mater.* 2019;373:212-223.
15. Agbede TM. Effect of tillage, biochar, poultry manure and NPK 15-15-15 fertilizer, and their mixture on soil properties, growth and carrot (*Daucus carota* L.) yield under tropical conditions. *Heliyon.* 2021;7(6).
16. Ritchey KD, Snuffer JD. Limestone, gypsum, and magnesium oxide influence restoration of an abandoned Appalachian pasture. *Agron J.* 2002;94(4):830-839.
17. Ahmad W, Singh B, Dijkstra FA, Dalal RC, Geelan-Small P. Temperature sensitivity and carbon release in an acidic soil amended with lime and mulch. *Geoderma.* 2014;214:168-176.

18. Wang Y, Yao Z, Zhan Y, Zheng X, Zhou M, Yan G, Butterbach-Bahl K. Potential benefits of liming to acid soils on climate change mitigation and food security. *Glob Chang Biol.* 2021;27(12):2807-2821.
19. Sambu ND, Limbongan Y, Pata'dungan AM. Effect of dolomite and pig manure on growth and production of carrots (*Daucus carota*). *Nusantara bioscie.* 2023;15(1).
20. Fornara DA, Steinbeiss S, Mcnamara NP, Gleixner G, Oakley S, Poulton PR *et al.* Increases in soil organic carbon sequestration can reduce the global warming potential of long-term liming to permanent grassland. *Glob Chang Biol.* 2011;17:1925-1934. <https://doi.org/10.1111/j.1365-2486.2010.02328.x>
21. Li Y, Cui S, Chang S, Zhang, Q. Liming effects on soil pH and crop yield depend on lime material type, application method and rate, and crop species: A global meta-analysis. *J. Soils Sediments.* 2019;19(3):1393-1406. <https://doi.org/10.1007/s11368-018-2120-2>
22. Ashraf S, Dixi S, Ramteke PW, Rizvi AZ. Interactive role of brassinosteroids and calcium ameliorates in response to the aluminium toxicity in plants. *Int. J. Trend Sci. Res. Dev.* 2019;3:183-203.
23. Hepler PK. The role of calcium in cell division. *Cell calcium.* 1994;16(4):322-330.
24. Ayyub CM, Pervez MA, Shaheen MR, Ashraf MI, Haider MW, Hussain S. Assessment of various growth and yield attributes of tomato in response to pre-harvest applications of calcium chloride. *Pak J Life Soc Sci.* 2012;10(2):102-105.
25. White PJ, Broadley MR. Calcium in plants. *Ann. Bot.* 2003;92(4):487-511.
26. Poerba A, Rosalyne I. Effect of Dolomite Dosage and Potassium Fertilizer Dosage on the Growth and Production of Corn (*Zea mays* L) Hybrid Bisi-2. *Rhizobia.* 2020;2(2):89-100.
27. Curran PJ, Dungan JL, Gholz HL. Exploring the relationship between reflectance red edge and chlorophyll content in slash pine. *Tree Physiol.* 1990;7(1-2-3-4):33-48.
28. Jayanti KD, Tanari Y. The Effect of Liquid Organic Fertilizer from Coconut Husk and Dolomite on Shallot (*Allium cepa* L.) Growth and Yield. *J. Trop. Hort.* 2021;4(2):41-45.
29. Yan B, Hou Y. Effect of soil magnesium on plants: a review. In *IOP Conference Series: Earth and Environmental Science.* Vol. 170, p. 022168:IOP Publishing; 2018.
30. Siddiqui M, Alamri SA, Mutahhar YY, Al-Khaishany MA, Al-Qutami HM, Khan MA. Nitric Oxide and calcium induced physio-biochemical changes in tomato (*Solanum lycopersicum*) plant under heat stress. *Fresen. Environ. Bull.* 2017;26(2a):1663-1672.
31. Zhang Z, Wu P, Zhang W, Yang Z, Liu H, Ahammed GJ, Cui J. Calcium is involved in exogenous NO-induced enhancement of photosynthesis in cucumber (*Cucumis sativus* L.) seedlings under low temperature. *Sci. Hortic.* 2020;261:108953.
32. Li Z, Tan XF, Lu K, Liu ZM, Wu LL. The effect of CaCl<sub>2</sub> on calcium content, photosynthesis, and chlorophyll fluorescence of tung tree seedlings under drought conditions. *Photosynthetica.* 2017;55(3):553-560.
33. Weng X, Li H, Ren C, Zhou Y, Zhu W, Zhang S, Liu L. Calcium regulates growth and nutrient absorption in poplar seedlings. *Front. Plant Sci.* 2022;13:887098.
34. Hepler PK, Winship LJ. Calcium at the cell wall-cytoplasm interface. *J. Integr. Plant Biol.* 2010;52(2):147-160.
35. Ekinci M, Estringü A, Dursun A, Yildirim E, Turan M, Karaman MR *et al.* Growth, yield, and calcium and boron uptake of tomato (*Lycopersicon esculentum* L.) and cucumber (*Cucumis sativus* L.) as affected by calcium and boron humate application in greenhouse conditions. *Turk J Agric For.* 2015;39(5):613-632.
36. Helal NA, AbdElhady SA. Calcium and potassium fertilization may enhance potato tuber yield and quality. *Middle East J.* 2015;4(4):991-998.
37. Handajarningsih M, Pujiwati H, Nasution DPP. Improving Yield and Performance of Shallot on Ultisol Through Application of Dolomite and Chicken Manure. In *International Seminar on Promoting Local Resources for Sustainable Agriculture and Development (ISPLRSAD 2020)* (pp. 310-313). Atlantis Press.

38. Agegnehu G, Amede T, Erkossa T, Yirga C, Henry C, Tyler R *et al.* Extent and management of acid soils for sustainable crop production system in the tropical agroecosystems: a review. *Acta Agric Scand B Soil Plant Sci.* 2021;71(9):852-869.

UNDER PEER REVIEW