

Original Research Article

Relationship of Maternal Serum Zinc Level with Preeclampsia

Abstract

A significant contributor to maternal and perinatal morbidities is preeclampsia. Zinc is a necessary trace element that is thought to affect renal salt transport and blood pressure management through its antioxidant capabilities. The goal of the study was to ascertain how preeclampsia and maternal serum zinc levels related to one another. This case-control study was conducted among 100 pregnant women at the Department of Obstetrics and Gynecology, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh. Among the 100 patients, 50 were pregnant women with preeclampsia (cases) and 50 were pregnant women without preeclampsia (controls). The serum zinc level was measured for all of them. The mean serum zinc level of the cases and controls were 59.58 ± 31.79 $\mu\text{g/dl}$ and 89.02 ± 31.14 $\mu\text{g/dl}$ respectively ($P < 0.001$). The respondents with zinc level < 68 $\mu\text{g/dl}$ had 6.05 times more chances to have preeclampsia compared to that of the respondents with zinc level ≥ 68 $\mu\text{g/dl}$ ($p < 0.001$; OR=6.048; 95% CI=2.540-14.399). There was significant negative correlation of serum zinc level with both systolic blood pressure ($r = -0.340$, $p = 0.001$) and diastolic blood pressure ($r = -0.404$, $p < 0.001$). In conclusion, low maternal serum zinc level was found related to preeclampsia.

Keywords: Preeclampsia, Serum zinc level.

Introduction

“Preeclampsia (PE) is characterized by new onset of hypertension (over 140 mmHg systolic or over 90 mmHg diastolic) after 20 weeks of pregnancy and the coexistence of 1 or more of the new-onset conditions: proteinuria, other maternal organ dysfunction like renal impairment, hepatic dysfunction, neurological complications or haematological complications (thrombocytopenia, disseminated intravascular coagulation) and/or uteroplacental dysfunction such as fetal growth restriction, abnormal umbilical artery doppler waveform analysis or stillbirth” [1]. An estimated 2-10% of pregnancies are complicated by preeclampsia globally, while its prevalence in developing nations ranges from 1.8% to 16.7% [2]. In Bangladesh, the overall prevalence of preeclampsia is 14.4% [3]. Complications of preeclampsia are maternal such as eclampsia, abruptio placentae, renal failure, cerebral haemorrhage, HELLP syndrome, pulmonary edema and visual loss. “Risks to the fetus include fetal growth restriction (FGR), premature birth, low APGAR score, stillbirth, neonatal intensive care unit admission and neonatal death. Since preeclampsia is caused by multiple factors, finding any factor related to this disorder can help in its prevention and can reduce the maternal and perinatal mortality and morbidity. As micronutrients are essential for normal physiological function, growth and development, the possible role of maternal serum zinc status also needs to be evaluated to understand the precise etiopathogenesis of preeclampsia and therefore limit its severity” [4].

“Proper implantation and angiogenesis are essential for the maintenance of a healthy pregnancy. The invading cells of embryo are called cytotrophoblasts. The process of invasion of cytotrophoblasts requires enzymatic digestion of the extracellular matrix (ECM) by matrix metalloproteinase enzyme (MMP-9) which is a zinc endopeptidase [5]. A study proves that zinc chelation diminishes the activity of MMP-2 and MMP-9” [5]. “Deficient concentrations

of zinc during pregnancy may cause impairment of antioxidant potential of cells by decreasing superoxide dismutase activity, as well as increased lipid peroxidation. Lipid peroxides formed at a primary site are then transported through the circulation by lipoproteins causing damage at distant tissues. The endothelial cells lining the high-pressured arterial vessels are particularly vulnerable to oxygen free radical initiated lipid peroxidation. Impaired function of generalized vascular endothelium may, in turn, cause vasospasm and an increase in sensitivity to vasopressors occurring in preeclampsia. This suggests the possibility of zinc deficiency in the incidence of preeclampsia” [6,7].

“In recent years, it was estimated that over 80% of pregnant mothers worldwide have inadequate zinc intake due to increased demand, low-zinc diets, and the consumption of foods that interfere with zinc absorption” (WHO, 2016) [8]. “Several groups of researchers emphasized the contribution of zinc deficiency in the aetiology of preeclampsia” [9,10,11]. Meanwhile, some studies found that the serum zinc concentrations were not significantly different between the preeclampsia patients and the healthy pregnant women [12,13,25]. So, this study has been designed to estimate the serum zinc level and determine its association with preeclampsia in women **in** our country.

Materials and Methods

This prospective case control study was conducted in the department of Obstetrics & Gynecology of Bangabandhu Sheikh Mujib Medical University (BSMMU), during March 2020 to February 2022 after obtaining Institutional Review Board approval. A total of 100 study subjects were enrolled upon fulfilling inclusion and exclusion criteria attending the outpatient and inpatient department of Obstetrics & Gynecology BSMMU. Preeclampsia patients (n = 50) and preeclampsia-free women (n = 50), who were matched for age and gestational age, made up the two groups that made up the study population. Cases had to be singleton pregnancies with preeclampsia and be between the ages of 18 and 40, with gestational ages ranging from 28 to 40 weeks. Control group bear the above-mentioned same characteristics except normotensive as determined by measurement of BP. Exclusion criteria are: multiple pregnancy, previous history of preeclampsia, pregnant women with raised uric acid level, diagnosed cases of chronic renal disease, diagnosed case of chronic hypertension, diagnosed case of cardiovascular disease, and patients who were taking zinc supplementation. Detailed medical, family, and socioeconomic histories were taken and recorded. Thorough physical examination of the subjects was done. Blood pressure was measured and recorded carefully. Urine analysis was done to measure the degree of proteinuria. Maternal blood samples were tested for serum zinc level in the Department of Biochemistry & Molecular Biology of BSMMU. Unpaired t-test, chi square test and Pearson’s correlation coefficient test were done for statistical analysis. Odds ratio (at 95% CI) was calculated to see the association of maternal serum zinc level with preeclampsia. A ‘p’value <0.05 was considered as statistically significant.

Results

The clinical characteristics as age, socioeconomic status, gravidity, gestational age, body mass index (BMI) and family history of hypertension are shown in table - 1 and table – 2 respectively. The mean maternal zinc level of the respondents is shown in table -3 which revealed that the zinc level of the cases was low (59.58 ± 31.79 µg/dl) than the control (89.02 ± 31.14 µg/dl) which is significant ($p = 0.025$). Table 4 shows that the pregnant women with zinc level <68 µg/dl had 6.05 times more chance to have preeclampsia compared to those of the pregnant women with zinc level ≥ 68 µg/dl ($p < 0.001$; OR=6.048; 95% CI=2.540-14.399). Negative correlation ($r = -0.340$.) was observed between maternal serum zinc level and

systolic blood pressure in the study subjects which is statistically significant ($p=0.001$) (Fig. 1). Similar negative correlation ($r= -0.404$) was observed between maternal serum zinc level and diastolic blood pressure which is also statistically significant ($p<0.001$) (Fig. 2).

Table 1. Sociodemographic data of the respondents

Characteristics	Case (n=50)	Control (n=50)	p-value
Age (Mean \pm SD)	27.18 \pm 6.03	26.56 \pm 5.46	0.591 ^a
Monthly family income (in taka) *			
Lower class (\leq 7,378)	8 (16.0)	4 (8.0)	0.363 ^b
Lower middle class (7,379–28,810)	26 (52.0)	25 (50.0)	
Upper middle class (28,811–89,280)	16 (32.0)	21 (42.0)	

^aChi square test was done to measure the level of significance.

^bUn-paired t-test was done to measure the level of significance.

Figure within parentheses indicates in percentage.

* World Bank Data Team on July 2020

Table 2. Clinical characteristics of the respondents

Characteristics	Case (n=50)	Control (n=50)	p-value
Gravida			
Primigravida	31 (62.0)	28 (56.0)	0.542 ^a
Multi gravida	19 (38.0)	22 (44.0)	
Gestational age in weeks (Mean \pm SD)	32.94 \pm 1.86	33.54 \pm 2.19	0.143 ^b
Family history of hypertension			
Present	17 (34.0)	9 (18.0)	0.068 ^a
Absent	33(66.0)	41 (82.0)	
BMI (kg/m^2) Mean \pm SD	26.68 \pm 1.78	26.02 \pm 1.56	0.053 ^b

^aChi square test was done to measure the level of significance.

^bUn-paired t-test was done to measure the level of significance.

Figure within parentheses indicates in percentage.

Table 3. Mean maternal zinc level of the respondents

Mean maternal zinc level ($\mu\text{g}/\text{dl}$)	Case (n=50)	Control (n=50)	p-value
Mean \pm SD	59.58 \pm 31.79	89.02 \pm 31.14	<0.001 ^a

^aUn-paired t-test was done to measure the level of significance.

Table 4. Odds ratios (OR) and 95% confidence intervals (CI) for preeclampsia according to maternal serum zinc level in pregnancy

Maternal	Case	Control	p-value	Odd's Ratio	95% CI (lower-
----------	------	---------	---------	-------------	----------------

serum zinc level ($\mu\text{g}/\text{dl}$)	(n=50)	(n=50)			upper
< 68 $\mu\text{g}/\text{dl}$	34 (68.0)	13 (26.0)	<0.001 ^a	6.048	(2.540-14.399)
\geq 68 $\mu\text{g}/\text{dl}$	16 (32.0)	37 (74.0)			

^aChi-square test was done to measure the level of significance.

Figure within parentheses indicates in percentage




Figure 1: Scatterplot diagram showing correlation between serum zinc level and systolic blood pressure (SBP) in patients with preeclampsia.




Figure 2: Scatterplot diagram showing correlation between serum zinc level and diastolic blood pressure (DBP) in the study subjects.

Discussion

In this study, the mean age of the respondents was 27.18 ± 6.03 years in cases and 26.56 ± 5.46 years in control. Majorities of the respondents from both the groups belonged to lower-middle-class social status (case: 52.0% and control: 50.0%). There was no significant difference in the distribution of the respondents in both groups regarding socio-demographic characteristics ($p > 0.05$). This result is consistent with the study of Akhtar et al. where the mean age was 25.11 ± 5.66 years in preeclamptic and 25.20 ± 4.85 years in normotensive pregnant women [14]. However, these findings contrast with the results of Morikaw et al. who recognized that an advanced maternal age ≥ 35 years and increased pre-pregnancy BMI were risk factors for pregnancy induced hypertension among pregnant Japanese women [15]. Differences in the study design and the demographic characteristics of the population may account for these differences in the results.

The present study revealed slightly lower gestational age among the cases (32.94 ± 1.86 weeks) compared to the controls (35.54 ± 2.19 weeks). About three-fifths (62.0%) of the preeclamptic patients and 56.0% healthy pregnant women were found to have primigravida but these were not statistically significant. In a similar study, Martadiansyah et al. also exhibited no difference in terms of gestational age ($p = 0.347$) and parity ($p = 0.940$) between cases and controls [16].

Family history of hypertension was present among 34.0% of the cases compared to 18.0% of the control group of respondents, this difference was found statistically not significant ($p = 0.068$) in the present study. In another study it was observed that the proportion of participants did not significantly differ with regards to family history of preeclampsia ($p = 0.704$) [17]. The mean BMI of the cases and control were 26.68 ± 1.78 kg/m^2 and 26.02 ± 1.56 kg/m^2 respectively which was not statistically significant ($p > 0.05$). In a systematic review and meta-analysis Tesfa et al. also noted a nonsignificant relation of BMI

between preeclampsia and the control group of patients ($p=0.057$) [18]. In contrast to this study, Al-Sakarneh and Mashal observed that during pregnancy the BMI of preeclamptic women was significantly higher than the normotensive women (34.4 ± 1.1 kg/m² vs. 30.08 ± 1.0 kg/m², $p=0.02$) [19].

The mean (\pm SD) serum zinc level was much lower in the case (59.58 ± 31.79 μ g/dl) than the control (89.02 ± 31.14 μ g/dl), which was statistically highly significant. Patients having zinc level < 68 μ g/dl had 6.05 times more chance to develop preeclampsia compared to those with zinc level ≥ 68 μ g/dl (OR=6.048, 95% CI = 2.540-14.399; p value < 0.001). Kumru et al. reported a significant reduction (43%) of serum zinc level in Turkish females with preeclampsia than in controls ($p<0.001$) [6]. A study by Akhtar et al. observed that there was a significant difference in zinc levels between pregnant women with preeclampsia and normal pregnancies (902.50 ± 157.15 μ g/L vs. 1153.33 ± 67.09 μ g/L; $p < 0.001$) [14]. Mohamed et al. exhibited zinc levels in the control group at 95.7 μ g /dl and in the preeclamptic group at 60.8 μ g dl with highly significant statistical difference ($p<0.001$) [20]. Emam and Attia evidenced that significantly lower levels of zinc (PE group: 58.94 ± 11.19 μ g/dl vs. control group 85.92 ± 13.8 μ g/dl, $p<0.001$) was associated with the development of preeclampsia (OR 4.47, 95% CI 3.05-6.52; p value < 0.001) and were inversely and significantly correlated with systolic ($r= -0.493$, $p=0.01$) and diastolic blood pressure ($r= -0.520$, $p=0.01$) [21]. In this study we also observed negative correlation ($r=-0.340$) between maternal serum zinc level and systolic blood pressure which was statistically significant ($p=0.001$). Similar negative correlation ($r= -0.404$) was observed between maternal serum zinc level and diastolic blood pressure which is also statistically significant ($p<0.001$).

In contrast, Adam et al. found no significant difference ($p>0.05$) in the mean serum zinc levels between preeclamptic (31.2 ± 5.2 μ g/dl) and normal pregnant women (34.1 ± 4.4 μ g/dl) [22]. Another study conducted by Magri et al. did not find a relationship between the serum levels of calcium, magnesium and zinc and gestational hypertension; therefore, they proposed that these elements might not clinically participate in the pathogenesis of the gestational hypertension [23]. These discrepancies of the study findings might be due to methods for measuring serum zinc not being sensitive or specific enough, variations in sample sizes and other demographic factors. A meta-analysis revealed that a lower level of maternal zinc was associated with increased risk of preeclampsia but not entirely consistent across countries and regions worldwide.[24].

With a few notable exceptions, the majority of studies found that preeclampsia patients had low serum zinc levels. Therefore, we may take some steps to improve the maternal and perinatal outcome, such as eating a diet rich in micronutrients and supplementing with zinc to overcome zinc insufficiency.

Conclusion

This study finding suggests that low maternal serum zinc level is significantly related with an increased risk for preeclampsia. The serum zinc level was found inversely correlated with both systolic and diastolic blood pressure. Therefore, this study concludes that a low level of serum zinc can be considered an important risk factor for developing preeclampsia.

Consent

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

Ethical Approval:

As per international standard or university standard guideline participant consent and ethical approval has been collected and preserved by the authors.

References

1. NICE. (2021) Pre-eclampsia. [Online] National Institute of Health and Care Excellence. Available: <http://pathways.nice.org.uk/pathways/hypertension-inpregnancy> [Accessed 26 September 2021].
2. Osungbade KO and Ige OK. Public Health Perspectives of Preeclampsia in Developing Countries: Implication for Health System Strengthening. *Journal of Pregnancy*. 2011; 2011:1-6.
3. Mou AD, Barman Z, Hasan M, Miah R, Hafsa JM, Das Trisha A et al. Prevalence of preeclampsia and the associated risk factors among pregnant women in Bangladesh. *Scientific Reports*. 2021; 11(1): 1-9.
4. Black RE. Micronutrients in pregnancy. *British Journal of Nutrition*. 2001; 85 (S2): S193-S197.
5. Thomas JM, Shenoy RP, Bhat PV and Rao P. Pathophysiology of preeclampsia and possible role of zinc in its genesis. *Current Women's Health Reviews*. 2014;10(1): 38-42.
6. Kumru S, Aydin S, Simsek M, Sahin K, Yaman M and Ay G. Comparison of serum copper, zinc, calcium and magnesium levels in preeclamptic and healthy pregnant women. *Biological Trace Element Research*. 2003; 94(2): 105-112.
7. Thakur S, Gupta N and Kakkar P. Serum copper and zinc concentrations and their relation to superoxide dismutase in severe malnutrition. *European Journal of Pediatrics*. 2004;163(12): 742-744.
8. WHO. (2016). WHO recommendations on antenatal care for a positive pregnancy experience. [Online] World Health Organization: Geneva, Switzerland, pp.1-172. Available:<https://apps.who.int/iris/bitstream/handle/10665/250796/97892415?sequence=1> [Accessed: 8 February 2022]
9. Akhter S, Ali T, Begum S and Ferdousi S. Micronutrient deficiency in severe preeclampsia. *Journal of Bangladesh Society of Physiologist*. 2013; 8(1):26-32.
10. Asliraf M, Rtd AS and Khurshid R. Maternal serum zinc concentration in gravidae suffering from pre-eclampsia. *APMC-Annals of Punjab Medical College*. 2007; 1(1): 24-27.
11. Nourmohammadi I, Akbaryan A, Fatemi S, Meamarzadeh A and Noormohammadi E. Serum Zinc Concentration in Iranian Pre-eclamptic and Normotensive Pregnant Women. *World Family Medicine*. 2008; 6(4):30-32.
12. Golmohammad LS, Amirabi A, Yazdian M and Pashapour N. Evaluation of Serum Calcium, Magnesium, Copper, and Zinc Levels in Women with Pre-eclampsia. *Iranian Journal of Medical Sciences*. 2008;33(4):231-234.
13. Rafeinia A, Tabandeh A, Khajeniazi S and Marjani AJ. Serum copper, zinc and lipid peroxidation in pregnant women with preeclampsia in gorgan. *The Open Biochemistry Journal*. 2014; 8: 83-88.

14. Akhtar S, Begum S and Ferdousi S. Calcium and zinc deficiency in preeclamptic women. *Journal of Bangladesh Society of Physiologist*. 2011;6 (2):94-99.
15. Morikawa M, Yamada T, Yamada T, Sato S, Cho K and Minakami H. Effects of nulliparity, maternal age, and pre-pregnancy body mass index on the development of gestational hypertension and preeclampsia. *Hypertension Research in Pregnancy*. 2013;1(2) :75-80.
16. Martadiansyah A, Maulina P, Mirani P and Kaprianti T. Zinc Serum Maternal Levels as a Risk Factor for Preeclampsia. *BioscientiaMedicina: Journal of Biomedicine and Translational Research*. 2021; 5(7) :693-701.
17. Chababa L, Mukosha M, Sijumbila G and Vwalika B. Relationship between serum zinc levels and preeclampsia at the University Teaching Hospital, Lusaka, Zambia. *Medical Journal of Zambia*. 2016; 43(3): 139-144.
18. Tesfa E, Nibret E and Munshea A. Maternal Serum Zinc Level and Preeclampsia Risk in African Women: a Systematic Review and Meta-analysis. *Biological Trace Element Research*. 2021;199(12): 4564-4571.
19. Al-Sakarneh NA and Mashal RH. Evaluation of Zinc and Homocysteine Status in Pregnant Women and Their Association with Pre-eclampsia in Jordan. *Preventive Nutrition and Food Science*.2021; 26(1): 21-29.
20. Mohamed ASS, El-Omda FA, Abdelfatah AT and Hashish MA. Comparative Study for Serum Zinc and Copper Levels in Cases with Normal Pregnancy Versus Preeclampsia. *The Egyptian Journal of Hospital Medicine*.2019;74(5): 1069-1074.
21. Emam AH and Attia AM. Selected Trace Elements in Egyptian Females with Preeclampsia. *International Journal of Medical Arts* .2020; 2(4): 811-819.
22. Adam B, Malatyalioglu E, Alvun M and Talu C. Magnesium, zinc and iron levels in pre-eclampsia'. *The Journal of Maternal-Fetal Medicine*.2001;10:1-5.
23. Magri J, Sammut M and Savona-Ventura C. Lead and other metals in gestational hypertension. *International Journal of Gynecology& Obstetrics*. 2003; 83(1): 29-36.
24. Jin S, Hu C, Zheng Y. Maternal serum zinc level is associated with risk of preeclampsia: A systematic review and meta-analysis. *Front Public Health*. 2022 Aug 1;10:968045. doi: 10.3389/fpubh.2022.968045. PMID: 35979462; PMCID: PMC9376590.
25. Sheikh ZS, Ali AM. Association of gestational diabetes mellitus with vitamin D deficiency among Saudi pregnant women: A case-control study. *Journal of Fatima Jinnah Medical University*. 2019 Dec 10;13(3):96-101.