

# Individual-based modelling for predicting dry matter accumulation in juveniles of Indian Olive raised under different soil treatments

Comment [a1]: Delete the extra space

## ABSTRACT

Dry matter accumulation data for juvenile Indian Olive saplings was modelled using linear and non-linear regression models. The data was gathered bimonthly and for two years i.e., twelve readings. Linear, Quadratic, Power, Sigmoid and Exponential models performed quite well for cumulative growth data (Adj.  $R^2 > 0.93$ ). The Sigmoid model (Chapman- Richard) demonstrated upper asymptotic values ranged between 179.96- 334.49 grams for all treatments ( $T_0 - T_4$ ) suggesting that olive saplings responded positively to different soil treatments. Both, age and size-dependent RGR manifested low values and decrease in RGR curve. Age-dependent RGR displayed a smooth decline in function, indicating extended longevity. Low RGR (along with low SLA), is a characteristic feature of slow growing and evergreen species and this trait was observed in Indian Olive growth too. Size- standardized RGR revealed that due to low rate of decline  $T_4$  had the superior RGR (and maximum dry matter accumulation) followed by other treatments. Moreover,  $T_0$  had the sharpest decrease and the lowest dry matter accumulation. Cross- validation of model parameters exhibited statistical stability.

Comment [a2]: , and

Comment [a3]: The low

Comment [a4]: Decline,

Keywords: Indian olive, regression models, Chapman- Richard, upper asymptote, RGR Relative growth rate"

Comment [a5]: Add the expression

## 1. INTRODUCTION

Plant simulation modelling is a robust and advance technique which is applied too often to model growth patterns of individual plant species as well as populations. The modelling system consists of a large number of mathematical functions which are mostly nonlinear in nature and with or without an upper asymptote. The upper stable value signifies that growth of an organism is always restricted due to competition between individuals, limited resources and environmental constrains. The growth curves of large number of organisms including plant groups forms a sigmoid curve where initially the growth is exponential; in the middle part the growth takes a linear trajectory and eventually in the final phase it levels off or straightens thus imitating an S-shaped curve. Generally, nonlinear and asymptotic models are used to simulate growth patterns of plants. The cumulative growth is represented by a typical sigmoid curve, and absolute growth by a bell-shaped curve, however relative growth curve is generally negative exponential or reverse sigmoidal in shape due to the fact that relative growth gradually decreases as the plant or organ size increases with age.

Comment [a6]: The

Comment [a7]: ,

The tree species selected for present investigation is an Indian Olive (*Olea ferruginea* Royle; syn. *O. cuspidate* Wall. Ex G. Don). It is a wild and indigenous species and has a wide distribution in the Himalayas from Kashmir to Kumaun, up to an altitude of 2400 m. It is one of the six species of *Olea* found in India. It is also noticed in the neighboring countries of Nepal, Pakistan and Afghanistan. It thrives well in wastelands where soil is not suitable for other plants, that is why this species has been utilized as stock in grafting experiments along with *Olea europaea* (an exotic species from Europe).

Comment [a8]: The present

Comment [a9]: The soil

Indian olive is a medium-sized, slow growing, broad-leaved and evergreen tree with an average height of approximately 10.5 m and an average diameter of 0.45 m. The trees constitute important communities of lesser Himalayas and provide wide variety of economic and ecological services to local people in their vicinity. Other fascinating characteristic includes: it is a zero-waste, multipurpose tree which provides construction wood, food, firewood, fodder and medicinal products to different ethnic groups since decades. Leaves are used as an antiperiodic in fever and debility [1], in mouth ulcer, toothache, sour throat, gonorrhoea, hoarse voice, as an astringent, antiseptic and a mild digestive aid [2.3]. Timber is heavy and resilient and is utilized for manufacturing agricultural tools [1]. Fruits are edible and are used as antidiabetics, appetizers, and a rich source of Oleic acid (from Olive oil). The oil is extensively used in jaundice, scabies, typhoid, burning of eyes, caries of the teeth and tooth ache [3.4]. The olive oil is further used in joint pains,

malaria, rheumatism, gonorrhoea, skin disease, cosmetics and for cooking as well [4, 5]. Root is used in rheumatism, headache, asthma and scorpion sting [6].

Due to its zero waste and multipurpose qualities, the Indian Olive growth pattern was investigated in an experimental plot at Almora applying different soil treatments. Healthy and disease-free saplings were selected and the dry matter accumulation data was recorded bimonthly for almost two years i.e., 24 months (12 readings). The cumulative data obtained from treatment T<sub>0</sub> for juvenile trees was initially screened using seven different regression models (both linear and nonlinear). Out of seven models tested, a sigmoid model (Chapman-Richard's curve) was applied to simulate cumulative and relative growth of trees for remaining treatments (T<sub>1</sub>- T<sub>4</sub>) Results of the modelling is being interpreted in the present manuscript.

Comment [a10]: Are

## 2. MATERIAL AND METHODS

### 2.1 Site details, experimental plot and data collection

The aim of this investigation was to model dry matter accumulation data over time during the initial developmental stages of Indian olive saplings. The present study was conducted at the experimental plots in Chausar village at Almora. Site coordinates were: 29.5892° N, 79.6467° E and it was situated at an elevation of 1642 m from sea level.

**Table 1. Mean dry matter accumulation (gm/sapling) for Indian Olive at all sampling stages i.e., Aug 2021 to June 2023 under different soil treatments (n=3, ±SE)**

Harvest (months)	T0	T1	T2	T3	T4
2	3.64±0.22	5.92±0.52	6.84±0.44	7.37±1.01	8.46±1.11
4	3.88±0.16	6.92±1.07	7.44±1.31	7.96±1.41	9.06±0.87
6	4.22±0.35	7.26±1.18	7.82±0.81	8.33±1.59	9.75±1.27
8	4.7±0.34	7.73±1.04	8.24±1.01	8.73±1.36	10.48±1.73
10	5.13±0.41	8.51±0.84	9.09±1.44	9.69±0.97	11.3±1.45
12	5.54±0.56	9.21±0.72	9.73±1.81	10.27±1.36	12.25±0.75
14	6.07±0.78	9.7±0.98	10.32±1.52	10.89±1.65	13.09±1.27
16	6.63±0.25	10.41±1.21	11.02±0.98	11.64±1.36	14.05±1.18
18	7±0.36	10.99±0.89	11.62±0.87	12.42±1.47	15.18±1.41
20	7.51±0.58	11.78±1.39	12.58±1.44	13.35±0.95	16.28±1.04
22	7.93±0.83	13.33±1.56	13.84±1.85	14.66±1.53	17.51±1.05
24	8.37±0.97	14.27±0.95	15.07±0.58	15.95±2.17	18.42±1.53

The experimental material consisted of 180 healthy, uniform and disease-free saplings of Indian Olive (*Olea ferruginea* Royle), procured from the Indo-Italian Government Demonstration farm, Dhakrani (Dehradun) and replanted in the experimental plots at village Chausar. The experiment was laid down in a randomized block design with thirty-six (36) saplings in each design and the data were analyzed accordingly. A total of 180 saplings were sacrificed to obtain the dry matter accumulation data of two years. Data was collected bimonthly that is, every sixty days interval. The experiment was conducted with three levels of Farm Yard Manure (FYM) viz. 0, 15 and 30 tons/ha and different combinations of N, P and K (Table 2) during 2021-23 at the experimental plots. Each of the combination was termed as treatment from T<sub>0</sub> to T<sub>4</sub>.

The details of the physio-chemical characteristic of the soil used for the purpose was as follows: Texture (sandy loam), Sand (67.16%), Silt (22.29%), Clay (10.55%), Bulk Density (1.36 gm/cc), Water Holding Capacity (40.65%), pH (6.8), Organic Carbon (0.48%), Organic Matter (0.83%), Total Nitrogen (0.08%), Available Nitrogen (138.3 Kg/ha) and Available Potassium (198.4 Kg/ha).

### 2.2 Modelling dry matter accumulation over time (24 months)

Initially the dry matter accumulation data over time for treatment T<sub>0</sub> was screened using seven (7) regression models. The cumulative growth function and performance statistics for all models is depicted in Table 4. In the subsequent step, a 3-parametric Chapman-Richard's function (C-R) (Table 3) was selected to calculate RGR for treatment T<sub>0</sub> along with CGR and RGR for remaining four treatments i.e., T<sub>1</sub>- T<sub>4</sub>. C-R function was applied to simulate the above data as all classical sigmoidal growth functions have two smooth transitions in a single formula and are quite flexible, versatile and reliable in nature.

**Table 2. Details of the treatments namely FYM (Tons/ha) and N, P and K (Kg/ha) on Indian Olive saplings at experimental plot at Almora**

Treatments	FYM (F) and N, P, K
T <sub>0</sub>	F <sub>0</sub> + N <sub>0</sub> + P <sub>0</sub> + K <sub>0</sub>
T <sub>1</sub>	F <sub>15</sub> + N <sub>100</sub> + P <sub>80</sub> + K <sub>50</sub>
T <sub>2</sub>	F <sub>15</sub> + N <sub>150</sub> + P <sub>120</sub> + K <sub>75</sub>
T <sub>3</sub>	F <sub>30</sub> + N <sub>0</sub> + P <sub>0</sub> + K <sub>0</sub>
T <sub>4</sub>	F <sub>30</sub> + N <sub>150</sub> + P <sub>120</sub> + K <sub>75</sub>

The function is frequently used to model growth of individual trees and populations. Here,  $\alpha$  is the upper asymptote,  $\beta$  is the growth rate or scale parameter and  $\delta$  is the shape parameter and inflection point. Cumulative growth was calculated for all treatments. Eventually relative growth was also calculated by using absolute growth values (Table 3).

**Table 3: Chapman- Richard's model[7] describing absolute and relative growth rate functions**

Model	Absolute Growth Rate	Relative Growth Rate
Chapman- Richard's	$y = \alpha\beta\delta e^{-\beta t} (1 - e^{-\beta t})^{\delta-1}$	$y = \beta\delta(1 - e^{-\beta t})^{-1}$

### 2.3 Model fitting, evaluation and validation

Nonlinear curve fitting was performed with Excel Solver which is an add-in function in Microsoft Excel, 2021. Three criteria were applied for model evaluation which were: Adjusted- R<sup>2</sup>[8], Akaike Information Criterion[9], and Bayesian Information Criterion[10]. Model with maximum Adjusted- R<sup>2</sup> and minimum AIC and BIC values were considered to perform the best. Cross- validation of model parameters was conducted through Jackknife resampling technique [11].

### 3.0 RESULTS AND DISCUSSION


Table 4 describes fitting and evaluation statistics of cumulative growth for treatment T<sub>0</sub>. Though, linear quadratic and exponential models were slightly better than Chapman- Richard's model but the latter was considered to further evaluate CGR and RGR values for remaining treatments due to the fact that it a sigmoid function which responded quite well to dry matter accumulation growth of juvenile olive saplings and was a combination of linear, power and exponential functions. The function was quite flexible, accurate and had an upper asymptote. Model parameters had biological meaning too. The upper asymptotic values for all treatments were in the range 179.96- 334.49 grams. The highest dry matter was estimated for treatment T<sub>4</sub> and lowest for treatment T<sub>0</sub>. The dry matter production had a wide range suggesting that olive saplings responded differently to different soil treatments. T<sub>0</sub> treatment, that is, normal soil (no FYM and NPK) displayed least value which confirmed that different treatments of NPK and FYM had a positive impact on dry matter accumulation in early stages of sapling growth. All the treatments followed the order: T<sub>4</sub>> T<sub>1</sub>> T<sub>3</sub>> T<sub>2</sub>> T<sub>0</sub>. Model parameters were statistically stable.

RGR functions are unique in capturing both age and size dependent changes in growth patterns. As RGR itself is size- dependent and declines as individual tree grows so size- correction or size- standardization is always a better option when species are compared at a common size. Both age and size dependent RGR demonstrated a decrease in RGR curve, which is commonly referred to as ontogenetic drift, however, the former displayed a smooth decline for all treatments indicating extended longevity (Fig. 2a). Self- shading of leaves, decrease in local concentration of soil nutrients and accumulation of non- photosynthetic biomass in the form of stem and roots could be the possible reasons for universal decrease in RGR as the tree grows in nature. The initial values for age- dependent RGR were low and ranged between (0.151- 0.172). Low RGR is a characteristic feature of trees with a slow growing nature and evergreen habit.

Size- standardization displayed the trend: T<sub>4</sub>>T<sub>3</sub>> T<sub>2</sub>> T<sub>1</sub>> T<sub>0</sub> i.e., due to low rate of decline T<sub>4</sub> had the superior RGR followed by other treatments (Fig. 2b). Moreover, T<sub>0</sub> had the sharpest RGR decline and the lowest dry matter accumulation (Fig. 2b). T<sub>0</sub> started with the highest RGR value but finished with lowest dry matter accumulation. At a common biomass value of 6.5 gm, T<sub>4</sub> had a superior RGR (0.15) than that of T<sub>0</sub> (0.02). The RGR for treatment T<sub>4</sub> was 13% more in magnitude than RGR for treatment T<sub>0</sub>.

**Table 4. Model performance of the dry matter accumulation data of juvenile Indian olive saplings for treatment T<sub>0</sub>, α, β and δ are model parameters and “t” denotes time of harvest in months**

Models	Functional Form	Performance criteria		
		Adj. R <sup>2</sup>	AIC	BIC
Linear	$y = \alpha + \beta t$	0.996	-54.204	-53.234
Logarithmic	$y = \alpha + \beta \log(t)$	0.832	-9.323	-8.354
Inverse	$y = \alpha + \beta/t$	0.601	1.068	2.038
Quadratic	$y = \alpha + \beta t + \delta t^2$	0.997	-58.178	-56.724
Power	$y = \alpha t^\beta$	0.936	-20.815	-19.845
Sigmoidal (C-R)	$y = \alpha(1 - e^{-\beta t})^\delta$	0.938	-20.777	-19.322
Exponential	$y = \alpha e^{\beta t}$	0.989	-42.206	-41.236



**Figure 1. Cumulative growth curves of juvenile Indian olive applying seven regression models\* Circles are observed values**

Evaluating growth rates is an important aspect in functional forestry. Relative growth rate (RGR) is a remarkable trait identifying plant growing strategy [12]. Species with high RGR are much more competitive in acquiring resources whereas slow growing species are more conservative with the limited resources they have obtained [13,14] or they are able to combat natural disturbances by allowing resources to storage rather than to grow [15]. In slow growing, evergreen trees low RGR is associated with a low SLA is a common feature. The present study also demonstrated low RGR for all treatments (T<sub>0</sub>- T<sub>4</sub>). Mean SLA was also calculated for 24 months (data not shown) and was also found to be consistently low with every harvest. As Indian Olive is an evergreen slow growing tree so low RGR (and SLA) values are not surprising at all. Ecological advantage of low RGR in slow-growing evergreen trees has been discussed in the past. Generally, evergreen trees thrive in temperate and sub-tropical area with less fertile soil. Some possible explanations for low RGR in slow growing and evergreen species could be: having low RGR, the plant functions close to its optimum growth rate in harsh conditions; the plants have lower rates of accumulation of minerals and photosynthates into structural material and finally due to their low demands slow growing species retains (holds) the resources for longer durations. Moreover, high leaf construction cost in evergreen trees could also be the reason for low SLA and RGR values.

Evergreen species are widely thought to have low RGR (slow-growing) and low SLA due to the need to produce thick leaves that sustain longer life spans [16]. For extended longevity and evergreen habit, the olive sapling needed to develop a mechanism which protects it from large number of biotic and abiotic stresses viz. nutrient leakage, drought, cold, herbivory and fungal infections to name a few. In order to maintain leaf longevity such species had to put extra investment in secondary compounds such as lignin and phenolics that eventually forms cell membranes. Moreover, leaf hairs and cuticle- waxes also increase the construction cost of leaves. Number of studies on different plant functional groups suggested that low RGR along with low SLA is an important feature linked with evergreen habits.

Comment [a11]: of

Comment [a12]: in a temperate

Comment [a13]: areas

Comment [a14]: the plants have lower rates of accumulation of minerals and photosynthates into a structural material. Finally, slow-growing species retain (hold) the resources for longer durations due to their low demands. Moreover, high leaf construction costs in evergreen trees could also be the reason for low SLA and RGR values.

Comment [a15]: That

Comment [a16]: A large

Comment [a17]: , and

**Table 5. Model fitting and validation statistics for all treatments (T<sub>0</sub>- T<sub>4</sub>) using a 3- parametric Chapman- Richard's function (\* ±SE in parenthesis); above values are the parameters for cumulative growth function and below for the relative growth function**

Treatments	Model Parameters		
	$\alpha$	$\beta$	$\delta$
T <sub>0</sub>	179.96 (6.27)	0.000033 (0.000008)	0.425 (0.006)
T <sub>1</sub>	254.15 (9.95)	0.00003 (0.000003)	0.407 (0.007)
T <sub>2</sub>	215.69 (8.41)	0.00003 (0.000004)	0.379 (0.007)
T <sub>3</sub>	213.62 (5.47)	0.000031 (0.000003)	0.373 (0.007)
T <sub>4</sub>	334.49 (9.94)	0.000024 (0.000005)	0.391 (0.006)
	-	$\beta$	$\delta$
T <sub>0</sub>	-	-0.357 (0.002)	0.777 (0.005)
T <sub>1</sub>	-	-0.239 (0.001)	1.389 (0.011)
T <sub>2</sub>	-	-0.239 (0.0003)	1.551 (0.0023)
T <sub>3</sub>	-	-0.229 (0.0009)	1.671 (0.0087)
T <sub>4</sub>	-	-0.184 (0.0068)	1.912 (0.0073)

#### 4.0 CONCLUSION

To conclude, growth modelling techniques are an important aspect of functional forestry where growth data can be collected over time, modelled and eventually interpreted to get the final outcome. In the present investigation, juveniles of Indian olive saplings were treated with five different soil treatments (T<sub>0</sub>- T<sub>4</sub>) and the dry matter accumulation data of two years was analyzed. T<sub>4</sub> came out with the best treatment with a smooth decline in RGR curve and maximum dry matter accumulation in two years duration. Here, low RGR (for all treatments) was found to be associated with low SLA which is a key feature in evergreen and slow-growing trees. Species with low SLA and RGR values are naturally conditioned for the conservation of acquired resources due to their large dry matter content, high concentration of cell walls and increased production of secondary metabolites as a part of defense infrastructure. Moreover, evergreen species with low SLA/RGR has high leaf and root longevity too. All such traits were observed in Indian olive species which naturally grows in wastelands and unfertile soils. Such characters are developed due to harsh micro-climatic conditions. As olive saplings responded positively to different soil treatments, so more combinations of FYM and NPK should be screened in future to come out with better growth, adaptation and survival models for the species.

**Comment [a18]:** The future

**Comment [a19]:** , and




Figure 2. Fitted values from a 3- parametric Chapman- Richard's function for (a) dry matter accumulation; (b) relative growth rate (RGR) on time basis and (c) RGR on size basis of juvenile Indian Olive (\*circles are observed values)

## REFERENCES

1. Anon., The Wealth of India, Vol. VII, CSIR, New Delhi, India, 1997.
2. Ahmad H., Ahmad A., Jan M.M. The medicinal plants of Salt Range, Journal of Biological Science. 2002;29: 175- 177.
3. Zabihullah Q, Rashid A., Akhtar N. Ethnobotanical survey in Kot Manzaray Baba valley, Malakand agency, Pakistan, Pakistan journal of Plant Sciences. 2006; 12:115- 121.
4. Ahmad S.S. Medicinal wild plants from Lahore- Islamabad motorway (M-2), Pakistan Journal of Botany. 2007; 39: 355- 375.
5. Ahmad S., Ali A., Bibi S., Marwat K.B, Hassan G. Ethnobotanical study on some medicinal plants of Ouch district lower Dir, Pakistan, Pakistan Journal of Plant Sciences. 2006; 12:65- 71.
6. Hussain F., Badshah L., Dastagir G. Folk medicinal uses of some parts of south Waziristan, Pakistan, Journal of Plant Sciences. 2006; 12: 27- 39.
7. Richards F.J. A flexible growth function for empirical use, Journal of Experimental Botany. 1959; 10: 290- 300.
8. Chenge, I.B. Height- Diameter relationship of trees in Omo Strict nature forest reserve, Nigeria, Trees Forest and People. 2021; 3.
9. Akaike H.A. A new look at the statistical model identification, IEEE Transactions on Automatic Control. 1974; 19(6): 716- 723. Doi: 10.1109/TAC.1974.1100705
10. Schwarz G. Estimating the dimension of a model, The Annals of Statistics. 1978; 6(2): 461- 464. <https://doi.org/10.1214/aos/1176344136>
11. Harris D.C. Non- linear least squares curve fitting with Microsoft Excel, Journal of Chemical Education. 1998; 75(1): 119- 121.
12. Reich P.B., Buschena C, Tjoelker M.G., et al., Variation in growth rate and ecophysiology among 34 grassland and savanna species under contrasting N supply: a test of functional group differences, New Phytologist. 2003a; 157: 617- 631.
13. Reich P.B., Wright I.J., Carine J.M., Oleksyn J, Westoby M, Walters M.B. The evolution of plant functional variations: traits, spectra and strategies, International Journal of Plant Sciences. 2003b; 164: S143- S164.
14. Poorter H, Garnier E. The ecological significance of variation in relative growth rate and its components In Pugnaire FI, Valladares F. eds. Functional Plant Ecology, 2<sup>nd</sup> edition Boca Raton, FL: CRC Press. 2007; 67- 100.
15. Poorter L, Kitajima K. Carbohydrate storage and light requirements of tropical moist and dry forest tree species, Ecology. 2007; 88: 1000- 1011.
16. Wright I.J., Reich P.B., Westoby M., et al., The worldwide leaf economics spectrum, Nature. 2004; 428: 821- 827.