

Influence of Phosphorus and Sulphur application on Sesamum yield in High Phosphorus soils of Telangana

Abstract:

The present study in pot experiments was conducted on “Influence of Phosphorus and Sulphur application on Sesamum yield in High Phosphorus soils of Telangana”. The experiment was conducted in two different levels of high phosphorus status soils with 67.29 kg P₂O₅ ha⁻¹ in soil 1 (S₁) and 83.46 kg P₂O₅ ha⁻¹ in soil 2 (S₂). The treatments were taken in factorial completely randomized design in combination of five levels of phosphorus (P₀₋₀, P₂₅₋₅, P₅₀₋₁₀, P₇₅₋₁₅ and P₁₀₀₋₂₀ kg ha⁻¹ of soil) four levels of sulphur (S₀₋₀, S₁₋₁₀, S₂₋₂₀ and S₃₋₃₀ kg ha⁻¹ of soil). A significant increase in seed and stalk yield of sesamum crop could be achieved by combined application of P₇₅ (15 kg P₂O₅ ha⁻¹) + S₂₀ (20 kg S ha⁻¹) in both high available phosphorus soils. Among the various treatments tested, maximum seed and stalk yield was obtained with combined application of P₇₅ (15 kg P₂O₅ ha⁻¹) + S₂₀ (20 kg S ha⁻¹) which was 11.89 g pot⁻¹ higher over control in S₁ and P₇₅ (15 kg P₂O₅ ha⁻¹) + S₂₀ (20 kg S ha⁻¹) which was 11.92 g pot⁻¹ higher over control in S₂.

Key words: Sesamum, high P soil, phosphorus and sulphurs

Introduction

Sesame (*Sesamum indicum* L.) belongs to family *Pedaliace* and it is fourth important edible oilseed crop next to the groundnut, rapeseed and mustard in India. About 78 per cent of the sesame produced in the country is used for oil extraction (Kalegore *et al.*, 2018). Sesame oil has 16-18% of carbohydrates and 42% essential linoleic acid. Its oil content generally varies from 46 to 52 per cent which is highly resistant to oxidative rancidity. As the protein content is around 25% it is called as “Queen of oilseeds”. Oilseeds are important constituent in human dietary system next to carbohydrates and proteins (Shelke *et al.*, 2014).oil is characterized for its stability and quality. Because of its excellent quality characters, sesame oil is some also referred to as “poor man’s substitute for ghee”. Sesame cake is eaten mixed with sugar by poor people and added to bread to improve palatability and nutritive value. It is also a valuable nutritious feed for milch animals and is ingredient of poultry feed as it contains 6.0-6.2% N, 2.0-2.2% P and 1.0-1.2% K. India ranks second in area, third in production but ranks seventh in productivity in world (FAO STAT, 2020). In India, sesame is cultivated in 17.22 lakh ha of area with production

6.57 lakh tonnes and productivity 474 kg ha⁻¹ during 2020-2021 (www.Indiastat.com). In Telangana the area under sesame was 0.23 L ha⁻¹ with production of 25,469 tonnes per acre and productivity 310 kg/acre during 2021-2022 (SYB, 2021). The major sesame growing districts of Telangana are Jagtial, Karimnagar, Nirmal, Nizamabad, Khammam Kamma Reddy, Pedda palli and Adilabad.

In intensively cultivated areas due to indiscriminate application of phosphorus was leading to phosphorus accumulation in top layers of soil. In some studies it was revealed that at high phosphorus status soils the recommended dose of phosphorus can be reduced without reduction in the economic yield of crop. Chaudhary and Thakur (2007) observed that P dose could be skipped in either of any crop or reduced to 50 per cent under rice-wheat sequence if soil contains high available phosphorus. Reduced P application by 25-50 per cent from the current RDP in high P soils is reported to be a possible saving measure without sacrificing the yields in crops like rice, sunflower and in rice-rice and rice-sunflower-cropping sequences (Babu *et al.*, 2005; Madhavi *et al.*, 2015; Ramya *et al.*, 2015 and Srinivas *et al.*, 2016).

Sulphur plays a key role in the plant metabolism, indispensable for the synthesis of essential oils, chlorophyll formation, required for development of cells and it also increases cold resistance and drought hardiness of crops especially for oilseed crops. Use of high analysis sulphur free fertilizers, heavy sulphur removal by the crops under intensive cultivation and neglect of sulphur replenishment contributed to widespread sulphur deficiencies in arable soils. Sulphur has become one of the major limiting nutrients for oilseeds in recent years due to its widespread deficiency. In many crop sequences involving oilseeds, sulphur application was also reported to be quite profitable.

Synergistic effect of application of P and S at lower levels and antagonistic effect at higher rates of P and S application to cowpea crop was reported by Shankarlingappa *et al.* (2000). The recent study conducted by Surendra Babu *et al.* (2015) indicated that application of S in small doses is useful to increase the yields of rice in high P soils even when the soil is having sufficient available S. The nutrient requirement and interaction of different nutrients under high phosphorus conditions were not worked out so far in sesamum track. To improve the productivity levels of sesamum it is imperative to assess the phosphorus and sulphur requirement under high P fertility soils to obtain the potential yield by the farmers.

Materials and Methods

Initially soils were screened for available phosphorus in and around Ranga Reddy district. Based on available phosphorus two soils were selected for conducting pot culture experiment with available phosphorus status S_1 ($67.29 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$) and S_2 ($83.46 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$) status. In the selected soils (S_1 and S_2) pot culture experiment was conducted in sesamum crop (JCS 1020) during summer 2022-23 at Agricultural Research Institute, Rajendranagar and Hyderabad. Initially experimental soil samples were analyzed for textural analysis (Bouyoucos, 1962) before conducting the experiment. The pH of the soil in 1: 2.5 soil water suspensions and electrical conductivity of the soil in 1:2.5 soil water extract (Jackson, 1973) and organic carbon (Walkley and Black, 1934) was done. Available nitrogen in the soil was determined by alkaline permanganate method (Subbiah and Asija, 1956) available phosphorus by Olsen's extractant (Olsen *et al.*, 1954), available potassium (Muhr *et al.*, 1965) and available sulphur was determined by turbidity method (Williams and Steinbergs, 1959).

Both the experimental soils were slightly alkaline with normal in reaction and low in organic carbon. The soils were low in available nitrogen, high in available potassium and medium in available sulphur. A total of twenty phosphorus and sulphur treatment combinations were adopted in both the soils with four levels of replication. The different treatment combinations comprises of five levels of phosphorus (P_0-0 , $P_{25}-5$, $P_{50}-10$, $P_{75}-15$ and $P_{100}-20 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ of soil) and four levels of sulphur (S_0-0 , $S_{10}-10$, $S_{20}-20$ and $S_{30}-30 \text{ kg S ha}^{-1}$ of soil) were presented in table 1 & 2 and it was laid out in Factorial Completely Randomized Design. The recommended dose of fertilizer of sesamum included $60-20-20 \text{ kg N, P}_2\text{O}_5 \text{ \& K}_2\text{O ha}^{-1}$ and the recommended dose of nitrogen (60 kg ha^{-1}), potassium (20 kg ha^{-1}) were applied uniformly to all the treatments through urea and potassium chloride while phosphorus and sulphur were applied through diammonium hydrogen phosphate and potassium sulphate. Nitrogen was applied 50 per cent basally while remaining was applied thirty days after application. Phosphorus, potassium and sulphur were applied as basally. All the prophylactic plant protection measures were adopted as and when needed during the crop growth period.

Table 1: Treatmental combinations in experimental soil 1 (67.29 kg P₂O₅ ha⁻¹)

Treatment	N(Kg ha ⁻¹)	P ₂ O ₅ (Kg ha ⁻¹)	K ₂ O(Kg ha ⁻¹)	S(Kg ha ⁻¹)
S ₁ P ₀ Su ₀	60	0	20	0
S ₁ P ₀ Su ₁₀	60	0	20	10
S ₁ P ₀ Su ₂₀	60	0	20	20
S ₁ P ₀ Su ₃₀	60	0	20	30
S ₁ P ₂₅ Su ₀	60	5	20	0
S ₁ P ₂₅ Su ₁₀	60	5	20	10
S ₁ P ₂₅ Su ₂₀	60	5	20	20
S ₁ P ₂₅ Su ₃₀	60	5	20	30
S ₁ P ₅₀ Su ₀	60	10	20	0
S ₁ P ₅₀ Su ₁₀	60	10	20	10
S ₁ P ₅₀ Su ₂₀	60	10	20	20
S ₁ P ₅₀ Su ₃₀	60	10	20	30
S ₁ P ₇₅ Su ₀	60	15	20	0
S ₁ P ₇₅ Su ₁₀	60	15	20	10
S ₁ P ₇₅ Su ₂₀	60	15	20	20
S ₁ P ₇₅ Su ₃₀	60	15	20	30
S ₁ P ₁₀₀ Su ₀	60	20	20	0
S ₁ P ₁₀₀ Su ₁₀	60	20	20	10
S ₁ P ₁₀₀ Su ₂₀	60	20	20	20
S ₁ P ₁₀₀ Su ₃₀	60	20	20	30

Table 2: Treatmental combinations in experimental soil 2 (83.46 kg P₂O₅ ha⁻¹)

Treatment	N(Kg ha ⁻¹)	P ₂ O ₅ (Kg ha ⁻¹)	K ₂ O(Kg ha ⁻¹)	S(Kg ha ⁻¹)
S ₂ P ₀ Su ₀	60	0	20	0
S ₂ P ₀ Su ₁₀	60	0	20	10
S ₂ P ₀ Su ₂₀	60	0	20	20
S ₂ P ₀ Su ₃₀	60	0	20	30
S ₂ P ₂₅ Su ₀	60	5	20	0
S ₂ P ₂₅ Su ₁₀	60	5	20	10
S ₂ P ₂₅ Su ₂₀	60	5	20	20
S ₂ P ₂₅ Su ₃₀	60	5	20	30
S ₂ P ₅₀ Su ₀	60	10	20	0
S ₂ P ₅₀ Su ₁₀	60	10	20	10
S ₂ P ₅₀ Su ₂₀	60	10	20	20
S ₂ P ₅₀ Su ₃₀	60	10	20	30
S ₂ P ₇₅ Su ₀	60	15	20	0
S ₂ P ₇₅ Su ₁₀	60	15	20	10
S ₂ P ₇₅ Su ₂₀	60	15	20	20

S ₂ P ₇₅ Su ₃₀	60	15	20	30
S ₂ P ₁₀₀ Su ₀	60	20	20	0
S ₂ P ₁₀₀ Su ₁₀	60	20	20	10
S ₂ P ₁₀₀ Su ₂₀	60	20	20	20
S ₂ P ₁₀₀ Su ₃₀	60	20	20	30

Results and Discussion:

Effect of different doses of P and S levels on seed and stalk yield of sesamum was presented in table 3 and 4. Application of different levels of phosphorus fertilizer to sesamum crop in high P soils significantly influenced the sesamum seed yield. It was found that increasing the phosphorus application from P₀ to P₇₅ (15 kg P₂O₅ ha⁻¹) enhanced the sesamum seed yield by 13.82% significantly from 10.01 to 11.39 g pot⁻¹. Subsequent enhancement of phosphorus application from P₇₅ (15 kg P₂O₅ ha⁻¹) to P₁₀₀ (20 kg P₂O₅ ha⁻¹) to sesamum crop recorded statistically on par seed yield of sesamum seed in these high P soils. Application of increased dose of sulphur from 0 to 30 kg ha⁻¹ enhanced the sesamum seed yield by 10.07% significantly from 10.23 to 11.26 g pot⁻¹. Subsequent enhancement of S application from S₂₀ to S₃₀ kg ha⁻¹ to sesamum crop did not affect the yield of sesamum seed and they were statistically on par (11.26 and 11.35 g pot⁻¹) in these high P soils. The interaction effect of P X S on sesamum seed yield was found to be significant. Maximum seed yield of 12.02 g pot⁻¹ was recorded due to combination of P₁₀₀ (20 kg P₂O₅ ha⁻¹) + S₃₀ (30 kg S ha⁻¹) treatment. Similar on par yields are also observed when P₁₀₀S₂₀ and P₇₅ with either S₃₀ or S₂₀. Sesamum seed yield was not significantly affected by two high P soils. The interaction effect of high P soils and doses of phosphorus application was found to be non-significant on seed yield of sesamum. The interaction effect of high P soils and sulphur doses applied to crop on sesamum seed yield was found to be statistically significant. The interaction effect of P, S and two high P soils on sesamum seed yield was found to be statistically significant. However, in S₁ (67.29 kg P₂O₅ ha⁻¹) significant increase in seed yield was recorded with P₇₅ while in S₂ (83.46 kg P₂O₅ ha⁻¹) it was recorded with P₅₀ compared to control. These differences in seed yield were found to be statistically significant.

Stalk yield of sesamum is significantly influenced by application of different levels of phosphorus fertilizer to sesamum crop in high P soils. The overall yield of sesamum stalk at

different levels of phosphorus application was in range of 19.07 to 21.62 g pot⁻¹ with mean of 20.78 g pot⁻¹. It was found that increasing the phosphorus application from P₀ to P₇₅ (15 kg P₂O₅ ha⁻¹) enhanced the stalk yield significantly from 19.07 to 21.51 g pot⁻¹. Subsequent enhancement of phosphorus application from P₇₅ (15 kg P₂O₅ ha⁻¹) to P₁₀₀ (20 kg P₂O₅ ha⁻¹) to sesamum crop did not effect the stalk yield of sesamum significantly in these high P soils. Application of increased dose of sulphur from S₀ to S₂₀ kg ha⁻¹ enhanced the sesamum stalk yield by 8.19% significantly from 19.63 to 21.24 g pot⁻¹. Subsequent enhancement of sulphur application from S₂₀ to S₃₀ kg ha⁻¹ to sesamum crop did not affect the yield of sesamum stalk and they were statistically on par (21.24 and 21.48 g pot⁻¹) in these high P soils. There was non-significant result in the interaction effect of P X S on sesamum stalk yield. Two high P soils had no significant effect on stalk yield of sesamum. The interaction effect of phosphorus doses and high P soils was found to be statistically non-significant on stalk yield of sesamum. The interaction effect of high P soils and sulphur doses applied to crop on sesamum stalk yield was found to be statistically significant. The interaction effect of P, S and different high P soils on stalk yield of sesamum was found to be statistically significant. However, in S₁ (67.29 kg P₂O₅ ha⁻¹) significant increase in stalk yield was recorded with P₇₅ while in S₂ (83.46 kg P₂O₅ ha⁻¹) it was recorded with P₅₀ compared to control. These differences in seed yield were found to be statistically significant.

Application of different doses of phosphorus to these high P soils resulted response to sesamum seed yield (Table. 3). Application of P₇₅ increased the sesamum seed yield significantly by 13.82% over control. Further increase in dose of phosphorus from P₇₅ to P₁₀₀ to sesamum crop did not influence the yield of sesamum seed significantly in these high P soils. Application of phosphorus fertilizer based on conventional soil test based P recommendation (25% reduction of RDP and 50 % reduction of RDP) to high P soils recorded on par yields with P₁₀₀ in S₁ (67.29 kg P₂O₅ ha⁻¹) and S₂ (83.46 kg P₂O₅ ha⁻¹) (Table.3). It is noticed that the enhancement in seed yield of sesamum was only 0.74% due to application of P₁₀₀ (20 kg P₂O₅ ha⁻¹) over P₇₅ (15 kg P₂O₅ ha⁻¹) in S₁ (67.29 kg P₂O₅ ha⁻¹) and the enhancement in seed yield of sesamum was only 1.57% due to application of P₁₀₀ (20 kg P₂O₅ ha⁻¹) over P₅₀ (10 kg P₂O₅ ha⁻¹) in S₂ (83.46 kg P₂O₅ ha⁻¹) to this high P soils. Based on the above results we can conclude that enhanced biomass accumulation and its efficient translocation from source to sink might have boosted growth parameters and yield contributing characters which in turn paved way to seed yield increase at

elevated fertilizer levels (Harshitha *et al.*, 2022). The above findings are in accordance with Vilakar (2021), Pagal *et al.* (2019) and Verma *et al.* (2014). Reduced P application by 25-50% from the current RDP in high P soils is reported to be a possible P saving measure without scarifying the yields in crops for e.g., in rice, sunflower, rice-rice, rice-sunflower (Babu *et al.*, 2005; Ramya *et al.*, 2015, Srinivas *et al.*, 2017). Thus the present results clearly establish once again that the phosphorus application can be saved in high P soils including the phosphorus loving crop like sesamum. These observations clearly indicate that it is possible to reduce current RDP by 25% to 50% in high P soils without scarifying the yield.

Application of S₂₀ (20 kg S ha⁻¹) increased the seed yield of sesamum by 10.08% over control. Further increase in dose of sulphur from S₂₀ to S₃₀ to sesamum crop didn't influenced the seed yield of sesamum in these high P soils. The bioactivity of sulphur might have played important role in improving yield attributes like capsule per plant, length of capsule and there by seed yield per plant ultimately increase in seed yield (Parmar *et al.*, 2018). Similar findings were reported by Yadav *et al.* (2020), Padasalagi *et al.* (2019).

The combination of P₇₅S₂₀ to sesamum crop results in highest seed yield in high P soils. The significant increase in seed yield per plant with increase in levels of phosphorus and sulphur and their interaction effect may be because of higher nutrient availability and their uptake with each increment level of phosphorus and sulphur which results in favorable condition for more seeds formation. Tekseng *et al.*, (2018). Similar findings were reported by Bhosale *et al.* (2011), Bhavana *et al.*, (2022) and Mishra *et al.*, (2022).

Two high P soils having different native available phosphorus status did not show any significant effect on the seed yield of sesamum (Table. 3).

Stalk yield of sesamum also significantly influenced by application of different doses of phosphorus in high P soils (Table. 4). Application of P₇₅ increased the sesamum stalk yield significantly by 12.75% over control. Further increase in dose of phosphorus from P₇₅ to P₁₀₀ to sesamum crop did not influence the stalk yield of sesamum crop significantly in these high P soils. The increment in stalk yield due to P₁₀₀ was only 0.51% with that of P₇₅. Application of phosphorus fertilizer based on conventional soil test based P recommendation (25% reduction of RDP and 50 % reduction of RDP) to high P soils recorded on par stalk yields with P₁₀₀ in S₁

(67.29 kg P₂O₅ ha⁻¹) and S₂ (83.46 kg P₂O₅ ha⁻¹) (Table. 4). It is noticed that the enhancement in stalk yield of sesamum was only 0.53% due to application of P₁₀₀ (20 kg P₂O₅ ha⁻¹) over P₇₅ (15 kg P₂O₅ ha⁻¹) in S₁ (67.29 kg P₂O₅ ha⁻¹) and the enhancement in stalk yield of sesamum was only 1.80% due to application of P₁₀₀ (20 kg P₂O₅ ha⁻¹) over P₅₀ (10 kg P₂O₅ ha⁻¹) in S₂ (83.46 kg P₂O₅ ha⁻¹) to this high P soils. These observations clearly indicate that it is possible to reduce current RDP by 25 to 50% in high P soils. Improvement in stalk yields with the application of phosphorus doses was reported by Dhage *et al.*, (2014) in soybean, Madhavi *et al.*, (2015) in sunflower and Pagal *et al* (2017) in sesamum.

Application of different doses of sulphur fertilizer influenced the stalk yield of sesamum crop significantly. It is noticed that the application of S₂₀ (20 kg S ha⁻¹) recorded higher stalk yield by 8.19% over control. Further increased dose of sulphur from S₂₀ to S₃₀ didn't showed any significant effect on stalk yield of sesamum in these high P soils. The above findings clearly indicate that sulphur might have played a crucial role in enhancing the stalk yield by its role in physiologically improved dry matter accumulation further led to hiking the stalk yield. (Kumar and Singh, 2021). As for the impact of sulphur fertilization on sesamum stalk yield, our results are also in accordance with Kalyani *et al.* (2020) and Ahmad *et al.* (2018). The interaction effect of P X S on sesamum stalk yield was found to be non-significant. Two high P soils having different native available phosphorus status did not show any significant effect on the stalk yield of sesamum (Table. 4).

Conclusion

Combination of P₇₅S₂₀ is sufficient to obtain higher yield in S₁ (67.29 kg P₂O₅ ha⁻¹). While combined application of P₅₀S₂₀ is sufficient to obtain higher yield in S₂ (83.46 kg P₂O₅ ha⁻¹). The extent of saving of recommended dose of phosphorus fertilizer is 75-100% and 50-100% in soil 1 (67.29 kg P₂O₅ ha⁻¹) and soil 2 (83.46 kg P₂O₅ ha⁻¹) respectively.

UNDER PEER REVIEW

Table 3. Effect of different levels of P and S on seed yield (g pot⁻¹) of sesamum crop in high P soils

3 Factor Tables													
	Soil 1(67.29 kg P ₂ O ₅ ha ⁻¹)					Soil 2(83.46 kg P ₂ O ₅ ha ⁻¹)							
P levels	S ₀ (Control)	S ₁₀ (10 kg S ha ⁻¹)	S ₂₀ (20 kg S ha ⁻¹)	S ₃₀ (30 kg S ha ⁻¹)	Mean	S ₀ (Control)	S ₁₀ (10 kg S ha ⁻¹)	S ₂₀ (20 kg S ha ⁻¹)	S ₃₀ (30 kg S ha ⁻¹)	Mean	Grand Mean		
P ₀ (Control)	9.37	9.95	10.21	10.27	9.95	9.50	10.12	10.26	10.37	10.06	10.01		
P ₂₅ (5 kg P ₂ O ₅ ha ⁻¹)	10.15	10.57	10.70	10.80	10.55	10.01	10.75	10.82	10.95	10.63	10.59		
P ₅₀ (10 kg P ₂ O ₅ ha ⁻¹)	10.55	10.81	10.93	11.16	10.86	10.38	10.94	11.92	11.98	11.31	11.08		
P ₇₅ (15 kg P ₂ O ₅ ha ⁻¹)	10.62	10.95	11.89	11.94	11.35	10.59	11.14	11.96	12.00	11.42	11.39		
P ₁₀₀ (20 kg P ₂ O ₅ ha ⁻¹)	10.64	11.13	11.96	12.00	11.43	10.62	11.30	11.99	12.05	11.49	11.46		
Mean	10.26	10.68	11.14	11.23	10.83	10.22	10.85	11.39	11.47	10.98	10.91		
2 Factor Tables													
	S ₀	S ₁₀	S ₂₀	S ₃₀	Mean		Soil 1	Soil 2	Mean		Soil 1	Soil 2	Mean
P ₀	9.44	10.03	10.24	10.32	10.01	P ₀	9.95	10.06	10.01	S ₀	10.26	10.22	10.24
P ₂₅	10.08	10.66	10.76	10.87	10.59	P ₂₅	10.55	10.63	10.59	S ₁₀	10.68	10.85	10.77
P ₅₀	10.47	10.87	11.42	11.57	11.08	P ₅₀	10.86	11.31	11.08	S ₂₀	11.14	11.39	11.26
P ₇₅	10.61	11.05	11.93	11.97	11.39	P ₇₅	11.35	11.42	11.39	S ₃₀	11.23	11.47	11.35
P ₁₀₀	10.62	11.22	11.98	12.02	11.46	P ₁₀₀	11.43	11.49	11.46	Mean	10.83	10.98	10.91
Mean	10.24	10.77	11.26	11.35	10.91	Mean	10.83	10.98	10.91	S	Soil	Soil	S X Soil
	P	S	P X S			P	Soil	P X Soil	SEm(±)	0.095	-		0.090
SEm(±)	0.107	0.095	0.156			SEm(±)	0.107	-	-	CD (P=0.05)	0.21	NS	0.20
CD (P=0.05)	0.24	0.21	0.35			CD (P=0.05)	0.24	NS	NS	Soil X P X S:SEd(±): 0.214			CD (P=0.05): 0.48

Table 4. Effect of different levels of P and S on stalk yield (g pot⁻¹) of sesamum crop in high P soils

3 Factor Tables													
	Soil 1(67.29 kg P₂O₅ ha⁻¹)					Soil 2(83.46 kg P₂O₅ ha⁻¹)							
P levels	S₀ (Control)	S₁₀ (10 kg S ha⁻¹)	S₂₀ (20 kg S ha⁻¹)	S₃₀ (30 kg S ha⁻¹)	Mean	S₀ (Control)	S₁₀ (10 kg S ha⁻¹)	S₂₀ (20 kg S ha⁻¹)	S₃₀ (30 kg S ha⁻¹)	Mean	Grand Mean		
P₀ (Control)	16.99	19.01	19.19	20.41	18.90	17.26	19.37	19.91	20.45	19.25	19.07		
P₂₅ (5 kg P₂O₅ ha⁻¹)	19.89	20.13	20.77	20.82	20.40	20.00	20.61	21.10	21.52	20.81	20.60		
P₅₀ (10 kg P₂O₅ ha⁻¹)	19.94	20.63	21.41	21.51	20.87	20.31	21.04	21.89	21.91	21.29	21.08		
P₇₅ (15 kg P₂O₅ ha⁻¹)	20.28	21.54	21.98	22.02	21.45	20.50	21.62	22.04	22.09	21.56	21.51		
P₁₀₀ (20 kg P₂O₅ ha⁻¹)	20.45	21.77	22.01	22.04	21.57	20.68	21.82	22.08	22.11	21.67	21.62		
Mean	19.51	20.62	21.07	21.36	20.64	19.75	20.89	21.40	21.61	20.91	20.78		
2 Factor Tables													
	S₀	S₁₀	S₂₀	S₃₀	Mean		Soil 1	Soil 2	Mean		Soil 1	Soil 2	Mean
P₀	17.13	19.19	19.55	20.43	19.07	P₀	18.90	19.25	19.07	S₀	19.51	19.75	19.63
P₂₅	19.94	20.37	20.94	21.17	20.60	P₂₅	20.40	20.81	20.60	S₁₀	20.62	20.89	20.75
P₅₀	20.12	20.84	21.65	21.71	21.08	P₅₀	20.87	21.29	21.08	S₂₀	21.07	21.40	21.24
P₇₅	20.39	21.58	22.01	22.05	21.51	P₇₅	21.45	21.56	21.51	S₃₀	21.36	21.61	21.49
P₁₀₀	20.57	21.79	22.04	22.07	21.62	P₁₀₀	21.57	21.67	21.62	Mean	20.64	20.91	20.78
Mean	19.63	20.75	21.24	21.49	20.78	Mean	20.64	20.91	20.78		S	Soil	S X Soil
	P	S	P X S				P	Soil	P X Soil	SEm(±)	0.112	-	0.089
SEm(±)	0.134	0.112	-			SEm(±)	0.134	-	-	CD (P=0.05)	0.25	NS	0.20
CD (P=0.05)	0.30	0.25	NS			CD (P=0.05)	0.30	NS	NS	Soil X P X S:SEm(±): 0.143		CD (P=0.05): 0.32	

REFERENCES

1. Ahmad, F., Ahmad, J., Nawaz, H., Abbas, M.W., Shah, M.A., Iqbal, S., Mehmood, Z. and Ali, M. 2018. Influence of sulphur and potassium levels on yield and yield attribute of sesame (*Sesamum indicum* L.). *Research in Agriculture Livestock and Fisheries*. 5(2): 147-150.
2. Babu, S.P., Reddy, V and Sathe, S.A. 2005. Phosphorus requirement and use efficiency by sunflower (*Helianthus annuus* L.) in P- accumulated Vertisols. *Journal of Oilseeds Research*. 22 (2):410-413.
3. Bhavana, T., Shankar, T., Maitra, S., Sairam, M. and Kumar, P.P. 2022. Impact of phosphorus and sulphur levels on growth and productivity of summer sesame. *Crop Research*. 57(3): 178-184.
4. Bhosale, N.D., Dabhi, B.M., Gaikwad, V.P. and Agarwal, M.C. 2011. Influence of potash and sulphur levels on yield, quality and economics of sesamum (*Sesamum indicum* L.). *International Journal of Plant Sciences* (Muzaffarnagar). 6(2): pp.335-337.
5. Bouyoucos, G.J. 1962. Hydrometer method improved for making particle size analysis of soils. *Agronomy Journal*.54:464-465.
6. Chaudhary, S.K and Thakur, R.B. 2007. Rational use of phosphorus in a rice-wheat cropping system in Bihar plains. *ORYZA*. 44 (3): 282-284.
7. Dhage, S.J., Patil, V.D. and Dhamak, A.L. 2014. Influence of phosphorus and sulphur levels on nodulation, growth parameters and yield of soybean (*Glycine max* L.) grown on Vertisol. *Asian Journal of Soil Science*. 9(2): 244-249.
8. Food and Agriculture Organization Statistical data bases (FAOSTAT, 2020).
9. Harshitha, C.H., Ramulu, C.H., Reddy, R.U. and Nagabhushanam, U. 2022. Effect of application of 100 and 150% RDF of HDPS cotton and varied RDF of sesame on yield attributes and yield of summer sesame on Alfisols. *The Pharma Innovation Journal*. SP-11(8): 1899-1902.
10. Jackson, M.L. 1973. Soil chemical analysis. Prentice Hall India Pvt. Ltd., New Delhi. 498.
11. Kalegore, N. K., Kirde, G. D., Bhusari, S. A., Kastle, S. V. and Shelke, R. I. (2018). Effect of different level of phosphorus and sulphur on growth and yield attributes of sesame. *International Journal of Economic Plants*. 5:163-66.

12. Kalyani, R., Nagavani, A.V., Chandrika, V., Prasanthi, A and Sagar, G.K. 2020. Performance of sesame (*Sesamum indicum* L.) varieties under varied levels of sulphur. *Andhra Pradesh Journal of Agricultural Sciences*. 6(2): 101-105.
13. Kumar, K.S. and Singh, S. 2021. Effect of different levels of sulphur and spacing on growth and yield of zaid sesame (*Sesame indicum* L.). *The Pharma Innovation Journal*. 10(10): 96-100.
14. Madhavi, A., Babu, P.S. and Reddy, P.V., 2015. Requirement of phosphorus and its use efficiency by sunflower in high phosphorus soils. *Journal of Progressive Agriculture*. 6(1): 27-30.
15. Mishra, A., Mishra, U.S., Sirothia, P. and Singh, O.K. 2022. Evaluation of interaction effect of phosphorus and sulphur on growth, yield and yield components of sesame (*Sesamum indicum* L.) under rainfed condition of Chitrakoot area. *The Pharma Innovation Journal*. 11(1): 1866-1870.
16. Muhr, R. Gilbert, Dutta, N. P.B., Sankara Subramoney, H., Dever, R. F., Laley, V. K and Donahue, R. L. 1965. Soil testing in India (United States Agency for International Development Mission to India, New Delhi).
17. Olsen, S. R., Cole, C. W., Watanable, F. S and Dean L. A. 1954. Estimation of Available Phosphorus in Soils by Extraction with Sodium Bicarbonate. United States Department Agricultural Circular No. 639.
18. Padasalagi, R.M., Lalitha, B.S., Jayadeva, H.M. and Raddy, G. 2019. Effect of sulphur and boron on growth and yield of sesame (*Sesamum indicum* L.). *Journal of Pharmacognosy and Phytochemistry*. 8(6): 1426-1431.
19. Pagal, A.K., Singh, A.P., Behera, S. and Meher, C. 2017. Effect of different level of sulphur and phosphorus on growth and yield attributes of sesamum. *International Journal of Current Microbiology and Applied Sciences*. 6(11): 3278-3285.
20. Parmar, N.N., Patel, A.P. and Choudhary, M. 2018. Effect of sources and levels of sulphur on growth, yield and quality of summer sesame under south Gujarat condition (*Sesamum indicum* L.). *International Journal of Current Microbiology and Applied Sciences*. 7(2): 2600-05.
21. Ramya, V.S., Madhavi, A., Babu, P.S and Madhavi, K. 2015. Zinc requirement to rice in high P soil with sufficient zinc status. *The Journal of Research PJTSAU*. 43(3): 33-35.

22. Shankarlingappa, B.C., Shivaraj, B and Viswanatha, P. 2000. Interaction effect of phosphorous and sulphur on uptake of nitrogen, phosphorous, potassium and sulphur by cowpea. *Karnataka Journal of Agricultural Science*. 13 (2): 295-298.
23. Shelke, R.I., Kalegore, N.K. and Wayase, K.P., 2014. Effect of levels of phosphorus and sulphur on growth, yield and quality of sesame (*Sesamum indicum* L.). *World Journal of Agricultural Sciences*. 10(3): 108-111.
24. Srinivas, A., Surendra Babu, P., Madhavi, A., Vidyasagar, G.E.C.H and Maruthi Sankar, G.R. 2016. Effect of phosphorus application on uptake, yield and fertilizer use efficiency under rice-rice and rice-sunflower systems in high P soils. *Agropedology*. 26 (02): 132-148.
25. Subbaiah, B. V and Asija, G. L. 1956. a. Rapid Procedure for the Determination of available nitrogen in Soil. *Current Science*. 25: 226-230.
26. Surendra Babu, P., Patnaik, M.C and Shankaraiah, M. 2015. Annual report of AICRP on micro and secondary nutrients and pollutant elements in soils and plants, *PJTSAU*, Hyd. 1-119.
27. SYB-2021. Statistical Year Book of Telangana, 2021.
28. Tekseng, O., Singh, P.K. and Bier, K. 2018. Effect of phosphorus and sulphur nutrition on growth, yield and quality attributes of sesame (*Sesamum indicum* L.) under acidic soil of Nagaland. *Journal of Pharmacognosy and Phytochemistry*. 7(1S): 315-319.
29. Verma, R.K., Yadav, S.S., Puniya, M.M., Yadav, L.R., Yadav, B.L. and Shivran, A.C. 2014. Effect of phosphorus and sulphur fertilization on growth and yield of sesame (*Sesamum indicum* L.) under loamy sand soils of Rajasthan. *Annals Agriculture Research New Series*. Vol 35(1):65-70.
30. Vilakar.K. 2021. Standardisation of fertilizer doses for summer sesame in northern telangana zone. M. Sc. thesis. PJTSAU, Telangana.
31. Walkey A and Black C A. 1934. a Proposed Modifications of the Chromic Acid Titration Method. *Soil Science*. 37: 29-38.
32. Williams, C. H and Steinbergs, A. 1959. Soil Sulphur Fraction as Chemical Indices of Available Sulphur in Some Australian Soils. *Australian Journal of Agricultural Research*. 10: 340-352.
33. www.Indiastat.com 2020-2021.

34. Yadav, P., Yadav, S.S., Garg, K. and Yadav, S. 2020. Effect of sulphur and zinc fertilization on growth, yield attributes and grain yield of sesame (*Sesamum indicum*). *Indian Journal of Agronomy*. 65(1): 120-124.

UNDER PEER REVIEW