

# **Original Research Article**

## **Study on Shifting Cultivation Cycle (Jhum Cycle) in Mokokchung District of Northeast India Using Multi Temporal and Land-Sat TM Satellite Data**

---

### **ABSTRACT**

Shifting cultivation has a long history that predates the development of agriculture as a whole. Locally known as jhum farming in Northeast India, this system's farmers are referred to as jhumias. The strategy is based on crop planting during the wet season and cutting and burning vegetation during the dry season. After growing crops for two to three years, the land is left fallow for a number of years before being further cultivated. The locations and area under various jhum cycles are currently inadequate. This was the first attempt to use GIS and remote sensing tools to generate data on various shifting cultivation fallow cycle periods in Mokokchung district of Northeast India between 1988 and 2013. To locate moving farmed areas, IRS and LISS-III sensor data from many time periods were used. While for the years 1988, 1991, 1993, 1994, 1996, 1997, 1999, and 2000, Land-sat TM satellite imagery was used. ERDAS envision and Arc GIS 10.0 software tool were used to determine the area under different jhum cycles, namely 20, 10, and 5 years, and to create the map. For the purpose of determining the spatial extent of the current shifting agricultural areas over the course of 20 years, visual interpretation of the images has been carried out and verified with field checking. Our findings indicate that the Northeast Indian district of Mokokchung's jhum lands declined from 8.99% to 6.31% of its overall geographic area. The highest jhum area in the research region was likewise estimated to be under 20 years cycle (817.23 ha), followed by 5 years cycle (783.02 ha), 10 years cycle (591.9 ha), and 15 years cycle (684.55 ha). A correct fallow cycle of shifting agriculture may help to increase watershed conservation while maximizing crop productivity, according to the fundamental concept of our work.

*Keywords: Shifting cultivation, jhum, fallow cycle, Northeast India.*

### **1. INTRODUCTION**

The Shifting Cultivation is also known as "slash and burn" and "bush fallow" agriculture in different parts of the world. It is the most common type of agriculture in North East India [1], which is locally known as jhum farming and the cultivators are known as jhumias [2]. Shifting agriculture is widely practiced in the tropical uplands, encompassing around 280 million hectares of land and likely supporting 200-300 million people [3]. Over 4,40,000 rural families in Northeast India rely on jhum agriculture for a portion or all of their income. [4] The approach entails clearing forest vegetation from a specified plot by cutting and burning, cultivating the area for one or two years, and then abandoning the field as fallow to regain soil fertility through natural vegetation regeneration. Though this farming system is being considered to be major factor responsible for loss of biodiversity and causing imbalance in the ecosystem, but the fact remains that the majority of the world mega biodiversity area coincide with the area occupied by the indigenous people practicing similar system of agriculture [5]. The role of jhum farming is being widely recognized for ecologically sustainable and economically viable form of agriculture [6]. Shifting cultivation systems are

ecologically viable as long as there is enough land for long (10–20 years) restorative fallow, and expectations of crop yield [7]. The length of this fallow period varies considerably and 5–20 years is common. In Northeast India shifting cultivation cycle of 10–15 years over the years has shortened to about 2–3 years, thus leading to ecological imbalance [8] Fallow duration and cultivation periodicity may be influenced by multiple factors, including ecological factors such as precipitation, soil conditions and topography as well as socio-economic factors [9]. Fallow times are severely reduced as populations increase, and the system deteriorate, leading in severe soil erosion and a decline in soil fertility and productivity. A reduction in the fallow cycle lowers the soil fertility and yield, increases soil erosion, and causes watershed siltation [10]. In Northeast India, the dynamics of crop-fallow rotation cycles in shifting cultivation fallows, however, are poorly understood [11]. The most preferred jhum fallows in the region remains 7–11 years [12] against most believed that jhum cycle has drastically reduced. The vulnerability assessment could offer us with a comprehensive examination of the state of ecological sustainability as a result of shifting cultivation at the landscape level. The length of shifting cultivation cycles can be used to determine vulnerability. The length of a shifting cultivation cycle appears as alternating fallow and cultivation seasons. This premise is widely accepted; nonetheless, empirical data of areas in various fallow cycles is limited. The primary goal of our research is to identify the places in Northeast India that are under distinct fallow cycles. As a result, fallow length could be a powerful predictor of crop productivity and the greatest proxy for measuring system sustainability. As a result, it is vital to bridge this gap by creating a fallow cycle map for this specific region to aid in its plans and management. The study is thus carried out with the goals of i) estimating the areas under different fallow cycles and ii) creating a fallow cycle map for the specific region.

## **2. MATERIAL AND METHODS**

### **2.1 STUDY AREA**

Mokokchung district of Nagaland covers an area of 1,615 sq km which falls under humid sub-tropical climate. It is situated at 94° 30.8078' E longitudes and 26°19.3222' N latitude (Fig.1). It is located at an elevation of 1325 meters above sea level. The annual average rainfall of the district is 250 cm. The maximum rainfall occurs during the months of June and July. Rainfall generally begins from April and continues till the end of September. The area enjoys a cold winter and mild summer. January and February are the coldest months when the night temperature comes down to 2°C. In summer it is not at all hot; rather it is cold in comparison to the adjoining plains of Assam. The average temperature during summer is 27°C and the temperature does not rise beyond 32°C. Soils found in Mokokchung district mainly include alluvial soil, non laterite red soil and forest soil which are in acidic in nature. The district covers 188.51 square kilometer area under agriculture and horticulture. The main agricultural products of the district are rice, maize, tomatoes, passion fruit, and oranges. The primary method of cultivation is jhum farming and about 70% of the total population is engaged in agriculture only.





Fig.1. Study sites of *jhum* cycle in Mokokchung district of Nagaland, Northeast India.

## 2.2 Integrating Remote Sensing, Ground Based Measurements and GIS for Calculating Jhum Cycle


Multi temporal satellite imagery of IRS (Indian Remote Sensing Satellite) LISS-III (Linear Imaging Self Scanning) sensor for the year 2003, 2006, 2009 and 2012 were used to identify shifting cultivated areas of Mokokchung district of Nagaland and observed the changes occurred in those areas. Whereas, Land-sat TM satellite imagery were used for the year 1991, 1993, 1994, 1996, 1997, 1999 and 2000 for the study area. Visual image interpretation technique has been used to identify the shifting cultivated areas and changes occurred during 20 years. The data (images) have been collected from NRSC, Hyderabad and the study area Digital processing of the satellite data pertaining to study area during March-April has been done using ERDAS IMAGINE 9.2 image processing software. The standard false colour composite (FCC) was generated by assigning blue, green and red colors to visible green, visible red and near infrared bands on 1:50,000 scale, respectively. Visual interpretation of the imagery has been done and verified with field checking for deriving information on spatial extent of current shifting cultivation areas. All the input vector layers (current shifting cultivation layers) were overlaid in GIS environment and change analysis was done using analysis tools. Analysis tool option is used from Arc tool box window to

overlay the layers for calculate the intersect area of jhum. The intersect areas are only considered by erasing repeated jhum areas within cycle. The area under different jhum cycle viz 20, 10 and 5 years were calculated by ERDAS imagines and Arc GIS 10.0 software tool and developed the maps. The brief methodology is presented in the flow chart (Fig.2).


UNDER PEER REVIEW




Overlay of layers



GIS analysis



GIS analysis



Estimation of fallow cycle

GIS analysis




Fig.2. Flow chart of Methodology for estimation of area under jhum cycle (Mokokchung district)

### 3. RESULTS AND DISCUSSION

Shifting cultivation is the practice of removing a section of forest by cutting it down and burning it, then farming the cleared area for a year or longer before moving to another section. After cropping, each patch is given a number of years to transition back to secondary growth before being cleaned and re-cultivated.

Jhum cycle is the total number of years under agriculture plus the number of years in fallow. The cycle may last only three to five years or up to twenty years. The jhum cycle varies in Northeast India, ranging from 3 to more than 20 years. The information in table 1 demonstrates that the Mokokchung district's total current jhum area varies from year to year and has been on the decline since 1991. In the years 1991 and 2012, respectively, the Mokokchung district's total jhum area decreased from 8.99% to 6.31% of the district's overall geographic area. The highest area (14519.54 ha) and lowest area (7337.81 ha) were discovered in 1991 and 2003, respectively. The results showed that in the Mokokchung district of Nagaland, Northeast India, the maximum jhum area is under 20 years cycle (817.23 ha), followed by 5 years cycle (783.02 ha), 10 years cycle (591.9 ha), and 15 years cycle (684.55 ha) (Table 1).

Several researchers also documented a decrease in jhum area and cycle. Several socioeconomic factors may have contributed to the shrinkage of the jhum region [12,13,14,15,16]. The cutting and burning of vegetation is a labor-intensive part of jhum farming [17]. The entire farming process is rain-fed, which increases the likelihood of crop failure. Jhum land generally has relatively low production, especially when the fallow period is shorter [18]. Additionally, the younger generation has a propensity to move away from commercial areas. Moreover, the government and non-governmental organizations pressured the jhumias towards settled farming. According to [19] where instantaneous carrying capacity has been attained in that ecosystem, a bigger population is to blame for the shorter fallow period under jhum farming. The highest theoretical population that can be sustained in a given region with the current technological and ecological conditions of production is known as the instantaneous carrying capacity. However, hill farmers continue to rely on it for their livelihoods, either entirely or partially. With rice, vegetables, cash crops, fruits, spices, etc., it provides staple foods for the majority of the months. This could be the

explanation for why jhum farming is regarded as one of the greatest livelihood options in mountainous areas. A sound approach is therefore needed to maintain and enhance the jhum farmer. Many workers come to the conclusion that a longer fallow cycle is crucial for maintaining the farming's environment, but we lack information on the locations that experience various fallow cycles. In regions where jhum farming is still practiced, the findings of total areas under various cycles may be valuable to policy makers for sustainable development.

Table 1. Total area under different jhum cycle in Mokokchung district of Nagaland

Year	Total <i>jhum</i> area (ha)	5 years cycle		10 years cycle		15 years cycle		20 years cycle	
		Year	Area (ha)	Year	Area (ha)	Year	Area (ha)	Year	Area (ha)
1991	14519.54	1996- 2000	739.02 (4.75)	1991- 2000	3407.5 (21.90)	1991- 2006	756.23 (7.94)	1991 to 2009- 10	814.11 (6.58)
1993	12819.31	2000- 01 to 2005- 06	555.33 (5.83)	1993- 2003	1488.6 (20.29)	1993- 2009	684.55 (5.53)	1993- 2012	817.23 (8.01)
1996	12617.23	2003- 04 to 2009	175.05 (1.42)	1996- 2006	874.88 (9.18)	-	-	-	-
2000	15561.63	2006- 07 to 2012	783.2 (7.68)	2003- 2012	591.9 (5.80)	-	-	-	-
2003	7337.81	-	-	-	-	-	-	-	-
2006	9523.9	-	-	-	-	-	-	-	-
2009	12363.39	-	-	-	-	-	-	-	-
2012	10190.12	-	-	-	-	-	-	-	-

Figures in parentheses indicates is % of total *jhum* area; - = Not calculated / no data

### 3.1 Mapping of jhum fields over the years

For identifying and defining of jhum fields from the multi-temporal Landsat data, LISS-III, and Land-sat TM Imagery of 1991 to 2012, the visual interpretation and on-screen digitization approaches were used. With the help of time series analysis of satellite data, mapping techniques, and overlaying, the jhum field's fallow age was ascertained (Fig. 3).




Fig.3. Maps depicting the fallow cycle over 5, 10, 15, and 20 years in Nagaland, Northeast India.

#### 4. CONCLUSION

Jhum, or shifting farming, is a key economic activity of the humid tropics in India's northeastern areas. Jhum is environmentally friendly as long as the fallow cycle is long enough to replenish the forest and soil nutrients lost during agriculture. According to our findings, the overall jhum area in Nagaland's Mokokchung district has been declining since 1991. However, the highest area of jhum cultivation in Mokokchung district is under a 20-year cycle, which is a good sign for jhum forest eco-restoration. The estimated jhum cycle data may be useful for good strategic jhum farming management in the region.

## REFERENCES

1. Upadhaya K, Barik SK, Kharbhih VM. Traditional Bun Shifting Cultivation Practice in Meghalaya, Northeast India. *Energy, Ecology and Environment*. 2020: 5:34–46.
2. Deka PK, Sarmah D. Shifting Cultivation and Its Effects in Regarding of Perspective in Northern India. *International Journal of Business and Management*. 2010: 3(2):157-165.
3. Nath AJ, Reang D, Sileshi GW. The Shifting Cultivation Juggernaut: An Attribution Problem. *Global Challenges*. 2022:6: 2200051.
4. NEC. Basic Statistics of Northeastern Region. Government of India, Published by Northeastern Council Secretariat (Evaluation and Monitoring Section), Nongrim Hills, Shillong, Meghalaya, India. 2015: <https://necouncil.gov.in>.
5. Kalita HM, Baruah MS, Datta D, Jini D, Alone RA. Status of Shifting Cultivation (Jhum) in Arunachal Pradesh, India. *Jhum Improvement for Sustaining Farm Livelihood and Natural Resource Conservation in Northeastern Hill Region: visit as and frontiers*; ICAR Research Complex for NEH Region publication, Umium, Meghalaya. 2017: 1-2.
6. FAO. The State of Food and Agriculture: Innovation in family farming. Food and Agricultural Organization of the United Nation Rome. 2014: <https://www.fao.org/3/i4040e/i4040e.pdf>
7. Lal R. Shifting cultivation. *Encyclopedia of Soils in the Environment, Science Direct*, Columbia University Press, New York, NY, USA. 2005:207.
8. Ranjan R, Upadhyay VP. Ecological Problems Due to Shifting Cultivation. *Current Science*. 1999:77(10):1246.
9. Mertz O. The Relationship between Length of Fallow and Crop Yields in Shifting Cultivation: A Rethinking. *Agroforestry Systems*. 2002:2:149-159.
10. Ziegler AD, Bruun TB, Guardiola-Claramonte M, Giambelluca TW, Lawrence D, Lam NT. Environmental Consequences of the Demise in Swidden Cultivation in Montane Mainland Southeast Asia: Hydrology and Geomorphology. *Human Ecology*. 2009:37:361–373.
11. Thong P, Pebam R, Sahoo UK. A Geospatial Approach to Understand the Dynamics of Shifting Cultivation in Champhai District of Mizoram, North-East India. *Journal of the Indian Society of Remote Sensing*. 2018:46:1713–1729.
12. Thong P, Sahoo UK, Pebam R. Spatial and Temporal Dynamics of Shifting Cultivation in Manipur, Northeast India Based on Time-Series Satellite Data. *Remote Sensing Applications: Society and Environment*. 2019:14:126-137.
13. Toky OP, Ramakrishnan PS. Cropping and Yields in Agricultural Systems of the North-Eastern hill Region of India. *Agriculture, Ecosystem & Environment*. 1981:7:11–25.

14. Roder W, Phengchanh S, Keoboulapha B. Relationships between Soil, Fallow Period, Weeds and Rice Yield in Slash and Burn Systems of Laos. *Plant and Soil*. 1995;176:27–36.
15. Silva-Forsberg MC, Fearnside PM. Brazilian Amazonian Caboclo Agriculture: Effect of Fallow Period on Maize Yield. *Forest Ecology and Management*. 1997;97:283–291.
16. Kato MSA, Kato OR, Denich M, Vlek PLG. Fire Free Alternatives to Slash-and-Burn for Shifting Cultivation in the Eastern Amazon Region: The Role of Fertilizers. *Field Crops Research*. 1999;62:225–237.
17. Datta J, Gangadharappa NR, Debnath A. Sustainability of Jhum Cultivation as Perceived by the Tribal People of Tripura. *International Journal of Social Science*. 2014;3:179-190.
18. Zonayet, Paul. Study on Productivity of Jhum Crops and Post-harvest Soil Nutrient Status by Using NPK Briquette. *International Journal of Bio-resource and Stress Management*. 2020;11(4):361-369.
19. Vermeer DE. Population Pressure and Crop Rotational Changes among the Tiv of Nigeria. *Annals of American Association of Geographers*. 2005;60:299-314.