

*Original Research Article*

**Exploring the Relationship Between Health Information Technology and Use of Prescribed Antihypertensive Medications**

Abstract

**Objective:** Our objective was to assess the correlation between various internet uses for health-related purposes and the utilization of prescribed antihypertensive medications. Additionally, we aimed to explore how socioeconomic status influences this relationship.

**Method:** This study was a cross-sectional analysis of 8,224 participants, representing 69,033,231 adults in the United States with hypertension who were prescribed antihypertensive medications. Out of these respondents, 7,837 individuals (88.8%) reported adherence to their prescribed medication regimen. The study examined several independent variables, including internet usage for (1) accessing health information, (2) filling prescriptions, (3) scheduling medical appointments, and (4) communicating with healthcare providers via email. The dependent variable under investigation was the usage of antihypertensive medications following prescription. To analyze the relationship between internet use for health-related purposes and the adherence to prescribed antihypertensive medications, as well as the influence of socioeconomic status on this relationship, a multiple logistic regression was employed. By utilizing this statistical approach, the researchers could assess the associations between the variables while avoiding potential issues of plagiarism.

**Result:** After controlling for other factors, individuals who utilized the internet for refilling prescription medications were found to have 1.65 times higher odds (95% CI 1.26, 2.16,  $p < .001$ ) of taking prescribed antihypertensive medications compared to those who did not use the internet for this purpose. Among hypertensive individuals who used the internet for prescription refills, specific subgroups showed even higher odds of taking prescribed antihypertensive medications. Specifically, those who were employed had an adjusted odds ratio (AOR) of 2.04 (95% CI 1.39, 2.99,  $p < .001$ ), college graduates had an AOR of 1.86 (95% CI 1.14, 3.04,  $p = .013$ ), and individuals earning  $\geq$  \$20,000/year had an AOR of 2.74 (95% CI 1.68, 4.46,  $p < .001$ ) compared to their counterparts who were unemployed, non-college graduates, or earned less than \$20,000/year.

**Conclusion:** The study suggests a potential association between online prescription refill and adherence to antihypertensive medications, with this relationship being particularly pronounced among individuals

with higher socioeconomic status. Further research is warranted to explore the connection between health-related internet usage and medication adherence.

Keywords: Health, information, technology, antihypertensives, adherence

## Introduction

Hypertension is a well-known cause of morbidity in the United States (US), with a prevalence of 45.4%, contributing to approximately 270,000 per year hypertension-related cardiovascular diseases according to a 2018 report by the Center for Disease Control (CDC)[<sup>1</sup>]. Given the high prevalence of hypertension-related complications, effective measures to manage hypertension remain of utmost importance.

Traditionally, the management of hypertension involves the use of lifestyle modifications and antihypertensive medications, measures that have been shown to effectively manage hypertension and as a result, reduce its complications. Despite these approaches, the prevalence of hypertension and hypertension-related cardiovascular diseases remain high, with challenges like antihypertensive medication non-adherence contributing significantly to it [<sup>2-4</sup>]. There have been several reported reasons for non-adherence to antihypertensive medications, some include poor patient-clinician communication, low socioeconomic factors, high pill burden, and poor access to health care [<sup>5-8</sup>]. Although different interventions that address these factors have shown encouraging results, newer approaches may yet improve adherence to antihypertensive medications, one of which is the use of health information technology.

Different health technological tools have been studied in recent times, all of which have the common theme of using health care information to make decisions that are expected to improve the quality of health care delivery[<sup>9</sup>]. Some of these tools include mobile applications, text messaging, and advanced functions of electronic health records. Reports of their efficacy in improving the quality of health care has been mixed and issues about accessibility of these tools remains of huge concern given the existing potential of worsening the disparities in health. In the Hypertension Adherence Program in Pharmacy

(HAPPy) trial, the use of prescription reminders via short message service (SMS) and telephone calls in addition to other interventions resulted in a 22.6% increase in adherence to antihypertensive medications [10]. In the study of a different technological approaches to improve medication adherence among asthmatics, the use of interactive voice recognition software resulted in a modest increase in inhaled corticosteroid (ICS) therapy adherence [11]. Contrastingly, other studies have reported negative findings, with two studies reporting that the use of the internet as a source of information led to poor adherence to medications and vaccines due to negative feelings users experienced from the information they found online [12,13].

**With regards to the accessibility of HIT and its impact on the overall quality of health care.** Poor access to HIT has been linked to low electronic health literacy, which in itself is associated with a poor socioeconomic status [14,15]. The implication of this is that individuals in a low socioeconomic class may benefit less from this technological intervention. This challenge echoes one of the concerns from experts of the Health Information Technology for Economic and Clinical Health (HITECH) Act, that the quality of health care may not improve as much as intended with the use of electronic health records with advance functions such as online medication refill requests, communication tools and appointment scheduling [16]. Given these benefits and concern, we set out to analyze the relationship between health information technology use, in this case the use of the internet for health purposes and use of prescribed antihypertensive medications, as well as the impact of socioeconomic status on this relationship. Our hypothesis is that the use health information technology will be associated with a positive effect on antihypertensive medication adherence.

## Method

### **Study design**

This is a cross-sectional study utilizing the sample adult core data of the 2018 National Health Interview Survey (NHIS). The NHIS is a survey of non-institutionalized civilian population of the United States (US) conducted by the Center for Disease Control and Prevention. The data contained sociodemographic,

health information and medical condition variables. Further details of this survey data are available elsewhere [17]. Since the data was publicly available, an IRB review was not required.

### **Study population**

We identified 8,224 respondents representing 69,033,231 US adults with hypertension from the NHIS who answered yes to the question “*Was any medicine ever prescribed by a doctor for your high blood pressure?*” Among them 7,837/8224 (88.8%) respondents representing 610,000,000 adults answered yes to the questions “*Are you now taking any medicine prescribed by a doctor for your high blood pressure?*” These individuals who answered yes to taking antihypertensive medication were classified as being adherent to antihypertensive medications.

### **Measures**

#### *Sociodemographic variables*

Age (18 -39, 40 – 64 and  $\geq$  65 years), race (white, black, Asians, others), gender (female and male), marital status (married, formerly married, and never married), smoking status (current smokers, former smokers, and never smoked) and body mass index (underweight, normal, overweight, and obese).

#### *Socioeconomic factors*

Income category ( $<$ \$20,000/year and  $\geq$  \$20,000/year), employment status (employed and unemployed) and educational status (did not graduate high school, graduated high school, graduated college).

#### *Independent variables*

Health information technology use: use of the internet to (1) fill up prescription (2) communicate with a health care provider (3) look up health information and (4) schedule medical appointment.

#### *Dependent variable*

Adherence to or taking prescribed antihypertensive medications.

## Statistical analysis

A sample weight was applied during the data analysis to ensure generalizability of the results. A descriptive statistic was used to assess the sociodemographic proportion of participants who use HIT in different forms and to analyze the distribution of different uses of HIT by taking prescribed antihypertensive medication status. A multiple logistics regression model was used to analyze the relationship between different uses of HIT and taking prescribed antihypertensive medications with confounders adjusted for and a second model was used to analyze the interaction between socioeconomic status and HIT on antihypertensive medication use. The analysis was performed using STATA 14.0 with a p-value of  $<.05$ .

## Results

Table 1 below shows the sociodemographic distribution of the study participants by different forms of health information use. With regards to all uses of HIT, respondents who use the various forms were mostly 40 – 64 years old, white, high school graduates, obese, employed, earn  $\geq$  \$20,000/year and have never smoked. The distribution differed only by gender such that respondents who use the internet to fill up prescriptions and communicate with a healthcare provider were mostly males, while those who use the internet to schedule medical appointments and look up health information were mostly females.

Table 1. Sociodemographic distribution of study participants by health information technology use

Variable	Variable Subset	Fill up prescriptions online N (weighted %)	Communicate with a healthcare provider N (weighted %)	Schedule medical appointment N (weighted %)	Look up health information N (weighted %)
<b>Age in years</b>	-	-	-	-	-
-	18 – 39	89 (9.4%)	93 (8.9%)	91 (10.9%)	291 (10.0%)
-	40 – 64	566 (54.7%)	653 (54.3%)	524 (55.0%)	1832 (54.7%)

	≥ 65	479 (35.9%)	546 (36.8%)	393(34.1%)	1592 (35.3%)
<b>Gender</b>	-	-	-	-	-
-	Female	559 (48.6%)	661 (48.9%)	538 (52.3%)	2077 (53.8%)
	Male	575 (51.4%)	631 (51.1%)	470 (47.7%)	1638 (46.2%)
<b>Race</b>	-	-	-	-	-
-	White	963 (84.0%)	1062 (81.7%)	810 (79.1%)	3001 (79.2%)
	Black	116 (10.9%)	145 (11.5%)	134 (13.5%)	461 (13.6%)
	Asians	28 (3.1%)	44 (4.5%)	35 (4.9%)	129 (4.3%)
	Others	27 (2.0%)	41 (2.4%)	29 (2.5%)	124 (2.9%)
<b>Educational level</b>	-	-	-	-	-
-	< high school (HS)	38 (3.5%)	31 (2.3%)	29 (3.5%)	167 (4.6%)
	Graduated HS	584 (52.0%)	629 (49.1%)	499 (50.4%)	2073(55.6%)
	Graduated college	510 (44.5%)	631 (48.6%)	478 (46.1%)	1471 (39.8%)
<b>Marital status</b>	-	-	-	-	-
-	Married	642 (67.5%)	708 (67.0%)	543 (64.6%)	1880 (62.0%)
	Formerly married	305 (18.2%)	378 (18.7%)	283 (19.0%)	1237 (22.7%)
	Never married	185 (14.3%)	205 (14.4%)	181 (16.4%)	592 (15.3%)
<b>BMI</b>	-	-	-	-	-
-	<18.5	1 (0.1%)	5 (0.3%)	5 (0.3%)	17 (0.4%)
	18.5 – 24.9	201 (16.7%)	231 (17.4%)	187 (17.6%)	668 (16.9%)
	25 -29.9	364 (31.7%)	427 (32.9%)	306 (30.5%)	1229 (33.9%)
	≥ 30	568 (51.6%)	629 (49.4%)	510 (51.7%)	1801 (48.9%)
<b>Employment</b>	-	-	-	-	-
	Unemployed	592 (48.5%)	626 (44.0%)	467 (42.8%)	1938 (48.0%)

-	Employed	541 (51.5%)	665 (56.0%)	541 (57.2%)	1775 (52.0%)
<b>Income per year</b>	-	-	-	-	-
-	<\$20,000	96 (14.1%)	104 (12.0%)	75 (11.4%)	334 (15.6%)
	≥ \$20,000	519 (85.9%)	632 (88.0%)	520 (88.6%)	1615 (84.4%)
<b>Smoking status</b>	-	-	-	-	-
-	Never	588 (54.6%)	723 (59.5%)	573 (60.0%)	1981 (55.9%)
	Former	439 (36.4%)	452 (32.2%)	335 (30.3%)	1256 (31.9%)
	Current	107 (9.0%)	117 (8.3%)	100 (9.7%)	447 (12.3%)

Table 2 below shows the proportion of respondents who use different forms of HIT by antihypertensive use status. Among 6349 respondents who reported taking prescribed antihypertensives, 3261 (41.3%) of them use the internet to look up health information, 1151 (15.0%) use the internet to communicate with a health care provider, 1040 (13.7%) use the internet to fill prescriptions and 897 (12.7%) use the internet to schedule medical appointments.

**Table 2. Weighted percentage of respondents who use different forms of HIT antihypertensive medications use status**

Variable	Currently not taking their antihypertensive medications N (weighted percentage)	Taking their antihypertensive medications N (weighted percentage)
Communicate with a health care provider	141 (2.0%)	1151 (15.0%)
Schedule medical appointment	111 (1.5%)	897 (12.7%)
Filled prescription	94 (1.3%)	1040 (13.7%)
Look up health information	454 (6.2%)	3261 (41.3%)

Table 3 below shows different forms of health information technology use and the odds of taking prescribed antihypertensive medications. Those who use the internet to fill up prescription medications as compared to those who do not have 47% higher odds (OR 1.47, 95% CI 1.14,1.90,  $p$  .003) of taking their prescribed antihypertensive medications. This odd further increased to 65% (AOR 1.65, 95% CI 1.26, 2.16,  $p$  <.001) after controlling for covariates. Individuals who use the internet to look up health information had 30% less odds (OR 0.71, 95% CI 0.59, 0.86,  $p$  <.001) of taking their antihypertensive medications as compared to those who do not. However, this was not significant after controlling for covariates (AOR 0.87, 95% CI 0.71, 1.06,  $p$  .164).

**Table 3. Crude and adjusted odds of antihypertensive medication adherence by different uses of health information technology**

Variable	Odds ratio (95% CI)	$p$	Adjusted odds ratio (95% CI)	$p$
Look up health information	0.71 (0.59 – 0.86)	<.001	0.87 (0.71 – 1.06) <sup>a</sup>	.164
Fill out medication prescriptions	1.47 (1.14 – 1.90)	.003	1.65 (1.26 – 2.16) <sup>b</sup>	<.001
Schedule medical appointments	1.07 (0.82 – 1.40)	.625	1.23 (0.93 – 1.65) <sup>c</sup>	.152
Communicate with a healthcare provider	0.93 (0.73 -1.19)	.576	0.99 (0.77 – 1.27) <sup>d</sup>	.950

a Adjusted for age, gender, race, marital status, body mass index and smoking status

b Adjusted for age, race, marital status, body mass index and smoking status

c Adjusted for age, marital status, body mass index and smoking status

d Adjusted for age, race, marital status, and smoking status

Table 4 below shows the effect of the interaction between socioeconomic factors on the relationship between health information technology and taking prescribed antihypertensive medications. Respondents who were employed and use the internet to fill out prescriptions had an insignificant odds (OR 1.13; 95% CI 0.80, 1.59,  $p$  .493) of taking their prescribed antihypertensive medications as compared to those who were unemployed and do not use the internet to fill out prescription. This odd became significant after adjusting for covariates (AOR 2.04; 95% CI 1.39, 2.99,  $p$  <.001).

As compared to respondents who did not graduate HS and do not use the internet to fill prescription, respondents who graduated college and use the internet to fill out prescription had a 67% higher odd (OR

1.67; 95% CI 1.04, 2.69,  $p$  .033) of being adherent to antihypertensive medications, and this odd increased to 86% after adjusting for covariates (AOR 1.86; 95% CI 1.14, 3.04,  $p$  .013).

Respondents who earn  $\geq$  \$20,000/year and use the internet to fill prescription as compared to respondents who earn  $<$  \$20,000/year and do not use the internet to fill prescription had a significant odd (OR 2.68; 95% CI 1.74, 4.12,  $p$   $<$ .001) of taking their antihypertensive medications. This odd was similar after adjusting for covariates (AOR 2.74; 95% CI 1.68, 4.46,  $p$   $<$ .001).

**Table 4. An interaction analysis of socioeconomic status and on-line medications refill on adherence to antihypertensive medications**

Variable	OR (95% CI)	$p$	AOR (95% CI)	$p$
<b>By employment status</b>	-	-	-	-
Respondents who are unemployed and do not use the internet to fill out prescriptions	1.00	-	1.00	-
Respondents who are employed and do not use the internet to fill out prescriptions	0.56 (0.46 – 0.69)	$<$ .001	0.95 (0.76 – 1.19)	.657
Respondents who are unemployed and use the internet to fill out prescriptions	1.12 (0.75 – 1.68)	.589	1.17 (0.77 - 1.77)	.460
Respondents who are employed and use the internet to fill out prescriptions.	1.13 (0.80 – 1.59)	.493	2.04 (1.39 - 2.99)	$<$ .001
<b>By education</b>	-	-	-	-
Respondents who did not graduate HS and do not use the internet to fill prescription	1.00	-	1.00	-
Respondents who graduated HS and do not use the internet to fill prescription	0.68 (0.50 – 0.91)	<b>.010</b>	0.74 (0.55 – 0.99)	<b>.045</b>
Respondents who graduated college and do not use the internet to fill prescription	0.85 (0.61 – 1.18)	.334	0.87 (0.62 -1.22)	.422
Respondents who did not graduate HS and use the internet to fill prescription	0.67 (0.24 - 1.89)	.448	0.77 (0.30 - 2.01)	.596

Respondents who graduated HS and use the internet to fill prescription	0.87 (0.58 – 1.29)	.483	1.09 (0.72 – 1.65)	.694
Respondents who graduated college and use the internet to fill prescription	1.67 (1.04 - 2.69)	<b>.033</b>	1.86 (1.14 – 3.04)	<b>.013</b>
<b>By income category</b>	-	-	-	-
Respondents who earn < \$20,000/year and do not use the internet to fill prescription	1.00	-	1.00	-
Respondents who earn >/\$20,000/ year and do not use the internet to fill prescription	1.42 (1.04 – 1.94)	<b>.027</b>	1.33 (0.92 – 1.94)	<b>.133</b>
Respondents who earn less than < \$20,000/ year and use the internet to fill prescription	1.74 (0.81 – 3.74)	.154	1.50 (0.65 – 3.47)	.343
Respondents who earn >= \$20,000/ year and use the internet to fill prescription	2.68 (1.74 – 4.12)	<b>&lt;.001</b>	2.74 (1.68 – 4.46)	<b>&lt;.001</b>

### Discussion

We set out to analyze the relationship between different forms of the use of the internet for health purposes and the odds of taking prescribed antihypertensive medications and in addition evaluate the interaction effect of socioeconomic status on this relationship. We found out that among the forms of HIT evaluated, the use of internet to fill up prescription was associated with a significant odd of taking prescribed antihypertensive medications. **As compared to respondents who do not use the internet to refill prescription medications and are either unemployed, earn < \$20,000 per year or did not graduate high school, respondents who do not use the internet to refill prescription medications and are either employed, have a higher level of education or earn  $\geq$  \$20,000 per year were not more likely to take prescribed antihypertensive medications.** All three measures of socioeconomic status however had a positive

interaction effect on the relationship between online prescription medication refills and taking prescribed antihypertensive medications. The findings of this study re-emphasizes three of five factors identified by the American College of Preventive Medicine as key in the adherence to prescription medications which are socioeconomic factors, patient-related factors and health care-system factors, with the others being medication condition-related factors, and therapy-related factors [18].

The available evidence suggests that improving access to health care can lead to an increase in medication adherence. In a review of management of adherence to antihypertensive medication, Burnier and Egan reported that the problem of antihypertensive medication adherence can be addressed by easing the access to antihypertensive medications [19]. In our study, as compared those who do not, individuals who use the internet to fill up prescription medications had 40% increase in the odds of taking prescribed antihypertensive medications, with this odd increasing further to 65% after adjusting for confounders. Other studies have reported similar positive associations of online refills and medication adherence. Sarkar et al, reported an absolute decline of 6% of non-adherence in individuals prescribed statins for dyslipidemia [20]. McInnes et al, reported an adherence of more than 90% to antiretroviral medications among veterans who use personal health record functions such as online refills [21], and Lyles et al, reported a significant improvement overtime of statin adherence by individuals who exclusively refill prescriptions online [22]. It is possible that being able to refill antihypertensive medications online creates an ease of access which could serve as an important strategy in improving medication adherence in individuals with hypertension and may extend to other populations. This increased likelihood to be adherent to prescribed medications may also be due to the higher engagement with electronic health record portals online refill of medications provide as compared to the other uses of the internet. This may imply that a higher engagement with the health care system through health portals may lead to an increase in healthy behaviors such as adherence to medications.

Interestingly, there was no difference in taking prescribed antihypertensive medications between those who use HIT in other forms besides online refill of prescription medications and those who did not. We

found no difference in taking prescribed antihypertensive medications with regards to the use of the internet to look up health information. This was not consistent with reports from other studies on use of the internet to gather health information, as findings of a positive [23], as well as a negative association with medication adherence have been reported [24]. This inconsistency was also found between the use of the internet to communicate with a health care provider and antihypertensive medication adherence, as studies have shown that patient-provider communication is associated with a positive medication adherence outcome [25-27]. However, it is important to mention that these studies with a reported positive association evaluated in-person communication as opposed to online communication which we analyzed. Another reason may be related to the less engaging nature of the health care system these other uses provide, and as such healthy habits are less likely to be encouraged with these methods.

Beyond these mixed findings, the interaction analysis showed that as opposed to a lower socioeconomic status, a higher socioeconomic status had a positive effect on the relationship between the use of health information to fill prescription medication and taking prescribed antihypertensive medications. Studies from Wong et al and Paccoud et al reported the positive effect a more affluent socioeconomic status has on the use of health information technology [28,29]. However, just like other studies that have been done, the relationship with medication adherence was not studied. This effect may be related to other factors such as the better health literacy of individuals in higher socioeconomic groups [30] and the higher likelihood of their use of the internet [31,32]. As individuals of a lower social status have been reported to use the internet in less effective and profitable ways [33]. Given these arguments, although the implication of our findings are positive with regards to medication adherence, encouraging the use of HIT to access health care may have a potential of marginalizing minorities.

This study has a few strengths worthy of mention. First, we evaluated the interaction of socioeconomic status on the relationship between a form of HIT use and taking prescribed antihypertensive medications. We felt that this was important as socioeconomic status has been shown to influence access to HIT and as such the relationship between HIT and adherence to antihypertensives may be affected by this influence

as shown in this study. We analyzed different uses of HIT in patient care and felt this was necessary as there are different uses of HIT and patients may use it for different reasons. Sampling weights were applied during the data analysis, which allowed for the generalization of the results to the US population. Despite these strengths, some weaknesses do exist. First, although sampling weights were applied, the proportion of HIT users who were and were not adherent to antihypertensive medications were not uniform, which may have influenced our results. Second, as this is a cross sectional study, causation cannot be implied, and just because people use HIT and are adherent to medications those not necessarily mean that this is the reason why, as there are other factors not controlled for such as the duration of hypertension, relationship with primary care provider, and access to the internet.

## **Conclusion**

Not all forms of health information technology use are associated with an increase in adherence to antihypertensive medications. Communicating with a health care provider online may not have the same effect on medication adherence as in-person communication as shown in other studies. Being able to refill medications online may improve adherence to antihypertensive populations and a higher socioeconomic status may be associated with a positive effect on the relationship between online medication refill and medication adherence. Further studies are needed to evaluate the relationship between different forms of health information technology use and adherence to medications, and to evaluate the potential for marginalization of minorities.

**Data Availability:** The data used in this study was from a publicly available data (NIS)

**Ethical approval:** This Paper does not require ethical approval.

## Disclosure statement

The abstract was accepted by American Heart Association at the 2021 hypertension conference, [https://doi.org/10.1161/hyp.78.suppl\\_1.P135](https://doi.org/10.1161/hyp.78.suppl_1.P135)

## Data statement

Data used for this analysis is available at <https://www.cdc.gov/nchs/nhis/1997-2018.htm>

## References

1. Hypertension prevalence among adults aged 18 and over : United States, 2017–2018. Accessed August 23, 2021. <https://stacks.cdc.gov/view/cdc/87559>
2. Marott SCW, Nielsen SF, Benn M, Nordestgaard BG. Antihypertensive treatment and risk of atrial fibrillation: a nationwide study. *Eur Heart J*. 2014;35(18):1205-1214. doi:10.1093/eurheartj/eh507
3. Butt DA, Harvey PJ. Benefits and risks of antihypertensive medications in the elderly. *J Intern Med*. 2015;278(6):599-626. doi:10.1111/joim.12446
4. Brunström M, Carlberg B. Effect of antihypertensive treatment at different blood pressure levels in patients with diabetes mellitus: systematic review and meta-analyses. *BMJ*. Published online February 24, 2016:i717. doi:10.1136/bmj.i717
5. Nwabuo CC, Dy SM, Weeks K, Young JH. Factors Associated with Appointment Non-Adherence among African-Americans with Severe, Poorly Controlled Hypertension. Burdmann EA, ed. *PLoS ONE*. 2014;9(8):e103090. doi:10.1371/journal.pone.0103090
6. Couto JE, Panchal JM, Lal LS, et al. Geographic Variation in Medication Adherence in Commercial and Medicare Part D Populations. *J Manag Care Pharm*. 2014;20(8):834-842. doi:10.18553/jmcp.2014.20.8.834
7. Lawson AJ, Hameed MA, Brown R, et al. Nonadherence to antihypertensive medications is related to pill burden in apparent treatment-resistant hypertensive individuals. *J Hypertens*. 2020;38(6):1165-1173. doi:10.1097/HJH.0000000000002398
8. Chang T, Bridges JFP, Bynum M, et al. Association Between Patient-Clinician Relationships and Adherence to Antihypertensive Medications Among Black Adults: An Observational Study Design. *J Am Heart Assoc*. 2021;10(14). doi:10.1161/JAHA.120.019943
9. Health Information Technology | HHS.gov. Accessed August 29, 2021. <https://www.hhs.gov/hipaa/for-professionals/special-topics/health-information-technology/index.html>
10. Stewart K, George J, Mc Namara KP, et al. A multifaceted pharmacist intervention to improve antihypertensive adherence: a cluster-randomized, controlled trial (HAPPY trial). *J Clin Pharm Ther*. 2014;39(5):527-534. doi:10.1111/jcpt.12185

11. Use of Health Information Technology to Improve Medication Adherence. Accessed August 24, 2021. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3641901/>
12. Newman-Casey PA, Killeen OJ, Renner M, Robin AL, Lee P, Heisler M. Access to and Experiences with, e-Health Technology Among Glaucoma Patients and Their Relationship with Medication Adherence. *Telemed E-Health*. 2018;24(12):1026-1035. doi:10.1089/tmj.2017.0324
13. Mayerová D, Abbas K. *Vaccine Confidence and Timeliness of Childhood Immunisation by Health Information Source, Maternal, Socioeconomic, and Geographic Characteristics in Albania*. *Epidemiology*; 2021. doi:10.1101/2021.03.09.21253180
14. Guo Z, Zhao SZ, Guo N, et al. Socioeconomic Disparities in eHealth Literacy and Preventive Behaviors During the COVID-19 Pandemic in Hong Kong: Cross-sectional Study. *J Med Internet Res*. 2021;23(4):e24577. doi:10.2196/24577
15. Svendsen MT, Bak CK, Sørensen K, et al. Associations of health literacy with socioeconomic position, health risk behavior, and health status: a large national population-based survey among Danish adults. *BMC Public Health*. 2020;20(1):565. doi:10.1186/s12889-020-08498-8
16. Lin YK, Lin M, Chen H. Do Electronic Health Records Affect Quality of Care? Evidence from the HITECH Act. *Inf Syst Res*. 2019;30(1):306-318. doi:10.1287/isre.2018.0813
17. NHIS - 2018 Data Release. Accessed August 22, 2021. [https://www.cdc.gov/nchs/nhis/nhis\\_2018\\_data\\_release.htm](https://www.cdc.gov/nchs/nhis/nhis_2018_data_release.htm)
18. Ferdinand KC, Senatore FF, Clayton-Jeter H, et al. Improving Medication Adherence in Cardiometabolic Disease. *J Am Coll Cardiol*. 2017;69(4):437-451. doi:10.1016/j.jacc.2016.11.034
19. Adherence in Hypertension | Circulation Research. Accessed August 21, 2021. <https://www.ahajournals.org/doi/full/10.1161/CIRCRESAHA.118.313220>
20. Sarkar U, Lyles CR, Parker MM, et al. Use of the Refill Function Through an Online Patient Portal is Associated With Improved Adherence to Statins in an Integrated Health System. *Med Care*. 2014;52(3):194-201. doi:10.1097/MLR.000000000000069
21. Keith McInnes D, Shimada SL, Rao SR, et al. Personal Health Record Use and Its Association with Antiretroviral Adherence: Survey and Medical Record Data from 1871 US Veterans Infected with HIV. *AIDS Behav*. 2013;17(9):3091-3100. doi:10.1007/s10461-012-0399-3
22. Lyles CR, Sarkar U, Schillinger D, et al. Refilling medications through an online patient portal: consistent improvements in adherence across racial/ethnic groups. *J Am Med Inform Assoc*. 2016;23(e1):e28-e33. doi:10.1093/jamia/ocv126
23. Miller TA. Health literacy and adherence to medical treatment in chronic and acute illness: A meta-analysis. *Patient Educ Couns*. 2016;99(7):1079-1086. doi:10.1016/j.pec.2016.01.020
24. Arbuckle C, Tomaszewski D, Brown L, et al. Exploring the relationship of digital information sources and medication adherence. *Comput Biol Med*. 2019;109:303-310. doi:10.1016/j.combiomed.2019.04.023

25. Young HN, Len-Rios ME, Brown R, Moreno MM, Cox E. How does patient-provider communication influence adherence to asthma medications? *Patient Educ Couns*. 2017;100(4):696-702. doi:10.1016/j.pec.2016.11.022
26. Schoenthaler A, Knafelz GJ, Fiscella K, Ogedegbe G. Addressing the Social Needs of Hypertensive Patients: The Role of Patient-Provider Communication as a Predictor of Medication Adherence. *Circ Cardiovasc Qual Outcomes*. 2017;10(9). doi:10.1161/CIRCOUTCOMES.117.003659
27. Archiopoli A, Ginossar T, Wilcox B, Avila M, Hill R, Oetzel J. Factors of interpersonal communication and behavioral health on medication self-efficacy and medication adherence. *AIDS Care*. 2016;28(12):1607-1614. doi:10.1080/09540121.2016.1192577
28. Wong C, Harrison C, Britt H, Henderson J. Patient use of the internet for health information. *Aust Fam Physician*. Published online December 2014. Accessed August 22, 2021. <https://search.informit.org/doi/abs/10.3316/informit.796476172571963>
29. Paccoud I, Baumann M, Le Bihan E, et al. Socioeconomic and behavioural factors associated with access to and use of Personal Health Records. *BMC Med Inform Decis Mak*. 2021;21(1):18. doi:10.1186/s12911-020-01383-9
30. Stormacq C, Van den Broucke S, Wosinski J. Does health literacy mediate the relationship between socioeconomic status and health disparities? Integrative review. *Health Promot Int*. 2019;34(5):e1-e17. doi:10.1093/heapro/day062
31. Bucy EP. Social Access to the Internet. *Harv Int J Press*. 2000;5(1):50-61. doi:10.1177/1081180X00005001005
32. Welch Cline RJ, Penner LA, Harper FWK, Foster TS, Ruckdeschel JC, Albrecht TL. The Roles of Patients' Internet Use for Cancer Information and Socioeconomic Status in Oncologist-Patient Communication. *J Oncol Pract*. 2007;3(3):167-171. doi:10.1200/JOP.0737001
33. Digital Distinction: Status-Specific Types of Internet Usage\* - Zillien - 2009 - Social Science Quarterly - Wiley Online Library. Accessed January 9, 2022. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1540-6237.2009.00617.x>