

# Original Research Article

## Standardization of screen aperture size for grading of dhaincha seeds

### Abstract

To standardize the sieve size for seed grading in dhaincha during Kharif, 2019–20 and 2020–21, an experiment was carried out. The native dhaincha species was processed using slotted perforated metal sieves with CRD design ranging S<sub>1</sub> - 1.4 mm (S), S<sub>2</sub>- 1.6 mm (S), S<sub>3</sub>- 1.8 mm (S), S<sub>4</sub> - 2.0 mm (S), and S<sub>5</sub>- 2.2 mm (S). The study realized that 2.0 mm and 2.2 mm sieves with the highest seed quality standards yield larger-sized seeds. Comparative to other sieves, the seed recovery % in the 1.4 mm screen was higher. The seed quality, nevertheless was better than the Minimum Seed Certification Standards as measured using 2.2 mm sieves. The SR (92.99%), PP (98.46%), PLS (90.37%), TW (26.16 g), SG (81.63%), TSL (25.2 cm), SDW (14.4 mg), SVI-I (2103) and SVI-II (1174) of the graded seeds gathered from the sieve 2.0 mm were noted in the combined average data of the two years. As an outcome, sorting dhaincha seeds via a 2.0 mm sieve is more effective and more economical than with a 2.2 mm sieve. The retrieval of good seed with percentage of germination and physical purity percentage over the permitted limits of Indian Minimum Seed Certification standard (IMSCS) can be accomplished with the sieve of 2.0 mm (S).

**Keywords:** Dhaincha, grading, screen size, dhaincha species

### Introduction

A conventional cultivation technique for preserving the fertility of the soil is referred to as "green manuring." While marginalizing the use of green manures in intensive crop cultivation, the rise of green farming has boosted the use of chemical fertilizers. Over time, the space beneath crops grown with green manure has shrunk, demonstrating this. Green manure crops are said to be grown over 1.23 million hectares of land in India (Anon., 2015). India's transition from a food-insecure to a food-secure region has been facilitated by increased fertilizer use at subsidized prices, but organic manure use, particularly green manure crops, has drastically dropped. In recent years, the overall price of inorganic fertilizer's has gone up, and soil productivity resilience is gaining momentum. Green manure crops are an affordable as well as effective way to bring down fertilizer expenditures while

maintaining the productivity and health of the soil. Almost every green manure crops, whether used in-situ or ex-situ, have all the plant nutrients necessary for boosting crop growth and maintaining the health of the soil.

Among the green manure crops dhaincha (*Sesbania aculeata*) is one of the most important green manure crop. The ease of establishment, fast growth leading to accumulation of large quantity of biomass, rich in nutrients especially nitrogen in short period and quick decomposition upon incorporation in paddy (puddle rice lands) can release nutrients as per the need of rice crop made dhaincha the most widely grown green manure crop. Further, it is promising for cultivation in salt affected ill drained soils and areas with higher rainfall (Parlawar *et al.*, 2003).

Seed processing is critical in scientific seed production because it preserves the physical integrity of seeds while also allowing for the recovery of ideal sized seeds for uniform crop establishment and growth. Seed size is a key determinant of seed vigour because it affects seed performance in soil. Although the seed mass after harvest contains a wide range of seed sizes, not all of them are equally valuable for planting. Farmers have long recognized the importance of utilizing homogeneous, high-viability seeds to achieve high emergence and growth. There is no recommended sieve size for sorting dhaincha seed at this time. The current grading process attempts to exclude non-viable seeds so that sound, disease-free seed of uniform size is accessible for planting, resulting in optimal plant population and increased yields.

One of the requirements in the Minimum Seed Certification Standard (MSCS) for seed approval by the Indian government is the determination of the best sieve size and kind of screen. The sieve size recommended for processing various agricultural seeds under the minimum seed certification standard appears to be more general and unsuitable for all newer types, resulting in low seed recovery (Anon., 1998). There is a pressing need to standardize the sieve size for grading dhaincha seed, as it is frequently noted that seed growers are losing a significant amount of good seed that is classified as a rejection. Given the high demand from farmers for certified seeds of dhaincha, this is an essential necessity.

As a result, the current study on standardizing sieves for grading dhaincha seed was planned and undertaken.

## Materials and methods

The investigation was conducted by using freshly gathered bulk dhaincha seeds from the National Seed Project (Crops), Seed Block, UAS, Raichur, Karnataka. The seeds' initial moisture content had been lowered to around 9 per cent by drying them out in the shade. At the Seed Quality Assurance and Research Laboratory, Seed Unit, University of Agricultural Sciences, Raichur, research have been carried out in the lab to standardize the seed size of the aperture for grading of dhaincha seeds during 2019–20 and 2020–21. Five separate sieve sizes were used for sorting the seeds. The seeds were graded using five different sieve sizes.

The experiment was laid out in Complete Randomized Design with four replications to impose treatments i.e., S<sub>1</sub>- 1.4 mm (S), S<sub>2</sub>- 1.6 mm (S), S<sub>3</sub>- 1.8 mm (S), S<sub>4</sub>- 2.0 mm (S) and S<sub>5</sub>- 2.2 mm (S) screen aperture sizes. The seeds obtained from different sieves were assessed for various seed quality seed parameters like seed recovery, physical purity, pure live seed, test weight, germination, seedling length, SDW and SVI. Gomez and Gomez (1984)'s Fischer method of Analysis of Variance approach was employed for the statistical examination as well as interpretation of the results of the experiment. For experiments in laboratories, the F test's 1% level of significance was employed.

## Results and discussion

By eliminating inferior seeds of the same species, sorting aims at improving the uniformity of the seed lot. Small seeds, which is believed to comprise void, underdeveloped, and low vigour seeds, are flung out during size grading. For practically all of the characteristics under inspection, highly significant variance was observed among the various sieve sizes. Menaka and Balamurugan (2008) reported on the value of seed size.

According to Suma *et al.* (2014), sorting is one of the essential after harvest management processes that homogenizes the seed lot and promotes consistent germination with increased planting value.

Seed size exerted a significant influence on the seed recovery, test weight, physical purity, pure live seed, germination, total seedling length, seedling dry weight and seedling vigour index (Table 1, 2 and 3).

## Seed recovery, physical purity and pure live seed

Assessing the results on SR (%) displayed in Table 1 reveals that varying screen sizes during 2019–20 and 2020–21 and aggregated data of two years substantially impacted seed recovery (%). The seeds retained on S<sub>1</sub> (1.4 mm) displayed maximum SR (98.18%) and lowest physical purity and PLS (95.47 and 68.69 %) and the lowest SR (87.95%) was in S<sub>5</sub> (2.2 mm) with higher Physical purity and PLS (99.16 % and 83.78 %), as per pooled data for two years.

From the above results seed recovery was significantly influenced by different sieve sizes. The seeds graded with 1.4 mm sieve size recorded higher SR (%) compared to seeds graded with 1.6, 1.8, 2.0 and 2.2 mm sieves. The reduction in SR is due to increase in screen aperture size which led to removal undersized seeds keeping good and bold seeds at top of the screen. As the screen size decreased from 2.2 to 1.4 mm, the per cent seed recovery increased. This is in conformity with the findings of Anuradha *et al.* (2009) in chickpea, Ganiger *et al.* (2016) in greengram and Kausal *et al.* (2008) in soybean. Significantly higher physical purity was recorded in processed seeds i.e., seeds retained on sieves S<sub>5</sub> (2.2 mm) and S<sub>4</sub> (2.0 mm) as compared to S<sub>1</sub> (1.4 mm), S<sub>2</sub> (1.6 mm) and S<sub>3</sub> (1.8 mm) due to effective processing (Fig. 1). Further, all the impurities were removed and whereas the seeds retained on smaller screens had impurities so less physical purity was observed. Similar observations of improved physical purity have been reported by Ganiger *et al.* (2016) in greengram and soybean.

## Test weight, germination, total seedling length and seedling dry weight

In pooled mean data of two years, highest test weight, germination, total seedling length, seedling dry weight, SVI I and II (27.20 g and 84.47 %, 26.3 cm and 15.3 mg, 2225 and 1292) was recorded in seeds retained on S<sub>5</sub> (2.2 mm) and the lowest was recorded in S<sub>1</sub> (1.4 mm) (22.32 g and 71.94 %, 21.3 cm, 11.1 mg, 1561 and 795), respectively (Table 2 and Fig. 2) The data was presented in Table 3.

.From the above obtained results, the seeds retained on 2.2 mm screen were larger in size and gave higher test weight and seed germination (Plate 1) which may be due to undesirable materials, low graded seeds are dropped through screens and retaining only goodgraded seeds which has better reserve food material available in the storage tissues.

These results are in conformity with findings of Dharmalingam and Ramakrishnan (1978) in black gram, Kausal *et al.* (1993) in sorghum and greengram, Negi *et al.* (1988) in soybean, Paul *et al.* (1997) in mustard, Vishwanth *et al.* (2006) in french bean, Kausal *et al.* (1993) in sorghum and greengram and Raghavendra Rao *et al.* (1993) in sorghum.

Total seedling length and seedling dry weight was recorded highest in seeds which were retained on large screen size i.e. 2.2 mm (Plate 2). While, the lowest was recorded in small screen size of 1.4 mm. This may be due to adequate quantity of reserve food material in the small to medium sized seeds endosperm and better supply of food to the embryo for a long period of time which helped for higher mobilization efficiency of reserve food material during germination leading to more number of cells per cotyledon in the form of reserve food resulting in increased root, shoot length (Guldan and Brun, 1985). As there has been better seedling growth, which might have led to an increase in mean seedling dry weight and similar results were also documented by Sudeep Kumar *et al.* (2010) in fieldbean.

## **Conclusion**

The study inferred that, a sieve size of 2.0 (S) mm registered SR of (92.99 %), PP (98.48 %), SG (81.63 %), TW (26.16 g), PLS (90.37 %), TSL (25.2 cm), SDW (14.4 mg) and SVI-I and II (2103 and 1174) was found to above the minimum seed certification standards. Hence the dhaincha can be processed using 2.0 mm (S) grading sieve for better seed recovery and quality.

## **References**

Anonymous, 1998, Minimum seed certification standards

Anonymous, 2015, Fertilizer association of India. *Fertilizer Statistics*. II-63.

Anuradha, R., Balamurugan, P., Srimathi, P. and Sumathi, S., 2009, Influence of seed size on seed quality of chick pea (*Cicer arietinum* L.). *Legume Res.*, 32: 133-135.

Dharmalingam, C. and Ramakrishnan, V., 1978, Seed quality in relation to seed size and seed coat colour variation in black gram (*Vigna mungo* L. Heppre). *Seed Res.*, 6(2): 101-109.

- Ganiger, B. S., Basavegowda, Lokesh, G. Y. and Rekha, 2016, Standardization of screen sizes for green gram seed processing. *Intl. Quart. J. Life Sci.*, 11(4): 2379-2381.
- Gomez, K. A. and Gomez, Z. A., 1984, Statistical procedures for agricultural research. A *Wiley Inter. Sci. Publication*, New York.
- Guldan, S. J. and Brun, W. A., 1985, Relationship of cotyledon cell number and seed respiration to soybean seed growth. *Crop Sci.*, 25: 951-996.
- Kausal, R. T., Changade, S. P. and Patil, V. N., 1993, Standardization of sieve sizes for grading crop seeds. *Seed Res.*, 2: 826-834.
- Kausal, R. T., Jeughale, G. S., Kakade, S. U. and Pravitrakar, N. R., 2008, Studies on optimum sieve size and type of screen for grading soybean seed. *Int. J. Agric. Sci.*, 4: 59-62.
- Menaka, C. and Balamurugan, 2008, Seed grading techniques in Amaranthus cv. CO-5. *Plant Arch.*, 8: 729-731.
- Negi, H. C. S., Kant, K. and Verma, M. M., 1988, Improving germination by grading in soybeans. *Seeds and Farms*, 14(7): 17-19.
- Parlawar, N. D., Giri, D. G. and Adpawar, R. M., 2003, Influence of seed rate, row spacing and phosphate level on nutrient uptake in dhaincha. *J. Soils and Crops*, 13(2): 364-67.
- Paul, S. R., Sarma, N. N. and Sarma, D., 1997, Effect of seed size on germination and seedling vigour of mustard. *Seed Tech. News*, 27(3): 6.
- Raghavendra Rao, D. V. S., Murlimohanreddy, B., Ankaiah, K., Saibabu, K, G. and Hussaini, S. H., 1993, Influence of sieve sizes on storability of CSH-5 sorghum hybrid. *J. Res., AndhraPradesh Agril. Univ.*, 21(3): 163-164.
- Sudeep Kumar, E., Channaveerswami, A. S., Merwade, M. N., Rudra, N. V. and Krishna, A., 2010, Influence of nipping and hormonal sprays on growth and seed yield in field bean [*Lablab purpureus* (L.) Sweet] genotypes. *Int. J. Econ. Plants*, 5(1): 8-14.

Suma, N., Srimathi, P. and Sumathi, S., 2014, Influence of size grading on seed and seedling quality characteristics of *Sesamum indicum*. *Int. J. Curr. Microbiol. App. Sci.* 3(6): 486-490.

Vishwanath, K., Kalappa, V. P. and Rajendra Prasad, S., 2006, Standardization of screen sizes for french bean seed processing. *Seed Res.*, 34(1): 77-81.

UNDER PEER REVIEW

**Table 1. Seed recovery, physical purity and pure live seed as influenced by different screen sizes in dhaincha**

Treatments	Seed recovery (%)			Physical purity (%)			Pure live seed (%)		
	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
S <sub>1</sub> (1.4 mm)	98.92	98.28	<b>98.18</b>	95.77	95.18	<b>95.47</b>	67.61	69.76	<b>68.69</b>
S <sub>2</sub> (1.6 mm)	98.44	97.92	<b>97.91</b>	96.48	96.43	<b>96.46</b>	71.63	72.87	<b>72.25</b>
S <sub>3</sub> (1.8 mm)	96.05	96.33	<b>95.33</b>	97.49	97.48	<b>97.49</b>	74.34	75.55	<b>74.94</b>
S <sub>4</sub> (2.0 mm)	94.08	94.03	<b>92.99</b>	98.47	98.46	<b>98.46</b>	79.50	81.25	<b>80.37</b>
S <sub>5</sub> (2.2 mm)	86.35	87.66	<b>87.95</b>	99.21	99.11	<b>99.16</b>	83.65	83.91	<b>83.78</b>
<b>Mean</b>	<b>94.77</b>	<b>94.84</b>	<b>94.47</b>	<b>97.48</b>	<b>97.33</b>	<b>97.41</b>	<b>75.35</b>	<b>76.67</b>	<b>78.01</b>
S.Em±	0.59	0.66	0.31	0.37	0.28	0.39	1.34	1.19	0.94
CD at 1 %	2.24	2.51	1.16	1.55	1.18	1.62	5.60	4.97	3.90

**Table 2. Test weight and germination as influenced by different screen sizes in dhaincha**

Treatments	Test weight (g)			Germination (%)		
	2019	2020	Pooled	2019	2020	Pooled
S <sub>1</sub> (1.4 mm)	21.99	22.45	<b>22.22</b>	70.58	73.30	<b>71.94</b>
S <sub>2</sub> (1.6 mm)	23.93	24.34	<b>24.13</b>	74.20	75.59	<b>74.89</b>
S <sub>3</sub> (1.8 mm)	25.30	25.37	<b>25.34</b>	76.27	77.51	<b>76.89</b>
S <sub>4</sub> (2.0 mm)	26.28	26.04	<b>26.16</b>	80.72	82.54	<b>81.63</b>
S <sub>5</sub> (2.2 mm)	27.21	27.19	<b>27.20</b>	84.28	84.66	<b>84.47</b>
<b>Mean</b>	<b>24.94</b>	<b>25.08</b>	<b>25.01</b>	<b>77.21</b>	<b>78.72</b>	<b>77.96</b>
S.Em±	0.33	0.40	0.36	1.02	1.14	0.30
CD at 1 %	1.39	1.69	1.51	4.26	4.76	1.24

**Table 3. Total seedling length and seedling dry weight as influenced by different screen sizes in dhaincha**


Treatments	Total seedling length (cm)			Seedling dry weight (mg)		
	2019	2020	Pooled	2019	2020	Pooled
S <sub>1</sub> (1.4 mm)	21.3	21.3	<b>21.3</b>	10.9	11.2	<b>11.1</b>
S <sub>2</sub> (1.6 mm)	24.1	23.4	<b>23.8</b>	11.8	12.2	<b>12.0</b>
S <sub>3</sub> (1.8 mm)	24.6	23.7	<b>24.1</b>	13.3	13.7	<b>13.5</b>
S <sub>4</sub> (2.0 mm)	25.5	24.9	<b>25.2</b>	14.4	14.3	<b>14.4</b>
S <sub>5</sub> (2.2 mm)	27.7	26.2	<b>27.0</b>	15.3	15.2	<b>15.3</b>
<b>Mean</b>	<b>24.6</b>	<b>23.9</b>	<b>24.3</b>	<b>13.1</b>	<b>13.3</b>	<b>13.2</b>
S.Em±	0.3	0.2	0.2	0.2	0.3	0.2
CD at 5 %	1.2	0.7	0.8	0.7	1.2	0.9

**Table 4. Seedling vigour index as influenced by different screen sizes in dhaincha**

Treatments	Seedling vigour index-I			Seedling vigour index-II		
	2019	2020	Pooled	2019	2020	Pooled
S <sub>1</sub> (1.4 mm)	1550	1572	<b>1561</b>	769	821	<b>795</b>
S <sub>2</sub> (1.6 mm)	1732	1778	<b>1755</b>	877	923	<b>900</b>
S <sub>3</sub> (1.8 mm)	1855	1848	<b>1851</b>	1013	1029	<b>1021</b>
S <sub>4</sub> (2.0 mm)	2131	2075	<b>2103</b>	1166	1181	<b>1174</b>
S <sub>5</sub> (2.2 mm)	2211	2239	<b>2225</b>	1294	1290	<b>1292</b>
<b>Mean</b>	<b>1896</b>	<b>1902</b>	<b>1899</b>	<b>1024</b>	<b>1049</b>	<b>1036</b>
S.Em±	32	30	21	27	8	15
CD at 5 %	132	126	63	115	34	61



**Plate 1. Germination (%) as influenced by different screen sizes in dhaincha**



**Plate 2. Total seedling length (cm) as influenced by different screen sizes in dhaincha**




Fig. 1. Seed recovery, physical purity and pure live seed as influenced by different screen sizes in dhaincha




Fig. 2. Germination and test weight as influenced by different screen sizes in dhaincha