

Effect of different dosages and scheduling time of plant growth regulators and time of application of defoliators on growth and yield of *Bt* cotton (*Gossypium hirsutum* L.) under high density planting system (HDPS)

Comment [H1]: Too long their rephrase

ABSTRACT: A field trial to determine the effect of different dosages, application time of plant growth regulators and defoliators on growth and yield of cotton under high density planting system was conducted at Regional Agricultural Research Station, Warangal during kharif-2022. The experiment consisted of eleven treatments laid out in Randomized Block Design with three replications. The treatment details were T₁: Application of Mepiquat chloride (MC) 25g a.i ha⁻¹ @40&55 DAE+ Etherel 2000ppm @40% boll burst, T₂: MC 25 g a.i ha⁻¹ @40&55 DAE+ Etherel 2000ppm @60% boll burst, T₃: MC 20g a.i ha⁻¹ @40DAE, MC 30g a.i ha⁻¹ @60DAE+ Etherel 2000ppm @ 40% boll burst T₄: MC 20g a.i ha⁻¹ @40DAE, MC 30g a.i ha⁻¹ @60DAE+ Etherel 2000ppm @ 60% boll burst T₅: MC 20 g a.i ha⁻¹ @40,55 & 75 DAE+ Etherel 2000ppm @40% boll burst, T₆: MC 20 g a.i ha⁻¹ @40,55 & 75 DAE+ Etherel 2000ppm @40% boll burst, T₇: MC 25 g a.i ha⁻¹ @40,55 & 75 DAE+ Etherel 2000ppm @40% boll burst, T₈: MC 25 g a.i ha⁻¹ @40,55 & 75 DAE+ Etherel 2000ppm @60% boll burst, T₉: 20g a.i ha⁻¹ @ 40DAE, 25g a.i ha⁻¹ @55DAE & 30g a.i ha⁻¹ @70DAE +Etherel 2000ppm @ 40% boll burst, T₁₀: 20g a.i ha⁻¹ @ 40DAE, 25g a.i ha⁻¹ @55DAE & 30g a.i ha⁻¹ @70DAE +Etherel 2000ppm @ 60% boll burst and T₁₁: Control. Results revealed that foliar application of MC @ 20 g a.i ha⁻¹ @ 40,55 & 70DAE in conjunction with Etherel 2000ppm @60% boll burst recorded higher seed cotton yield (?????kg). Plant growth, dry matter accumulation and stalk yield were recorded highest in control plot and lowest with spraying of : 20g a.i ha⁻¹ @ 40DAE, 25g a.i ha⁻¹ @55DAE & 30g a.i ha⁻¹ @70DAE +Etherel 2000ppm @ 40% boll burst. Among Mepiquat chloride treatments with similar doses, all the agronomic traits of cotton crop are positively influenced with application of 2000ppm of etherel @60% boll burst. The spraying of M.C @ 20 g a.i ha⁻¹ at 40,55 and 70 DAE + Etherel 2000ppm @ 60% boll burst would be economically ideal to the farming community.

Key words: Plant growth regulators, Mepiquat chloride, Defoliators, Dry matter production, Etherel.

INTRODUCTION

Cotton (*Gossypium hirsutum* L.) is an important cash crop of the world. Because of its grate contribution to Indian agriculture in terms of employment generation, industrial development and national economy it is referred as "King of fibres" (Uma and Murali 2019). According to 2021-22 data global area under cotton is 32.7 million ha with production of 121.6 million bales, where each bale weighs 217.72 kgs. (www.pitsau.edu.in cotton outlook January-2022). During 2021-22 in India the cotton area, production and productivity projected as 12,371 thousand ha, 31,117 thousand bales of each 170 kgs, 428 kg ha⁻¹ (www.indiastat.com). In Telangana, in 2021-22 it was recorded that area under cotton is 18.89 Lakh ha, production of cotton is 48.78 Lakh bales and Productivity of 439 kg ha⁻¹ (www.aiccip.cicr.org.in). In Telangana, Nalgonda is leading district with area of 294074.18 ha under cotton cultivation.

Cotton progress depends on environment and seasonal practices, impacting growth, development, and yield. Maximizing cotton yield involves adjusting plant density per unit area, but it varies with cultivars used. Optimal plant spacing plays a pivotal role in adjusting plant density to enhance cotton productivity, particularly under irrigated conditions (Zaman *et al.*, 2021). High density planting system (HDPS) of cotton is a method of cultivation, where the space between the rows and plants in a row is decreased in order to increase number of plants per unit area, which will result in more number of

bolts per unit area and subsequent increase in yield. But the problem with HDPS is over crowding of plants causing excessively taller height, more vegetative [growth](#); mutual shading which may intern leads to reduction in yield. (Priyanka *et al.*, 2021). Thus [plant growth regulators have](#) to be applied to HDPS cotton which will reduce the plant height of cotton and helps in increasing productivity and profitability of the HDPS cotton. Mepiquat chloride (1,1-dimethyl-piperidinium chloride), a common plant growth regulator, is extensively employed to control cotton growth structure, modulate plant growth, and expedite maturation in conditions of high planting densities (Stuart *et al.*, 1984). The application of growth regulators led to dose-dependent reductions in plant height, decreased height to node ratio, increased boll weight, and delayed maturity. Mepiquat chloride enhanced compactness and reduced nodes (Reddy *et al.*, 1990). Environmental factors, notably temperature impact mepiquat chloride's influence on cotton potentially causing varied outcomes among different locations (Rosolem *et al.*, 2013).

Plant defoliant is the chemical which will promote the development of abscission layers in the leaf petioles, leading to the premature desiccation and shedding of foliage compared to its natural occurrence. [They](#) will reduce the trash content in the lint and enhance the lint quality. Defoliant is a chemical that modifies plant metabolism, inducing leaf shedding. In agriculture, they clear crop leaves to ease harvesting. Ethylene-based agents promote leaf drop, synchronous boll opening, and sunlight exposure, readying cotton for mechanical harvest. Timely and precise defoliation is vital for a successful harvest (Sravanthi *et al.*, 2022). Various defoliant including Dropp ultra, Etherel, NaCl, Paraquat and Urea are employed in cotton at varying rates depending on environmental conditions and cultivar used. [The experiment aimed to determine the effect of different dosages, application time of plant growth regulators and defoliant on growth and yield of cotton under high density planting system.](#)

METHODS AND MATERIALS

Field experiment was conducted at RARS, Warangal during *kharif* -2022 season. The experimental site is geographically located in central agro climatic zone of Telangana. The soil of experimental site is clay loam in texture with nearly neutral p_H in reaction (7.2), E.C (0.53 dsm^{-1}), low in organic carbon (0.43%), medium in available N (209 kg ha^{-1}) & phosphorous (28 kg ha^{-1}) and high in available potassium (334 kg ha^{-1}).

The experiment was laid out in a Randomized [Block Design](#) with eleven treatments. The treatment details are T₁: Mepiquat chloride (MC) 25 g a.i ha^{-1} @ 40 & 55 DAE + Etherel 2000ppm @ 40% boll burst, T₂: MC 25 g a.i ha^{-1} @ 40 & 55 DAE+ Etherel 2000ppm @ 60% boll burst, T₃: MC 20 g a.i ha^{-1} @ 40 DAE, MC 30 g a.i ha^{-1} @ 60DAE+ Etherel 2000ppm @ 40% boll burst T₄: MC 20 g a.i ha^{-1} @ 40DAE, MC 30 g a.i ha^{-1} @ 60DAE+ Etherel 2000ppm @ 60% boll burst T₅: MC 20 g a.i ha^{-1} @ 40,55 & 75 DAE+ Etherel 2000ppm @ 40% boll burst, T₆: MC 20 g a.i ha^{-1} @ 40,55 & 75 DAE + Etherel 2000ppm @ 40% boll burst, T₇: MC 25 g a.i ha^{-1} @ 40, 55 & 75 DAE+ Etherel 2000ppm @ 40% boll burst, T₈: MC 25 g a.i ha^{-1} @ 40, 55 & 75 DAE+ Etherel 2000ppm @ 60% boll burst, T₉: 20 g a.i ha^{-1} @

[Type here]

40DAE, 25 g a.i ha⁻¹ @ 55 DAE & 30 g a.i ha⁻¹ @ 70DAE +Etherel 2000ppm @ 40% boll burst, T₁₀: 20 g a.i ha⁻¹ @ 40 DAE, 25 g a.i ha⁻¹ @ 55 DAE & 30 g a.i ha⁻¹ @ 70DAE + Etherel 2000ppm @ 60% boll burst, T₁₁: Control. The treatments were sown at a spacing of 80×20 cm on 25/06/2022. Recommended dose of fertilizers 120-60-60 NPK ha⁻¹, urea, di ammonium phosphate, murjate of potash were used in this experiment. Adequate plant protection measures were taken as per requirement. Randomly five plants were selected and tagged from each plot for recording growth and yield parameters periodically and at harvest through destructive and non destructive sampling. The rainfall of 1223 mm was received in 57 rainy days during the entire period of crop growth. The mean maximum temperature and minimum temperature recorded was 30.6°C and 21.7°C respectively. Data recorded was analyzed as suggested by Gomez and Gomez 1984.

RESULTS AND DISCUSSIONS

Initial and final plant population (No. ha⁻¹)

Results relating to initial and final plant population as influenced by application of different doses and various time of application of Mepiquat chloride and different stages of Etherel application under HDPS was reduced in all the treatments over initial plant population due to heavy rainfall during early crop growth stages, plants lost during intercultural operations and dry matter collection.

Growth parameters

Plant height has importance as it shapes node and internode structure enabling the growth of essential sympodial branches that impact productivity. Table 2 displays plant height data across growth stages i.e., 30, 60, 90, 120 days after sowing (DAS) and at harvest. As application of growth regulators started at 40 Days after emergence, at 30 DAS plant height did not significantly differ among treatments. However as spraying of growth regulator treatments stopped at 75 Days after emergence, from 90 DAS of sowing to till the harvest change in plant height followed the same trend. At harvest control treatment showed highest plant height (121.73 cm), while lowest plant height was recorded with (T₁₀) M.C 20, 25, 30 g a.i ha⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 60% boll burst (95.3cm). Similar results were observed by Priyanka et al., (2022), Khetre et al., (2018) and Collins et al (2017).

Mepiquat chloride is a growth regulator used in cotton, which hinders the biosynthesis of gibberellic acid within the plant. Gibberellic acid is a key regulator of cell elongation and division, primarily responsible for stem elongation. Inhibition of gibberellic acid counters the elongation effect, resulting in shorter stems and overall reduced plant height (Suttle (1995) and Rademacher (2000)). Mepiquat chloride treatments application of 2000ppm Etherel at 40% boll burst resulted in lowest plant height compared to application of 2000ppm Etherel at 60% boll burst. This might be due to the reason that use of defoliant at early stages, potentially promoted the accelerated shedding of leaves, leading to a

[Type here]

decrease in the absorption of photosynthates and ultimately causing a reduction in the height of the plants. [Similar results were](#) observed by Sravanthi *et al.*, (2022), Kulvir and Pankaj (2015), Singh *et al.* (2017) and Mrunalini *et al.* (2019).

Dry matter production serves as the foundation for plant growth, development and yield. The use of plant growth regulators impacted the accumulation of dry matter [and its](#) accumulation [was](#) influenced by various dosages and time of application of Mepiquat chloride and various stages of application of Etherel [Table 3]. However dry matter accumulation followed same trend as plant height from 90 DAS to till harvest of the crop. At harvest control treatment was superior (5988.26 kg ha⁻¹) [outperformed](#) other treatments in dry matter accumulation and lowest [figure](#) was recorded with [the](#) application of M.C 20, 25, 30 g a.i ha⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst (T₉) (4253.26 kg ha⁻¹). Similar findings were noted by Priyanka *et al.* (2022), Kaul *et al.* (2016) [and](#) Paslawar *et al.* (2015). The decrease in the accumulation of dry matter could be because of both the observed decline in plant height and the disruption in the source sink relationship resulting from the application of Mepiquat chloride Priyanka *et al.*, (2021).

Significant enhancement in dry matter was observed when defoliant were applied at a later stage compared to their early application. Delaying the defoliation process enabled increased carbon assimilation and the allocation of photosynthates towards the growth of cotton bolls and greater biomass accumulation. These findings align with the outcomes [as](#) reported by Kulvir *et al.* (2015), Mrunalini *et al.* (2019) [and](#) Sravanthi *et al.* (2022). The decrease in the dry matter accumulation over control as effected by different Mepiquat chloride and Etherel treatments was depicted in the Fig 1- where reduction in plant dry matter ranged from 11.82% with Spraying of MC 25 g a.i ha⁻¹ @ 40 & 55 DAE+ Etherel 2000ppm @ 60% boll burst, T₃: MC 20g a.i ha⁻¹ @40DAE (T₂) to 28.97% in Spraying of M.C 20, 25, 30 g a.i ha⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst (T₉).

YIELD

[Results](#) in Table 3 [showed](#) highest seed cotton yield with spritzing of MC 20 g a.i ha⁻¹ @40,55 & 75 DAE+ Etherel 2000ppm @60% boll burst (T₆) (3163.57 Kg ha⁻¹) and lowest in [the](#) control treatment (T₁₁) (2262.17 Kg ha⁻¹). According to Patel *et al.*, (2021), Brar *et al.*, (2020) [the](#) application of higher doses of M.C at higher no of doses resulted in higher seed cotton yield. In contrary the application [of](#) 20 g of a.i ha⁻¹ at 3 stages resulted in higher economic yield, this might be due to higher level of hindrance to gibberlin synthesis at higher doses of M.c , which in [turn](#) resulted in very low vegetative growth, which is necessary to support reproductive growth. [This could be due ot the](#) better [partit](#)ioning of photosynthates to reproductive [plant](#) parts. In cotton excessive vegetative growth occurs at the cost of economic yield, application of Mepiquat chloride regulates the excessive vegetative growth, and resulted in more number of yield components and highest seed cotton yield Uma *et al.* (2019). Application of 20 g a.i ha⁻¹ resulted in higher yields, this might be due higher level of hindrance to Gibberelin synthesis at high doses of M.C.

Comment [H2]: Of what????

Among the [different](#) stages of [spraying](#) of 2000ppm of etherel at 60% boll burst resulted in higher seed cotton yield compared to etherel application at 40% boll burst stage. [This could be due to](#) early

[Type here]

leaf defoliation that adversely affect all agronomic characteristics of crop, on contrary application of defoliators at correct time preferably not before 60% boll burst allowed the crop to adequately partition the carbohydrates which ultimately resulted in higher yields. Similar results were observed with

TREATMENTS	Plant population (No. ha ⁻¹)
------------	---

Sravanthi *et al.* (2022), Raghavedra and Rama (2020) and Singh *et al* (2017).

Results in Table 4 showed highest stalk yield in control (T₁₁) treatment (6673.33 Kg ha⁻¹) and lowest in (T₁₀) M.C application of 20,25 and 30 g a.i ha⁻¹ @ 40,55 and 60 DAE + Etherel 2000ppm @ 60% boll burst due to the reduction in plant height, accompanied by diminished accumulation of dry matter. Similar outcomes were noticed by Priyanka *et al.* (2022) and Patel *et al.* (2021). Stalk yield varied with stage of application of etherel, among both stages i.e., 40% , 60% boll burst stages such that application of etherel at 60 % boll burst resulted in higher stalk yield in all of M.C applied treatments.

Delayed defoliation would facilitate time for more carbon accumulation and better partitioning of photo assimilates to all the plant parts and similar results were observed by Sravanthi *et al.* (2022) and Mrunalini *et al.* (2019). Higher Harvest index was observed with with T₆ (MC 20 g a.i ha⁻¹ @40,55 & 75 DAE application + Etherel 2000ppm spraying @40% boll burst) (36.89) and lowest recorded with control (25.28). Priyanka *et al.* (2020) found similar outcome.

Table 1: Initial and final of *Bt* cotton as influenced by application of different doses and application time Mepiqaut chloride and different stages of application of Etherel under HDPS.

[Type here]

No	TREATMENT imposed	Initial Plant population	Final Plant population
T ₁	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 40% boll burst.	61771	59271
T ₂	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 60% boll burst.	61354	59271
T ₃	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55 DAE + Etherel 2000 ppm @ 40% boll burst.	61458	59375
T ₄	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55DAE + Etherel 2000 ppm @ 60% boll burst.	61354	58854
T ₅	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	61563	59201
T ₆	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	61458	59375
T ₇	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	61979	59757
T ₈	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	61250	58750
T ₉	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst.	61563	59479
T ₁₀	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 60% boll burst.	61458	59375
T ₁₁	Control (Water spray @ 40, 55 & 70 DAE)	61667	59167
	<u>Statistics missing</u>		

[Type here]

Table 2: Plant height of *Bt* cotton as influenced by application of different doses and application time Mepiqaut chloride and different stages of application of Etherel under HDPS.

Treatments		Plant height (cm)				
No	Treatment imposed	30 DAS	60 DAS	90 DAS	120 DAS	Harvest
T ₁	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 40% boll burst.	11.11	19.91	58.00	99.27	110.20
T ₂	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 60% boll burst.	11.40	19.86	59.23	100.07	110.37
T ₃	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55 DAE + Etherel 2000 ppm @ 40% boll burst.	10.83	19.18	56.40	97.73	109.97
T ₄	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55DAE + Etherel 2000 ppm @ 60% boll burst.	11.79	19.17	57.73	98.23	109.30
T ₅	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	11.59	21.90	52.60	87.20	99.07
T ₆	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	11.76	21.26	53.07	88.00	98.83
T ₇	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	11.23	19.70	50.77	86.20	97.20
T ₈	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	10.62	19.71	51.90	87.53	96.90
T ₉	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst.	11.60	20.77	48.50	84.00	95.87
T ₁₀	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 60% boll burst.	11.14	20.64	49.77	85.07	96.23
T ₁₁	Control (Water spray @ 40, 55 & 70 DAE)	10.93	27.31	66.43	112.70	121.73
SEm ±		0.30	0.94	1.96	4.05	3.70
C.D (p=0.05)		NS	2.8	5.79	11.95	10.93

Comment [H3]: I am not sure

[Type here]

Table 3: Dry matter of *Bt* cotton as influenced by application of different doses and application time Mepiqaut chloride and different stages of application of Etherel under HDPS

Treatments		Crop dry matter (Kg. ha ⁻¹)				
No	Treatment imposed	30 DAS	60 DAS	90 DAS	120 DAS	Harvest
T ₁	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 40% boll burst.	115.76	1696.47	3479.80	4219.60	5176.27
T ₂	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 60% boll burst.	121.36	1686.60	3516.35	4300.23	5280.27
T ₃	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55 DAE + Etherel 2000 ppm @ 40% boll burst.	119.41	1627.70	3294.00	4068.87	5082.20
T ₄	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55DAE + Etherel 2000 ppm @ 60% boll burst.	121.79	1621.23	3360.33	4183.54	5166.87
T ₅	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	119.67	1847.53	2804.15	3557.77	4437.77
T ₆	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	109.41	1822.97	2851.28	3604.97	4521.63
T ₇	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	115.23	1656.53	2688.00	3398.97	4378.97
T ₈	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	116.17	1653.57	2735.10	3494.11	4440.77
T ₉	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst.	119.05	1785.93	2598.50	3309.90	4253.26
T ₁₀	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 60% boll burst.	123.49	1732.93	2637.88	3362.57	4355.78
T ₁₁	Control (Water spray @ 40, 55 & 70 DAE)	124.78	2137.75	4231.50	5011.60	5988.26
SEm ±		5.13	77.63	220.40	234.59	237.09
C.D (p=0.05)		NS	228.99	650.08	691.93	699.31

Comment [H4]: Not sure

[Type here]

Table 4: Yield of *Bt* cotton as influenced by application of different doses and application time Mepiquat chloride and different stages of application of Etherel under HDPS.

Treatments		Seed Cotton Yield (kg ha ⁻¹)	Stalk yield (kg ha ⁻¹)	Harvest Index (%)
No	Treatment imposed			
T ₁	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 40% boll burst.	2756.57	6028.13	31.24
T ₂	M.C 25 g a.i ha ⁻¹ @ 40 & 55 DAE + Etherel 2000 ppm @ 60% boll burst.	2783.07	6027.60	31.46
T ₃	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55 DAE	2613.47	5923.07	30.43


[Type here]

	+ Etherel 2000 ppm @ 40% boll burst.			
T ₄	M.C 20 g a.i ha ⁻¹ @ 40 DAE and 30 g a.i ha ⁻¹ at 55DAE + Etherel 2000 ppm @ 60% boll burst.	2643.93	5982.20	30.43
T ₅	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	3129.73	5420.80	36.58
T ₆	M.C 20 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	3163.57	5408.27	36.89
T ₇	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 40% boll burst.	3013.50	5362.27	35.91
T ₈	M.C 25 g a.i ha ⁻¹ @ 40, 55 & 70 DAE + Etherel 2000 ppm @ 60% boll burst.	3041.20	5352.33	36.10
T ₉	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 40% boll burst.	2921.67	5317.07	35.33
T ₁₀	M.C 20, 25, 30 g a.i ha ⁻¹ @ 40, 55 & 70 DAE respectively + Etherel 2000 ppm @ 60% boll burst.	2942.67	5306.47	35.41
T ₁₁	Control (Water spray @ 40, 55 & 70 DAE)	2262.17	6673.33	25.28
SEm ±		113.15	203.12	1.07
C.D (p=0.05)		333.75	599.11	3.16

Comment [H5]: ????

[Type here]

Fig 1: Change in dry matter of *Bt* cotton as influenced by application of different doses and application time Mepiquat chloride and different stages of application of Etherel under HDPS



How about control treatment

CONCLUSION

Application of higher doses of Mepiquat chloride at 3 stages resulted in lowest plant height, dry matter accumulation and stalk yield. Spraying of Etherel 2000ppm at 60% boll burst resulted in superior agronomic characteristics compared to etherel application at all M.C application treatments. However, higher economic yields were observed with spraying of M.C @ 20 g a.i ha⁻¹ at 40,55 and 70 DAE + Etherel 2000ppm @ 60% boll burst.

Formatted: Line spacing: 1.5 lines

References

Brar, H.S., Kumar, D and Singh, P. 2020. Dataset of source-sink manipulation through growth retardant for enhancing productivity and profitability of cotton in north west, India. *Data in brief*, 31 :105914.

Comment [H6]: Listed but missing please check

Collins, G.D., Edmisten, K.L., Wells, R. and Whitaker, J.R., 2017. The effects of mepiquat chloride applied to cotton at early bloom and physiological cutout. *Journal of Cotton Science*, 21(3), pp.183-189.

Gomez, K.A. and Gomez, A.A.: Statistical Procedures for Agriculture Research, 2nd Ed. John Willey and Sons, New York, p. 680 (1984).

Kadiyam Priyanka (2021), Performance of Cotton (*Gossypium hirsutum* L.) under HDPS to various plant growth regulators, Msc (Agri) Thesis, ANGRAU, Guntur, Andhra Pradesh.

Comment [H7]: Listed but not cited

Kataria, G.K. and Khanpara, M.D., 2012. Effect of Cycocel and Mepiquat Chloride on physiology, growth and yield of irrigated *Bt* cotton (*Gossypium hirsutum* L.). *International Journal of Scientific Research*, 1(1), pp.90-91.

Comment [H8]: Listed but not cited

[Type here]

Kaul, A., Deol, J.S. and Brar, A.S., 2016. Response of different Bt cotton (*Gossypium hirsutum* L.) hybrids to canopy modification practices. *Journal of Applied and Natural Science*, 8(3), pp.1188-1197.

Khetre, O.S., Shinde, V.S., Asewar, B.V. and Mirza, I.A.B., 2018. Response of growth and yield of Bt cotton to planting densities as influenced by growth regulators. *International Journal of Chemical Studies*, 6(4), pp.485-488.

Kulvir, S., Pankaj, R and Singh, K. 2015b. Dose and time dependent efficacy alteration of different defoliant on seed cotton yield. *Journal of Environmental Biology*. 36: 891-895.

Mrunalini, K., Sree Rekha, M., Murthy, V.R.K and Jayalalitha, K. 2019. Impact of harvest-aid defoliant on yield and economics of high-density cotton. *Indian Journal of Agricultural Research*. 53 (1): 116-119.

Nawalkar, D.P., Ban, Y.G and Kumar, V. 2015. Influence of ethylene and maleic hydrazide on morpho-phenological events and yield in cotton hybrids. *Research in Environment and Life Sciences*. 8(4): 697-700.

Comment [H9]: Listed but not cited

Paslawar, A.N., Meena, A.K., Deotalu, A.S., Bhale, V. M., Ingole, P.G and T.H. Rathod. 2015. Foliar application of mepiquat chloride under high density planting system on different species of cotton. National Symposium on Future Technologies: Indian cotton in the next decades at Acharya Nagarjuna University, Guntur Dec 17-19.

Patel, B.R., Chaudhary, P.P., Chaudhary, M.M. and Reddy, T.V., 2021. Effect of mepiquat chloride on yield attributes, yield and economics of Bt cotton under high density planting system.

Comment [H10]: Listed but not cited

Priyanka, K., Rekha, M.S., Lakshman, K and Rao, C.S. 2021. Influence of plant growth regulators in cotton under HDPS. *The Pharma Innovation Journal*. 10 (7) : 329-331.

Priyanka et al., (2022) Cited but not listed

Rademacher (2000) Cited but not listed

Raghavendra, T. and Reddy, Y.R., 2020. Efficacy of defoliant on yield and fibre quality of American cotton in semi-arid conditions. *Indian Journal of Agricultural Research*, 54(3), pp.404-407.

Reddy, D.N., Baker and Hodges, H.F. 1990. Temperature and mepiquat chloride effects on cotton canopy architecture. *Agronomy Journal*. 82: 190-195

Rosolem, C.A., Oosterhuis, D.M. and Souza, F.S.D., 2013. Cotton response to mepiquat chloride and temperature. *Scientia Agricola*, 70, pp.82-87.

Singh, K., Singh, H.P., Rathore, P., Singh, K. and Mishra, S.K., 2017. Manipulations of source sink relationships through mepiquat chloride for enhancing cotton productivity and monetary returns in north western India. *J. Cotton Res. Dev*, 31(1), pp.62-68.

Snipes, C.E., and C.C. Baskin. 1994. Influence of early defoliation on cotton yield, seed quality, and fibre properties. *Field Crop Research*. 37:137- 143.

Comment [H11]: Listed but not cited check

[Type here]

Pravanthi (2020), Studies on defoliants in High density planting cotton- green gram sequence , PhD thesis, ANGRAU, Guntur, Andhra Pradesh.

Comment [H12]: Listed but not cited

Pravanthi, S., Rekha, M.S., Venkateswarlu, B., Rao, C.S. and Jayalalitha, K., 2022. Effect of defoliants on percent defoliation and yield of American cotton (*Gossypium hirsutum*). *Research on Crops*, 23(2), pp.458-465.

Subbiah, B.V. and Asija, G.L. 1956. A rapid procedure for the determination of available nitrogen in soil. *Current Science*. 25: 259-260.

Comment [H13]: Listed but not cited

Stuart *et al.*, 1984 Cited but not listed

Suttle, J.C. 1985. Involvement of ethylene in the action of the cotton defoliant thidiazuron. *Plant Physiology*. 78: 272 – 276.

Uma Maheswari, M and Murali Krishnasamy, S. 2019. Effect of crop geometries and plant growth retardants on physiological growth parameters in machine sown cotton. *Journal of Pharmacognosy and Phytochemistry*. 8(2): 541-545.

Zaman, I., Ali, M., Shahzad, K., Tahir, M.S., Matloob, A., Ahmad, W., Alamri, S., Khurshid, M.R., Qureshi, M.M., Wasaya, A. and Baig, K.S., 2021. Effect of plant spacings on growth, physiology, yield and fiber quality attributes of cotton genotypes under nitrogen fertilization. *Agronomy*, 11(12), 2589.

[Type here]