

Integrated Disease Management (IDM) Modules against karnal bunt (*Tilletia indica*) of wheat

Abstract

Wheat, (*Triticum aestivum* L.) belonging to family “Gramineae” and genus “*Triticum*”, is one of the world’s most widely cultivated food grain crop, due to its wider adaptability to different agro-climatic and soil conditions. Karnal bunt (*Tilletia indica*) is an important wheat disease with implications for wheat grain quality and inflicts changes in chemical composition of infected grains. IDM modules evaluated under pot and field condition revealed that all the thirteen modules were significantly effective and observed lower disease incidence of karnal bunt. Module M₁₁ and M₈ recorded nil (0.00%) disease incidence in both condition. The maximum incidence was recorded (0.425%) and (0.81%) in M₁₂ followed by M₅ (0.395%) and (0.68%) whereas minimum incidence was observed in M₉ (0.002%) and (0.05%) under pot and field respectively. However, in the field maximum yield (44.65 q/ha) and test weight (36.04 g) was recorded in module M₁₁, followed by (44.30q/ha) and (35.91g) in M₈, while, minimum yield (40.35q/ha) and test weight (35.30g) was observed in M₁₂.

Keywords: Karnal bunt, *Tilletia indica* (give space e.g. *Tilletia indica*), Incidence, Module, Treatments

Introduction

Wheat, (*Triticum aestivum* L.) belonging to family “Gramineae” and genus “*Triticum*”, is one of the world’s most widely cultivated food grain crop, due to its wider adaptability to different agro-climatic and soil conditions. Globally, the nutrient rich cereals are grown altogether in 215.48 million hectare area with the annual production 731.40 million metric tons. In (give space) India, during 2020-21, wheat has been cultivated in 30.22 million hectare and production has made another landmark achievement by producing 99.9 million metric tons with an average national productivity of 33.71 q/ha. (Anonymous, 2021). Many (give space) factors of biotic and abiotic stresses pose serious threats to sustain production, productivity and quality of wheat in Indian subcontinent and through-out the world (Chatrath *et al.*, 2007). Out of these different biotic stresses, karnal bunt (*Tilletia indica*) is an important wheat disease with implications for wheat grain quality and inflicts changes in chemical composition of infected grains (Aujla *et al.*, 1980). Karnal bunt is also a disease of quarantine interest and it affects the international trade of commercial wheat grain and movement of wheat throughout the world. With the advent of new stringent laws in import/export, there is zero tolerance limits on shipment of wheat from karnal bunt prone regions (Butler, 1990). The (give space) loss due to Karnal Bunt is difficult to estimate because the disease reduces seed quality, inflicts changes in the chemical composition of infected grains and renders seed useless for consumption. Nevertheless, in India, loss in yield due to the disease has been calculated as 1/3 x yield x per cent infection as the disease covers one third of an area under wheat cultivation in India (Munjal, 1975). The financial losses caused by the disease are substantial, ranging from 5-20%. The losses in grain quality due to Karnal bunt have serious economic repercussions, even though the losses in grain yield are minimal. The economic importance of this disease is generally not measured by the loss in grain quantity but by grain quality which gets deteriorated due to *Tilletia indica* (Sharma *et al.* 2021).

Methods and Materials

An experimental trial (field and pot) for the evaluation of different (give space) Integrated Disease Management (IDM) Modules (treatments) against (give space) karnal bunt of wheat was conducted at Crop Research Center Chiraudi (CRC), Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut during the year 2021 and 2022. The (give space) trial was conducted in 4.0x3.0 m² plots under irrigated condition with recommended package and practices. The experiment was laid out in a Randomized Block Design (RBD) with 3 replications and 13 treatments (Modules) using (give space) karnal bunt susceptible variety HD-2967. When crop reached to boot leaf stage each of the ten heads (give space) in each plot was (give space) inoculated (4-5 times at one day intervals) with 3 ml of sporidial suspension of *T. indica* (10,000 spores/ml). Sterilized water sprayed in sub-plots served as control. After 48 hours of artificial inoculation, each (give space) fungicides was sprayed at their recommended dosage rates at two crop stages first at boot stage and second at ear emergence stage (Table 1). Inoculated plants were tagged and labeled and field moisture was maintained. At maturity, the inoculated heads were harvested, hand threshed and percent seed infections for each module was determined. The data collected included the numbers of infected and uninfected seeds every ten spikes to evaluate the incidence of the disease (% seeds infected). The incidence of the disease for each entry was calculated by using the following formula Aujla *et al.* (1989).

$$\text{Percent disease incidence} = \frac{\text{Number of bunted grains in 10 spike}}{\text{Total number of grain in 10 spike}} \times 100$$

And Percent Disease control was calculated with the help of the following formula:

$$\text{Percent disease control} = \frac{\text{PDI in control} - \text{PDI in treated}}{\text{PDI in control}} \times 100$$

Whereas, PDI = Percent Disease Incidence



Fig 1. 1. Culture of *Tilletia indica* in PDB, 2. Artificial inoculation of spore suspension, 3. Fungicide spray on experimental field and 4. Infected grains of wheat

Table 1: Details of the IDM modules:

Module	Seed Treatment	Soil Treatment	Foliar Spray	
M ₁	Hot water (52 ⁰ C) for 10 min.	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Propiconazole 25% EC @ 0.1% at ear emergence	
M ₂	Thiram 75% DS @ 2.5 g/Kg seed	<i>Bacillus subtilis</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Tebuconazole 25.9% EC @ 0.1% at ear emergence	
M ₃	Late sowing	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Pressmud @ 10t/ha.	Azoxytrobins 23% SC @ 0.1% at ear emergence	
M ₄	Hot water (52 ⁰ C) for 10 min.	<i>Bacillus subtilis</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Pressmud @ 10t/ha.	Tebuconazole 25.9% EC @ 0.1% at ear emergence	
M ₅	Salt (NaCl) @ 20%	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Azadirachtin (Neem Oil) @ 0.3% at ear emergence	
M ₆	Thiram 75% DS @ 2.5 g/Kg seed	<i>Bacillus subtilis</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Azadirachtin (Neem Oil) @ 0.3% at ear emergence	
M ₇	Hot water (52 ⁰ C) for 10 min.	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Pressmud @ 10t/ha.	Azadirachtin (Neem Oil) @ 0.3% at ear emergence	
M ₈	Salt (NaCl) @ 20%	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Bootstage	Ear emergence
			Propiconazole 25% EC @ 0.1%	Azadirachtin (Neem Oil) @ 0.3%
M ₉	Thiram 75% DS @ 2.5 g/Kg seed	<i>Bacillus subtilis</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Tebuconazole 25.9% EC @ 0.1%	Azadirachtin (Neem Oil) @ 0.3%
M ₁₀	Hot water (52 ⁰ C) for 10 min.	<i>Pseudomonas fluorescens</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Azoxytrobins 23% SC @ 0.1%	Azadirachtin (Neem Oil) @ 0.3%
M ₁₁	Thiram 75% DS @ 2.5 g/Kg seed	<i>Bacillus subtilis</i> (cfu 2×10 ⁸ /g) @ 5Kg/ha. + Vermicompost @ 10t/ha.	Propiconazole 20% EC @ 0.1%	Azoxytrobins 23% SC @ 0.1%
M ₁₂	Late sowing without treatment			
M ₁₃	Control	Control	Control	

Results and Discussion

IDM modules evaluated under pot condition revealed that all the thirteen modules were significantly effective and observed lower disease incidence of karnal bunt (Table 2). Module (give space) M₁₁ [seed treatment with Thiram 75% DS @ 2.5 g/kg seed, soil treatment with *Bacillus subtilis* (CFU 2×10⁸/g) @ 5Kg/ha. + Vermicompost @

10t/ha. and Propiconazole 20% EC @ 0.1% booting (give space) stage + Azoxystrobin 23% SC @ 0.1% at ear emergence stage] and M₈[seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha and Propiconazole 25% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage) recorded nil (0.00%) disease incidence. After control the maximum incidence was recorded (0.425%) in M₁₂[late sowing] followed by M₅(0.395%) [Seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha + Vermicompost @ 10t/ha. and foliar spray with Azadirachtin (Neem Oil) @ 0.3% at ear emergence] whereas minimum incidence was observed in M₉ (0.001%). [seed (give space) treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha + Vermicompost @ 10t/ha and foliar spray with].

Table 2: Evaluation of IDM modules against *T. indica* (Pot)

Modules	Disease Incidence (2021)	Disease Incidence (2022)	Plant Disease over Control (2021)	Plant Disease over Control (2022)	Average DI (2021-22)
M ₁	0.026	0.028	94.90	94.71	0.027
M ₂	0.019	0.022	96.27	95.84	0.020
M ₃	0.210	0.200	58.82	62.26	0.205
M ₄	0.090	0.110	80.39	79.24	0.100
M ₅	0.390	0.400	23.52	24.52	0.395
M ₆	0.250	0.270	50.98	49.05	0.260
M ₇	0.310	0.300	39.22	43.39	0.305
M ₈	0.000	0.000	100	100	0.000
M ₉	0.000	0.005	100	99.05	0.002
M ₁₀	0.150	0.140	70.58	73.58	0.145
M ₁₁	0.000	0.000	100	100	0.000
M ₁₂	0.430	0.420	15.68	20.75	0.425
M ₁₃	0.510	0.530	0	0	0.520
CD at 5%	0.03	0.03			

Studies on management of karnal bunt were conducted to evaluate thirteen IDM modules, under field condition revealed that all the modules were significantly effective and recorded lower disease incidence of karnal bunt (Table 3). However, under natural condition module M₁₁[seed treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha. and Propiconazole 20% EC @ 0.1% booting stage+ Azoxystrobin 23% SC @ 0.1% at ear emergence stage] and M₈[seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha. and Propiconazole 25% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage] observed nil (0.00%) incidence followed by M₉ [seed treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha + Vermicompost @ 10t/ha. and foliar spray with Tebuconazole 25.9% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage] (0.005%) was recorded. The other modules were recorded higher disease incidence viz., M₁₂ (0.81%) [Late sowing], M₅ (0.68%) [Seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha + Vermicompost @ 10t/ha. and foliar spray with Azadirachtin (Neem Oil) @ 0.3% at ear emergence] and M₇ (0.60%) [seed treatment with Hot water (52°C) for 10 min., soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha + Pressmud @ 10t/ha and foliar spray with Azadirachtin (Neem Oil) @ 0.3% at ear emergence].

Table 3: Evaluation of IDM Modules against karnal bunt of wheat (Experimental Field)

Module	Disease Incidence 2021	Plant Disease over control (2021)	Disease Incidence 2022	Plant Disease over control (2022)	Average Disease Incidence
M ₁	0.15	83.14	0.17	80.46	0.16
M ₂	0.09	89.88	0.11	87.35	0.10
M ₃	0.47	47.19	0.45	48.27	0.46
M ₄	0.21	76.40	0.20	77.01	0.20
M ₅	0.67	24.71	0.60	20.68	0.68
M ₆	0.53	40.45	0.56	35.63	0.54

M ₇	0.59	33.71	0.61	29.88	0.60
M ₈	0.00	100	0.00	100	0.00
M ₉	0.04	95.50	0.06	93.10	0.05
M ₁₀	0.33	62.92	0.35	59.77	0.34
M ₁₁	0.00	100	0.00	100	0.00
M ₁₂	0.83	6.74	0.79	9.19	0.81
M ₁₃	0.89	0	0.87	0	0.88
CD at 5%	0.06	0.06			

Data from table 4 also revealed that all the modules were significantly effective and recorded higher yield and test weight compared to unprotected control. However, the maximum yield and test weight was recorded (44.65 q/ha) and (40.59 g) in module M₁₁ [seed treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha. and Propiconazole 20% EC @ 0.1% booting stage+ Azoxystrobin 23% SC @ 0.1% at ear emergence stage] followed by (44.30 q/ha) and (40.38 g) M₈ [seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha and Propiconazole 25% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage], after control the minimum yield and test weight was observed (40.35q/ha) and (37.50 g) in M₁₂ (Late sowing) respectively.

Table 4: Effect of IDM modules on yield (q/ha) and 1000 grain weight (test weight) of wheat

Module	Yield (q/ha)		Increase yield over control		Average yield (2021-22)	1000 grain weight (gram)		
	2021	2022	2021	2022		2021	2022	Ave.
M ₁	43.60	43.20	9.00	7.73	43.40	39.92	39.68	39.80
M ₂	43.90	43.70	9.75	8.98	43.80	40.09	39.87	39.98
M ₃	42.30	42.10	5.75	4.98	42.20	39.19	38.76	38.97
M ₄	43.10	42.80	7.75	6.73	42.95	39.86	39.39	39.62
M ₅	40.60	40.50	1.50	0.99	40.55	37.96	37.71	37.83
M ₆	41.90	41.70	4.75	3.99	41.80	38.77	38.43	38.68
M ₇	41.50	41.30	3.75	2.99	41.40	38.23	38.09	38.16
M ₈	44.40	44.20	11.00	10.22	44.30	40.44	40.33	40.38
M ₉	44.10	43.90	10.25	9.47	44.00	40.26	40.11	40.18
M ₁₀	42.70	42.50	6.75	5.98	42.60	39.63	39.17	39.40
M ₁₁	44.80	44.50	12.00	10.97	44.65	40.67	40.52	40.59
M ₁₂	40.30	40.40	0.75	0.75	40.35	37.56	37.47	37.50
M ₁₃	40.00	40.10	0	0	40.05	37.10	37.21	37.15
CD at 5%	5.20	5.18				1.78	1.72	

DI= Disease Incidence, Ave. = Average

Karnal bunt pathogen is predominantly soil-borne rather than seed-borne like the dwarf bunt; seed treatment with fungicide is ineffective for the total eradication of infection (Hoffmann, 1982; Brooks and Buckley, 1977). However, it might reduce the likelihood of infection (El-Naimi *et al.*, 2000). The biggest issue with utilize existing fungicides is that when the chemical washes off the spore, karnal bunt spores may develop (Sharma & Basandrai, 2000).

Foliar sprays of Propiconazole, Tebuconazole, Hexaconazole, Thifluzamide, Diniconazole etc. were shown to be effective against natural infection in India (Singh *et al.*, 1999).

All the fungicides singly or in combinations significantly increased the disease control and yield increased as compared to control. Efficacy of such combination products in managing many fungal diseases has been reported by various workers across the world (Wanyera *et al.*, 2009; Muniraju *et al.*, 2017). Different fungicides vary in their efficacy to control karnal bunt. Nagy and Moldovan (2006) reported that fungicides containing Difenconazole (Dividend 030 FS 1.0 l/t), Tebuconazole (Raxil 060 FS 0.5 l/t), Fludioxonil + Epoxiconazole (Maxim Star DS 1.5 kg/t), Tebuconazole + Thiram (Raxil T 515 FS 2.0 l/t) had a very good efficiency in controlling the common bunt even under artificial infections. Singh *et al.* (2000) compared the efficacy of different fungicides found that the maximum disease control (99.8%) was achieved Propiconazole (give space) (0.1) by a single spray controlled (96.46%) disease, followed by Hexaconazole (92.87%) in the post-inoculation treatment. Singh *et al.* (2018) evaluated the efficacy of seven fungicides viz; Tilt 250, Folicur, Bavistin, Thiram, Vitavax, Benlate and Dithane M-

75 of different groups against *Tilletia indica* and reported that out of seven, two fungicides viz; Tilt 250 EC and Folicur to be the most effective as they have inhibited the fungal growth completely.

CONCLUSION

Among all the 13 modules (treatments) evaluated at field conditions, the best module was found to be effective, against karnal bunt of wheat was module M₁₁ [seed treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha. and Propiconazole 20% EC @ 0.1% booting stage+ Azoxystrobin 23% SC @ 0.1% at ear emergence stage] and M₈ [seed treatment with salt (NaCl) @ 20%, soil treatment with *Pseudomonas fluorescens* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha. and Propiconazole 25% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage] observed nil incidence (0.00%), whereas, minimum disease incidence was observed in M₉ (0.005%) [Seed treatment with Thiram 75% DS @ 2.5 g/Kg seed, soil treatment with *Bacillus subtilis* (cfu 2×10⁸/g) @ 5Kg/ha. + Vermicompost @ 10t/ha and foliar spray with Tebuconazole 25.9% EC @ 0.1% at booting stage and Azadirachtin (Neem Oil) @ 0.3% at ear emergence stage]. Similar results were also recorded with test weight (40.59g) and yield (44.65q/ha) in module M₁₁ and M₈ (40.38g & 44.30q/ha).

FUTURE SCOPE

The fungus *Tilletia indica* inciting Karnal bunt disease of wheat is a serious concern for import of wheat to Karnal bunt free countries. The grain quality deterioration caused by Karnal bunt fungus is known to have serious implications in the world wheat trade due to strict quarantine regulations. Understanding pathogen population is an important aspect for exploiting resistance, especially when dealing with a heterothallic fungus. Several diagnostic techniques have been devised for the accurate identification of the fungus. This seed, soil and air borne pathogen have less chemical control measures so an integrated disease management strategy is the best approach to combating the disease.

Reference

- Anonymous (2021). Economic survey of India.
- Aujla, S. S., Grewal, A. S., Gill, K. S. and Sharma, I. 1980. A screening technique for karnal bunt disease of wheat. *Crop Improvement*, 7(2): 145-146.
- Aujla, S. S., Sharma, I. and Singh, B. B. 1989. Rating scale for identifying wheat varieties resistant to *Neovossia indica*. *Indian Phytopathology*, 42: 161-162.
- Butler, L. 1990. Karnal bunt, quarantine and the international shipment of CIMMYT wheat seed. Proceedings Workshop on Smut Fungi, 7th.
- Brooks, D. H. and Buckley, N. G. 1977. Results in practice -1. *Cereals and grasses*. Pages 213-233 in: Systemic Fungicides. Marsh, R. W., ed. Longman, New York. 401pp.
- Chatrath, R., Mishra, B., Ortiz-Ferrara, G., Singh, S. K. and Joshi, A. K. 2007. Challenges to wheat production in South Asia. *Euphytica*, 157: 447-456.
- El-Naimi, M., Toubia-Rahme, H. and Mamluk, O. F. 2000. Organic seed treatment as a substitute for chemical seed treatment to control common bunt of wheat. *European Journal of Plant Pathology*, 106: 433-437.
- Hoffmann J. A. 1982. Bunt of wheat. *Plant Disease*, 66: 979-986.
- Khan, M. A. and M. B. Ilyas. 2007. Chemotherapy of Plant Diseases. (Laboratory Manual). Department of Plant Pathology University of Agriculture Faisalabad Pakistan.
- Munjal, R.L. (1975). Status of Karnal bunt (*N. indica*) of wheat in North India during 1968-69 to 1969-70. *Indian Journal of Mycology and Plant Pathology*, 5, 185-187.
- Muniraju, K. M., Pramesh, D., Malleth, S. B., Mallikarjun, K. and Guruprasad, G. S. 2017. Novel fungicides for the management of false smut disease of rice caused by *Ustilaginoidea virens*. *International Journal of Current Microbiology and Applied Sciences*, 6(11): 2664-2669.
- Nagy, E. and Moldovan, V. 2006. The effect on fungicides treatments on the wheat common bunt (*Tilletia* spp.) in Transylvania-Romania. *Czech Journal of Genetics and Plant Breeding*, 42: 56-61.
- Sharma, B. K. and Basandrai, A. K. 2000. Effectiveness of some fungicides and biocontrol agents for the management of karnal bunt of wheat. *Indian Journal of Mycology and Plant Pathology*, 30: 76-78.
- Singh, H., Grewal, T. S., Pannu, P. P. S. and Dhaliwal, H. S. 1999. Genetic of resistance to karnal bunt disease of wheat. *Euphytica* 105: 125-131.
- Singh, R., Beniwal, M. S. and Karwasra, S. S. 2000. Field evaluation of fungicides against karnal bunt of wheat through foliar spray. Department of Plant Pathology, Chaudhary Charan Singh Haryana Agricultural University, Hisar - 125 004, India. : Wellesbourne, UK: Association of Applied Biologists. *Tests of Agrochemicals and Cultivars*, 21: 9-10.
- Kumar S., Singroha G., Singh G. P. and Sharma P. 2021. Karnal bunt of wheat: Etiology, breeding and integrated management *Crop Protection*, 139: 105376
- Wanyera, R., Macharia, J.K., Kilonzo, S.M. and Kamundia, J.W. 2009. Foliar fungicides to control wheat stem rust, race TTKS (Ug99), in Kenya. *Plant Disease*, 93: 929-932.