

Original Research Article

Analysis of Parent Offspring Regression of Selected Cowpea (*Vigna unguiculata* [L. Walp.) Agronomic Traits in Phosphorus Limiting Soil.

Comment [a1]: Remove toggle key as per do it as per Journal style

ABSTRACT

Cowpea is a significant grain legume pulse, vegetable, and fodder crop. However, due to biotic and abiotic stresses, yields have been consistently low. Phosphorus (P) deficiency-related yield losses are one of the major abiotic stresses. The objectives therefore were i) to evaluate the mean performance response of selected cowpea traits across genotypes in optimum and P- limiting soil media and ii) to determine the cowpea heritability response of agronomic traits in P-limiting soil. Twenty cowpea genotypes comprising of 12 F₁ crosses and eight progenitors were evaluated at the University of Zambia Greenhouse in Lusaka, using a completely randomized design with three replications and two-level sets of phosphorus (i.e. 0 kg P₂O₅ and 60 kg P₂O₅). The 60 kg P₂O₅ level set was used as a control. The results showed that the mean performance across genotypes for all measured traits was higher in control experiment at an applied rate of 60 kg P₂O₅/ha than at applied rate of 0 Kg P₂O₅/ha experimental sets. Significant better performances were obtained with plant height, shoot biomass, plant biomass and yield measured traits. The narrow sense heritability for measured traits ranged from 11- 35 % and 12 - 72% for 0 Kg P₂O₅ and 60 Kg P₂O₅ experimental set respectively. The highest narrow sense heritability scores in experimental set were pod length ($h^2= 0.35$) and root biomass ($h^2= 0.28$) while in the control set the variable yield ($h^2= 0.72$) had the highest score. This implies that the identified trait (pod length) with high narrow sense heritability in P-limiting soil can initially be used to aid in selecting for high performing genotypes in P limiting soil. However, this should be supplemented by yield response especially in late generation selection (F₅ onwards).

Comment [a2]: remove no. ...keep sentence in one line

Comment [a3]: remove no. ...keep sentence in one line

Comment [a4]: italic

Keywords: *Parent offspring regression, Phosphorus, Narrow sense and *Vigna unguiculata**

Comment [a5]: full botanical name

INTRODUCTION

Cowpea [*Vigna unguiculata* (L.) Walp] is an essential economic crop, native to Africa and belongs to the family Fabaceae. It is rich in diverse nutrients, with high levels of protein, vitamins, fiber, macro and micronutrients (i.e. protein, carbohydrates, vitamins, and minerals) [1,2,3,4]. Cowpea is multifunctional future smart climate-resilient crop contributing to food and nutritional security to resource poor farmers. Importantly, cowpea forms symbiosis with the root nodule bacterium, *Rhizobium*, and fixes 70 to 350 kg/ ha of atmospheric nitrogen and some 40 to 80 kg of this is deposited into soils as a natural source of mineral nitrogen, which contribute to soil health [5].

Comment [a6]: italic

Global production of cowpea is estimated to be 7.4 million tons per annum on a 12.5 million hectare land [6]. The mean grain yield of cowpea in sub-Saharan Africa is less than 700 kg/ ha which is far less than the potential yield of 3000 kg/ha [7,8]. The yield gap is attributable to a multitude of biotic and abiotic stresses. Among the abiotic stresses, yield losses due to phosphorus (P) deficiency is an important factor. Phosphorus is one of the most important plant nutrients, accounting for 0.05 % to 0.5 % of plant dry weight [9]. However, P is one of the least available plant nutrients and is deficient in many soils of the world [10]. Presently, the use of P-formulated fertilizers by some farmers appears to be a quick and easy fix for P deficient soils. However, the current option has not been widely adopted by most smallholder cowpea growers because it is expensive, neither sustainable nor environmentally safe. Therefore, the most sustainable strategy is to develop and choose cowpea genotypes that have high yield in P-limiting soil and are easily adopted by low-resource farmers.

Comment [a7]: loss...not losses

Knowledge on narrow sense heritability of the agronomic variables is key in understanding a breeding strategy to employ in screening for cowpea genotypes in P-limiting soils. As an indicator of selection effectiveness and a measure of the likelihood that advantageous additive genes will be passed down from progenitor to their progeny,

information on narrow-sense heritability is crucial to breeders [11]. It is important because of the effectiveness of selection is dependent on the additive components of genetic relative to total variance [12,13]. However, heritability estimates are usually specific to the trait, population, and environment being investigated [14]. This allows breeders to develop the best breeding techniques and selection criteria for desirable trait. This research endeavors to generate fundamental knowledge for utilization in breeding cowpea varieties efficient at utilizing P in P-limiting soils. Therefore, the objectives of this study were i) to evaluate the mean performance response of selected cowpea traits across genotypes in optimum and P- limiting soil media, and ii) to determine the cowpea heritability response of agronomic traits in P-limiting soil.

Comment [a8]: remove numbers

MATERIALS AND METHODS

Experimental site, material used and experimental details.

Twenty genotypes comprising of 12 F1 crosses and eight progenitors were used for this study. The F₁ crosses were generated as by Chikalipa *et al.* [15] (**Table 1**). The study was conducted at the University of Zambia, Green house in Lusaka (15°23'S and 28°25'E, at 1250 m above sea level). Soil samples used in the study were collected from the top layer (0–30 cm) at Liempe Farm (15° 22' S and 28° 26' E, at 1171m above sea level) in Chongwe District, as described by [15]. It contained 7.30 mg/kg available phosphorus, 1.92 % organic matter, 0.34 % nitrogen (N) and 0.28 cmol/kg potassium (K). The average pH was measured to be 5.25.

Table 1. Genotype used in the study

Genotypes Codes	Genotypes	Type
AXH	<u>LT 3-8-4-6</u> X <u>LT 11-3-3-13</u>	Cross
BXG	<u>LT 11-5-1-1</u> X <u>LT 10-7-1-12</u>	Cross
AXD	<u>LT 3-8-4-6</u> X <u>LT 4-2-4-1</u>	Cross
CXE	<u>LT 11-5-2-2</u> X <u>LT 3-8-4-1</u>	Cross
DXA	<u>LT 4-2-4-1</u> X <u>LT 3-8-4-6</u>	Cross
EXB	<u>LT 3-8-4-1</u> X <u>LT 11-5-1-1</u>	Cross
FXE	<u>LT 16-7-2-5</u> X <u>LT 3-8-4-1</u>	Cross
AXE	<u>LT 3-8-4-6</u> X <u>LT 3-8-4-1</u>	Cross
GXC	<u>LT 10-7-1-12</u> X <u>LT 11-5-2-2</u>	Cross

CXG	<u>LT 11-5-2-2 X LT 10-7-1-12</u>	Cross
EXD	<u>LT 3-8-4-1 X LT 4-2-4-1</u>	Cross
HXA	<u>LT 11-3-3-13 X LT 3-8-4-6</u>	Cross
A	LT 3-8-4-6	Parent
B	LT 11-5-1-1	Parent
C	LT 11-5-2-2	Parent
D	LT 4-2-4-1	Parent
E	LT 3-8-4-1	Parent
F	LT 16-7-2-5	Parent
G	LT 10-7-1-12	Parent
H	LT 11-3-3-13	Parent

The study was a two-set experiment and laid out as a completely randomized design (CRD), with three replications. The experimental plot was 15cm diameter polythene plastic bags filled with 3kg of soil and Single Super Phosphate (SSP) was applied three weeks after planting. The quantity of SSP added was calculated to achieve the soil P at two levels 60kg P₂O₅ /ha (optimal) was added to control set while 0kg P₂O₅/ha (suboptimal) was added to experimental set. However, the recommended quantities of potassium (K) (30 kg K/ha) and nitrogen (N) (20 kg N/ha) was applied as basal dressing in form of potassium sulphate and urea (46% N) fertilisers at the rates of 64 kg/ha and 25 kg/ha respectively. In each bag, three identical genotypes were planted which were later thinned to two plants per bag at the true-leaf stage. Other crop management practices were carried out as per the best agronomic practices recommended for cowpea production.

Data collection

The following variables were collected on each experimental unit in both sets; Plant height(cm); Shoot biomass(g); Root biomass(g); Plant biomass(g); 100 Seed weight (g) was calculated in grams from hundred sampled seeds; Pod Length (cm); Root length (cm); Number of pods per plant were counted at harvest maturity; Number of seeds per pod were counted at harvest maturity; Yield/ha; Yield per hectare was estimated on a plot basis and converted the plot yield of the seed dry weight to kg/ha.

Comment [a9]: check fount spacing before (g)

Comment [a10]: yield per ha

Comment [a11]: remove

Comment [a12]: (kg/ha)

Statistical Data Analysis

Heritability in the narrow sense was estimated by regressing mid-parent means on the progeny means as suggested by Falconer and Mackay [11]. The equation being;

$$Y = bX + C$$

Where; **Y** is the average performance of progeny, **b** is the gradient representing narrow sense heritability estimate, **X** is the average performance of mid-parent, and **C** is the constant.

Mean genotypic performance for each measured variable in each experimental set was recorded and the standard error bars were computed in excel. Data analysis was performed using GenStat (18th edition) and excel statistical software.

RESULTS AND DISCUSSION

Genetic evaluation of traits

The results showed that the mean performance across genotypes for all measured traits was higher in control experiment at an applied rate of 60kg P₂O₅/ha than at applied rate of 0Kg P₂O₅/ha (Figure 1). Significant better performances were obtained with plant height, shoot biomass, plant biomass and yield measured traits. This implies that Phosphorous plays a key role in plant productivity, cowpea inclusive. Previous studies have demonstrated that phosphorous is an important element in leguminous crop [16,17]. Furthermore, mean genotypic responses of measured variables exhibited varying narrow sense heritability (h²) ranging from 11- 35 % and 12-72 % for 0 Kg P₂O₅ experimental trial and 60 Kg P₂O₅ control trial respectively across measured traits (Table 2).

Comment [a13]: space

Comment [a14]: space

Comment [a15]: remove

Comment [a16]: plant productivity in cowpea, inclusively.

Comment [a17]: remove




Figure 1. Mean performance of measured traits at P- limited and optimum medium PH=Plant Height (cm), NDP= Number of Pods per Plant, PDL= Pod Length (cm), NSPD= Number of Seed per Pod, 100SW=Hundred seed weight, RBM =Root Biomass, PBM= Plant Biomass (g), SBM=Shoot biomass, Yield*= Yield Kg x 10²/ ha.

Table 2. Narrow sense heritability of measured variables

Measured variables	h^2 0Kg P ₂ O ₅	h^2 60Kg P ₂ O ₅
PH	0.19	0.12
NDP	0.13	0.22
PDL	0.35	0.16
NSPD	0.11	0.46
100SW	0.12	0.24
RBM	0.28	0.16
PBM	0.12	0.38
SBM	0.13	0.37

Yield	0.11	0.72
-------	------	------

h²=Heritability, **PH**=Plant Height, **NDP**= Number of Pods per Plant, **PDL**= Pod Length, **NSPD**= Number of Seed per Pod, **100SW**=Hundred seed weight, **RBM** =Root Biomass, **SBM**=Shoot biomass, **PBM**= Plant Biomass, **Yield**= Yield Kg/ha.

The narrow sense heritability for agronomic traits ranged from 11-35 % and 12-72% for 0 Kg P₂O₅ and 60 Kg P₂O₅ respectively (**Table 2**). Narrow sense heritability estimates in crops are classified as high (>0.20), medium (0.10 - 0.20), and low (<0.10) [18]. The narrow-sense heritability observed in the experimental trial, suggest that there is potential for improvement through progeny selection and advancement [14].

Comment [a18]:comma

Estimates for narrow sense heritability obtained are similar to reported narrow sense heritability ranging from 20-85% in cowpea [19, 20]. In the experimental set (0 Kg/ ha P₂O₅) variables, pod length ($h^2 = 0.35$) and root biomass ($h^2 = 0.28$) had the highest heritability values while in the control set, the variable yield ($h^2 = 0.72$) had the highest value. The results entail that selection for yield in P-limiting medium may be challenging for breeders especially if applied in early generation selection and efforts may be boosted through applying genomics or marker-assisted breeding [21, 22, 23]. Responses to pod length (PDL) and root biomass (RBM) could be used as an indirect selection criteria for advancing cowpea crosses in P-limiting soil especially at early generation selection. Early generation selection can be applied to traits that have high heritability. However, for yield ($h^2 = 0.11$) under P- limited soil, late generation selection (F₅ onwards) screening should be employed as heritability for yield was found to be low. Previous studies have shown that PDL and RBM can be used as an indirect selection criteria for dry bean agreeing with our finding [24]. The higher yield heritability value ($h^2 = 0.72$) in control set (**Table 2**) could be due to sufficient P which permitted full expression of genes associated with yield [18].

Comment [a19]: remove an

Comment [a20]: delete

CONCLUSION

In this study, results showed that the mean performance across genotypes for all measured traits was higher in optimum than the limiting soil medium. Significant better performances were obtained with plant height, shoot biomass and plant biomass traits. The narrow sense heritability for measured traits ranged from 11- 35 % and 12-72% for 0 Kg P₂O₅ and 60 Kg P₂O₅ experimental set respectively. The highest narrow sense heritability scores in a limiting P soil medium were for pod length ($h^2= 0.35$) and root biomass ($h^2= 0.28$) while in the control set (optimum soil medium) the yield trait ($h^2= 0.72$) had the highest score. This implies that the identified variable (pod length) with high narrow sense heritability in P-limiting soil can initially be used to aid in selecting for high performing genotypes in P limiting soil during early generation selection. However, this should be supplemented by yield response especially at late generation selection (non-segregating generations), F₅ onwards.

Comment [a21]: set,

Comment [a22]: selection in early generation

Comment [a23]: F₅do it in complete article

Comment [a24]: do stricly as per Journal format

REFERENCES

1. Bai Z, Huang X, Meng J, Kan L, Nie S. A comparative study on nutritive peculiarities of 24 Chinese cowpea cultivars. Food chemistry toxicology journal, 2020; 146: 111- 841.
2. ElMasry G, Mandour N, Ejez Y, Demilly D, Al-Rejaie S, Verdier J, Belin E, Rousseau D. Multichannel imaging for monitoring chemical composition and germination capacity of cowpea (*Vigna unguiculata*) seeds during development and maturation. The Crop Journal, 2021; 10(5): 1399-1411.
3. Silva VM, Nardeli AJ, de Carvalho Mendes NA, de Moura Rocha M, Wilson L, Young SD, Broadley MR, White PJ, Dos Reis AR. Agronomic biofortification of cowpea with zinc: variation in primary metabolism responses and grain nutritional quality among 29 diverse genotypes. Plant Physiology and Biochemistry Journal, 2021;162: 378–387.

4. Tembo L, Pungulani L, Sohati HP, Mataa CJ, Munyinda K. Resistance to *Callosobruchus maculatus* developed via gamma radiation in cowpea. *Journal of Agriculture and Crops*, 2017; 3(8): 65-71.
5. Meena H, Krishna KR, Singh B. Character associations between seed yield and its components traits in cowpea (*Vigna unguiculata* L.) Walp. *Indian Journal Agricultural Research*, 2015; 49(6): 567-70.
6. FAOSTAT. Food and Agriculture data; 2022. Available at <https://www.fao.org/faostat/en/#home> Accessed June 11. 2022.
7. Horn LN, Shimelis H, Laing M. Participatory appraisal of production constraints, preferred traits and farming system of cowpea in the northern Namibia: implications for breeding. *Legume Research – An International Journal*, 2015; 38, 691–700.
8. Gerrano AS, van Rensburg SWJ, Adebola OP. Preliminary evaluation of seed and germination traits in cowpea (*Vigna unguiculata*) genotypes, *South African Journal of Plant and Soil*, 2017;34(5), 399-402.
9. Vance CP, Uhde-Stone C, Allan DL. Phosphorus acquisition and use: critical adaptations by plants for securing a non renewable resource. *New Phytology*, 2003; 157: 423–447.
10. White PJ, Brown PH. Plant nutrition for sustainable development and global health: *Annals of Botany*, 2010; 105(7), 1073-1080.
11. Falconer DS, Mackay TFC. *Introduction to Quantitative Genetics* (4th ed.), Longman Group Ltd. 1996.
12. Acquah G. *Principles of Plant Genetics and Breeding*. Oxford: Wiley- Blackwell. 2007.
13. Bernardo R. *Essentials of Plant Breeding*. Woodbury, USA: Stemma Press. 2014.
14. Horn LN, Shimelis H. Production constraints and breeding approaches for cowpea improvement for drought prone agro-ecologies in Sub-Saharan Africa. *Annals of Agricultural Sciences*, 2020; 65(1), 83-91.

Comment [a25]: remove

15. Chikalipa E, Mwala SM, Tembo L. Evaluation of cowpea parents and hybrids through multivariate analysis under phosphorus limited soil. *Journal of Genetics, Genomics and Plant breeding*, 2022; 6(4) 102-110.
16. Li SX, Wang ZH, Stewart BA. Differences of some leguminous and non-leguminous crops in utilization of soil phosphorus and responses to phosphate fertilizers. *Advances in Agronomy*, 2011; 110, 125-249.
17. Tairo EV, Ndakidemi PA. Possible benefits of rhizobial inoculation and phosphorus supplementation on nutrition, growth and economic sustainability in grain legumes. *American Journal of Research Communication*, 2013;1(12), 532-556.
18. Ajayi A. Genetic Variability of Quantitative Traits in F2 Hybrids of Cowpea and Parent Lines. *Journal of Genetic Resources*, 2023; 9(1), 1-10.
19. Jacinto-Hernandez C, Azpiroz-Rivero S, Acosta-Gallegos, J A, Hernandez-Sanchez H, Bernal-Lugo I. Genetic analysis and random amplified polymorphic DNA markers associated with cooking time in common bean. *Crop Science Society of America*, 2003; 43, 329–332.
20. Mashi DS. Genetic studies on seed coat texture and cooking time in some varieties of cowpea (*Vigna Unguiculata* (L.) Walp) (Doctoral dissertation). 2006.
21. Mbwando A, Lungu DM, Tryphone GM, Tembo L. Nature of resistance of cowpea *Alectra vogelii* infestation. *African Crop Science Journal*, 2016; 24, 389-395.
22. Tan H, Tie M, Luo Q, Zhu Y, Lai J, Li H. A review of molecular markers applied in cowpea (*Vigna unguiculata* L. Walp.) breeding. *Journal of Life Sciences*, 2012; 6(11), 1190.
23. Raina A, Laskar RA, Wani MR, Khan S. Role of Mutation and Molecular Breeding in the Improvement of Cowpea, a Primary Pulse Crop. *Biotechnologies and Genetics in Plant Mutation Breeding: Mechanisms for Genetic Manipulation of Plants and Plant Mutants*. 2023.

24. Cichy KA, Wiesinger JA, Matthew B, Nchimbi-Msolla S, Deidre F, Porch TG. The role of genotype and production environment in determining the cooking time of dry beans (*Phaseolus vulgaris L.*). Legume Science Journal, 2019; 1:e13.

UNDER PEER REVIEW