

Some Fixed Point results of Rational Type-Contraction Mapping in S-Metric Space

Abstract:

In this paper, we demonstrate the existence of some fixed points of rational type contraction in context of S-metric space and we examine the T-stability of the P-property for some mapping. Also, we present few examples to illustrate the validity of the results obtained in the paper.

Keywords: Fixed point, rational type contraction, S-metric space.

MSC: 54H25, 47H10

1. Introduction and Preliminaries

Fixed point theory is an active area of research with various application in real life. One of the main approaches used in this theory to demonstrate the existence and uniqueness of fixed point is contraction. In 1989, Bakhtin [7] was the first who introduced the concept of b-metric space. In 1993, Czerwik [9] extended the results of Bakhtin [10] and gave generalization of Banach fixed point theorem in b-metric spaces. In 2012, the idea of S-metric space was established by Sedghi *et al.* [21], who also proved fixed point theorems there in. More well-known results in the direction of S-metric space are involved in (refer [22]-[25]).

Furthermore, we proceed by reviewing some important definitions and key terms that would be used throughout our discussion.

Definition 1.1 [21]: “Let \mathcal{X} be a non-empty set. An S-metric on \mathcal{X} is a mapping $\mathcal{S}: \mathcal{X} \times \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}^+$ which satisfies the following condition:

(\mathcal{S}_1) $\mathcal{S}(u, v, w) = 0$ if and only if $u = v = w = 0$;

(\mathcal{S}_2) $\mathcal{S}(u, v, w) \leq \mathcal{S}(u, u, a) + \mathcal{S}(v, v, a) + \mathcal{S}(w, w, a)$, for all $u, v, w, a \in \mathcal{X}$.

The pair $(\mathcal{X}, \mathcal{S})$ is called an S-metric space.”

Example 1.2 [21]: “Let $\mathcal{X} = \mathbb{R}$. Then $\mathcal{S}(u, v, w)$ is an S-metric on \mathbb{R} given by $\mathcal{S}(u, v, w) = |u - w| + |v - w|$, which is known as usual S-metric space on \mathcal{X} .”

Manoj K. *et al.* [18], proved fixed point theorem by using altering distance function in S-metric space.

Theorem 1.3 [18]: “Let $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping on a complete S-metric space $(\mathcal{X}, \mathcal{S})$ such that

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq \lambda \mathcal{S}(u, u, v) + \eta \frac{\mathcal{S}(u, u, \mathcal{T}v)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)},$$

for all $u, v \in \mathcal{X}$, $\lambda, \eta > 0$, $\lambda + \eta < 1$. Then \mathcal{T} possess a fixed point $w \in \mathcal{X}$ which is unique.”

Theorem 1.4 [18]: “Let $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping on a complete S-metric space $(\mathcal{X}, \mathcal{S})$ such that

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq \lambda \mathcal{S}(u, u, v) + \eta \frac{\mathcal{S}(v, v, \mathcal{T}v)[1 + \mathcal{S}(u, u, \mathcal{T}u)]}{1 + \mathcal{S}(u, u, v)},$$

for all $u, v \in \mathcal{X}$, $\lambda, \eta > 0$, $\lambda + \eta < 1$. Then \mathcal{T} possess a fixed point $w \in \mathcal{X}$ which is unique.”

Lemma 1.5 [21]: “If $(\mathcal{X}, \mathcal{S})$ is an S-metric space on a non-empty set \mathcal{X} , then $(\mathcal{X}, \mathcal{S})$ satisfy the symmetric condition, that is $\mathcal{S}(u, u, v) = \mathcal{S}(v, v, u)$, for all $u, v \in \mathcal{X}$.”

Definition 1.6 [21] “Let $(\mathcal{X}, \mathcal{S})$ be an S-metric space. For $r > 0$ and $u \in \mathcal{X}$ we define the open ball $B_s(u, r)$ and closed ball and $B_s[u, r]$ with a center u and radius r as follows:

$$B_s(u, r) = \{v \in \mathcal{X} : \mathcal{S}(v, v, u) < r\}$$

$$B_s[u, r] = \{v \in \mathcal{X} : \mathcal{S}(v, v, u) \leq r\}.”$$

Definition 1.7 [22]: “A sequence $\{u_n\}$ in $(\mathcal{X}, \mathcal{S})$ is said to be convergent to some point $u \in \mathcal{X}$, if $\mathcal{S}(u_n, u_n, u) \rightarrow 0$ as $n \rightarrow \infty$.”

Definition 1.8 [22]: “A sequence $\{u_n\}$ in $(\mathcal{X}, \mathcal{S})$ is said to be Cauchy sequence if $\mathcal{S}(u_n, u_n, u_m) \rightarrow 0$ as $n, m \rightarrow \infty$.”

Definition 1.9 [22]: “An S -metric space $(\mathcal{X}, \mathcal{S})$ is said to be complete if every Cauchy sequence in X is convergent in \mathcal{X} .”

Lemma 1.10 [22]: “Let $(\mathcal{X}, \mathcal{S})$ be an S-metric space. If $u_n \rightarrow u$ and $v_n \rightarrow v$ then $\mathcal{S}(u_n, u_n, v_n) \rightarrow \mathcal{S}(u, u, v)$.”

Lemma 1.11 [23]: “Let $(\mathcal{X}, \mathcal{S})$ be an S-metric space and $\{u_n\}$ is a convergent sequence in \mathcal{X} . Then $\lim_{n \rightarrow \infty} u_n$ is unique.”

Definition 1.12 [23]: “Let $(\mathcal{X}, \mathcal{S})$ be S-metric space. A map $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ is said to be contraction if there exists a constant $k \in [0, 1)$ such that

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq k \mathcal{S}(u, u, v), \text{ for all } u, v \in \mathcal{X}.”$$

Lemma 1.13 [23]: “If $\{u_n\}$ is a sequence of elements from S-metric space $(\mathcal{X}, \mathcal{S})$ satisfying the following property $\mathcal{S}(u_n, u_n, u_{n+1}) \leq k \mathcal{S}(u_{n-1}, u_{n-1}, u_n)$, for each $k \in [0, 1)$ where $n \in \mathbb{N}$, then $\{u_n\}$ is a Cauchy sequence.”

Finally, in this article we will apply the property introduced by G.S. Jeong and B.E. Rhoades in [15] which they called the property P in metric spaces.

Definition 1.14: [15] “Let \mathcal{S} be a self-mapping of S-metric space $(\mathcal{X}, \mathcal{S})$ with a nonempty fixed point set $F(\mathcal{T})$. Then \mathcal{T} is said to satisfy the property P if $F(\mathcal{T}) = F(\mathcal{T}^n)$ for each $n \in \mathbb{N}$.”

2. Main Results

In this section, we establish fixed points of rational type contractions in the context of S-metric spaces and demonstrates that the P property is T-stable for some mappings. In order to show the relevance of the conclusions drawn in this work, we also provide a few examples.

Theorem 2.1: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping such that

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}u) + \mathcal{S}(v, v, \mathcal{T}v)\mathcal{S}(u, u, \mathcal{T}v)}{\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u)}, \quad (2.1)$$

for all $u, v \in \mathcal{X}$ and $a_1, a_2 \geq 0$, $\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u) \neq 0$ with $a_1 + a_2 < 1$. Then, \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Let u_0 be an arbitrary in \mathcal{X} , we define a sequence $\{u_n\}$ in \mathcal{X} such that $\mathcal{T}u_n = u_{n+1}$ for all $n = 1, 2, \dots$. From condition (2.1) with $u = u_n$ and $v = u_{n-1}$, Therefore

$$\begin{aligned} \mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \end{aligned}$$

$$\begin{aligned}
& + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1}) + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})} \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \\
& + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_n, u_n, u_n) + \mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1}) + \mathcal{S}(u_n, u_n, u_n)} \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_n, u_n, u_{n+1}).
\end{aligned}$$

It follows that

$$(1 - a_2)\mathcal{S}(u_n, u_n, u_{n+1}) \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.2)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1}{1-a_2}\right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \left(\frac{a_1}{1-a_2}\right)$. In view of $a_1 + a_2 < 1$, then $0 \leq \lambda < 1$. Thus, by Lemma 1.13, $\{u_n\}$ is a Cauchy sequence in \mathcal{X} such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

By (2.2), it is easy to see that

$$\begin{aligned}
\mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) & = \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*) \quad (2.3) \\
& \leq a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u^*, u^*, \mathcal{T}u_n) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u_n, u_n, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}u_n)} \\
& \leq a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u_n, u_n, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, u_{n+1})}.
\end{aligned} \quad (2.4)$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.4), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) = 0$.

That is, $u_n \rightarrow \mathcal{T}u^*$. Hence, $u^* = \mathcal{T}u^*$, u^* is a fixed point of \mathcal{T} .

Finally, we prove the uniqueness of the fixed point. Indeed, if there is another fixed point v^* , then by (2.1), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) & = \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
& \leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*) + \mathcal{S}(v^*, v^*, \mathcal{T}v^*)\mathcal{S}(u^*, u^*, \mathcal{T}v^*)}{\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}u^*)}
\end{aligned}$$

$$\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(v^*, v^*, u^*) + \mathcal{S}(v^*, v^*, v^*)\mathcal{S}(u^*, u^*, v^*)}{\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, u^*)}$$

$$\mathcal{S}(u^*, u^*, v^*) \leq a_1 \mathcal{S}(u^*, u^*, v^*). \quad (2.5)$$

Since $a_1 + a_2 < 1$ implies $a_1 < 1$.

Therefore, we obtain that $\mathcal{S}(u^*, u^*, v^*) = 0$, i.e., $u^* = v^*$.

Hence the fixed point is unique.

This completes the proof. \blacksquare

Example 2.2: Let $\mathcal{X} = [0, 1]$ be equipped with complete S-metric space define by

$$\mathcal{S}(u, v, w) = (|u - v| + |u - w| + |v - w|)^2.$$

Consider a mapping $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ defined by

$$\mathcal{T}(u) = \frac{1}{36} u^2 e^{-u^2},$$

for all $u, v, w \in \mathcal{X}$.

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) &= (|\mathcal{T}u - \mathcal{T}u| + |\mathcal{T}u - \mathcal{T}v| + |\mathcal{T}u - \mathcal{T}v|)^2 \\ &= (2 |\mathcal{T}u - \mathcal{T}v|)^2 \\ &= 4 \left| \frac{1}{36} u^2 e^{-u^2} - \frac{1}{36} v^2 e^{-v^2} \right|^2 = \left| \frac{1}{18} u^2 e^{-u^2} - \frac{1}{18} v^2 e^{-v^2} \right|^2 \\ &\leq \frac{1}{9} |u^2 e^{-u^2} - v^2 e^{-v^2}|^2 \\ &\leq \frac{4}{9} |u - v|^2 = \frac{1}{9} |2(u - v)|^2 \\ &\leq \frac{1}{3} \mathcal{S}(u, v, w) \\ &\leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}u) + \mathcal{S}(v, v, \mathcal{T}v)\mathcal{S}(u, u, \mathcal{T}v)}{\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u)}. \end{aligned}$$

Clearly by taking $a_2 = \frac{1}{2}$, we have $a_1 + a_2 = \frac{1}{3} + \frac{1}{2} = \frac{5}{6} < 1$. Then, from Theorem 2.1 we conclude that, \mathcal{T} has a unique fixed point. Also, 0 is the only fixed point of \mathcal{T} .

Theorem 2.3: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping such that

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) &\leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}v)\mathcal{S}(v, v, \mathcal{T}u)}{\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u)} \\ &\quad + a_3 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}u) + \mathcal{S}(v, v, \mathcal{T}v)\mathcal{S}(u, u, \mathcal{T}v)}{\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u)}, \end{aligned} \quad (2.6)$$

where a_1, a_2, a_3 are non-negative constant with $a_1 + a_2 + a_3 < 1$. Then, \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Choose $u_0 \in X$ and construct a Picard iterative sequence $\{u_n\}$ as $\mathcal{T}u_n = u_{n+1}$. If there exists $n_0 \in \mathbb{N}$ such that $u_{n_0} = u_{n_0+1}$, then $u_{n_0} = u_{n_0+1} = \mathcal{T}u_{n_0}$, i.e., u_{n_0} is a fixed point of \mathcal{T} . Next, without loss of generality, let $u_n \neq u_{n+1}$ for all $n \in \mathbb{N}$, Using (2.6), we get

$$\begin{aligned} \mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \\ &\quad + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})}{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})} \\ &\quad + a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1}) + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})} \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \\ &\quad + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1}) + \mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_n, u_n, u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1}) + \mathcal{S}(u_n, u_n, u_n)} \\ &\quad + a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_n, u_n, u_n) + \mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1}) + \mathcal{S}(u_n, u_n, u_n)} \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_3 \mathcal{S}(u_n, u_n, u_{n+1}). \end{aligned}$$

It follows that

$$(1 - a_3)\mathcal{S}(u_n, u_n, u_{n+1}) \leq (a_1 + a_2)\mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.7)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1 + a_2}{1 - a_3}\right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \frac{a_1 + a_2}{1 - a_3}$. In view of $a_1 + a_2 + a_3 < 1$, we have $0 \leq \lambda < 1$. Thus, from Lemma 1.13 $\{u_n\}$ is Cauchy sequence in \mathcal{X} . Since, $(\mathcal{X}, \mathcal{S})$ is a complete S-metric space, so there exists some point $u^* \in \mathcal{X}$ such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

Again from (2.6) it is easy to see that

$$\begin{aligned}
\mathcal{S}(u^*, u^*, \mathcal{T}u^*) &\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) & (2.8) \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*) \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) \\
&\quad + a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}u_n)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}u_n)} \\
&\quad + a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u^*, u^*, \mathcal{T}u_n) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u_n, u_n, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}u_n)} \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) \\
&\quad + a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, u_{n+1})}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, u_{n+1})} \\
&\quad + a_3 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u_n, u_n, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, u_{n+1})}. & (2.9)
\end{aligned}$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.9), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u^*, u^*, \mathcal{T}u^*) = 0$.

Hence, $\mathcal{T}u^* = u^*$ it follows that u^* is a fixed point of \mathcal{T} .

Next, we claim the uniqueness of fixed point.

Indeed, if there is another fixed point v^* , then by (2.6), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) &= \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}v^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*)}{\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}u^*)} \\
&\quad + a_3 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*) + \mathcal{S}(v^*, v^*, \mathcal{T}v^*)\mathcal{S}(u^*, u^*, \mathcal{T}v^*)}{\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}u^*)} \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, v^*)\mathcal{S}(v^*, v^*, u^*)}{\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, u^*)} \\
&\quad + a_3 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(v^*, v^*, u^*) + \mathcal{S}(v^*, v^*, v^*)\mathcal{S}(u^*, u^*, v^*)}{\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, u^*)} \\
\mathcal{S}(u^*, u^*, v^*) &\leq a_1 \mathcal{S}(u^*, u^*, v^*). & (2.10)
\end{aligned}$$

Since $a_1 + a_2 + a_3 < 1 \Rightarrow a_1 < 1$, we obtain that $\mathcal{S}(u^*, u^*, v^*) = 0$, i.e., $u^* = v^*$.

Hence the fixed point is unique.

This completes the proof. \blacksquare

Theorem 2.4: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space. Let $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping satisfying

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)} + a_3 \frac{\mathcal{S}(v, v, \mathcal{T}v)[1 + \mathcal{S}(u, u, \mathcal{T}u)]}{1 + \mathcal{S}(u, u, v)}, \quad (2.11)$$

for all $u, v \in \mathcal{X}$ and a_1, a_2, a_3 are non-negative constant with $a_1 + a_2 + a_3 < 1$. Then \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Choose $u_0 \in X$. Construct a sequence $\{u_n\}$ in \mathcal{X} by $\mathcal{T}u_n = u_{n+1}$. For all $n \in \mathbb{N}$, from condition (2.11) with $u = u_n$ and $v = u_{n-1}$, we have

$$\begin{aligned} \mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_n, u_n, \mathcal{T}u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\quad + a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)[1 + \mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})]}{1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_n, u_n, u_{n+1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\quad + a_3 \frac{\mathcal{S}(u_n, u_n, u_{n+1})[1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)]}{1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_n, u_n, u_{n+1}) + a_3 \mathcal{S}(u_n, u_n, u_{n+1}). \end{aligned}$$

It follows that

$$(1 - a_2 - a_3)\mathcal{S}(u_n, u_n, u_{n+1}) \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.12)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1}{1 - a_2 - a_3} \right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \frac{a_1}{1 - a_2 - a_3}$. In view of $a_1 + a_2 + a_3 < 1$, we have $0 \leq \lambda < 1$. Thus, from Lemma 1.13 $\{u_n\}$ is Cauchy sequence in \mathcal{X} . Since, $(\mathcal{X}, \mathcal{S})$ is a complete S-metric space, so there exists some point $u^* \in \mathcal{X}$ such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

Again from (2.11) it is easy to see that

$$\mathcal{S}(u^*, u^*, \mathcal{T}u^*) \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) \quad (2.13)$$

$$\begin{aligned}
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*) \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)} \\
&+ a_3 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)[1+\mathcal{S}(u_n, u_n, \mathcal{T}u_n)]}{1+\mathcal{S}(u_n, u_n, u^*)} \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)} \\
&+ a_3 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)[1+\mathcal{S}(u_n, u_n, \mathcal{T}u_{n+1})]}{1+\mathcal{S}(u_n, u_n, u^*)}. \tag{2.14}
\end{aligned}$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.14), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u^*, u^*, \mathcal{T}u^*) = 0$.

Hence, $\mathcal{T}u^* = u^*$ it follows that u^* is a fixed point of \mathcal{T} .

Next, we claim the uniqueness of fixed point.

Indeed, if there is another fixed point v^* , then by (2.11), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) &= \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_1 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(v^*, v^*, \mathcal{T}v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_1 \frac{\mathcal{S}(v^*, v^*, \mathcal{T}v^*)[1+\mathcal{S}(u^*, u^*, \mathcal{T}u^*)]}{1+\mathcal{S}(u^*, u^*, v^*)} \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(v^*, v^*, v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_3 \frac{\mathcal{S}(v^*, v^*, v^*)[1+\mathcal{S}(u^*, u^*, u^*)]}{1+\mathcal{S}(u^*, u^*, v^*)}
\end{aligned}$$

$$\mathcal{S}(u^*, u^*, v^*) \leq a_1 \mathcal{S}(u^*, u^*, v^*).$$

Since $0 < a_1 + a_2 + a_3 < 1 \Rightarrow a_1 < 1$, thus, we obtain $\mathcal{S}(u^*, u^*, v^*) = 0$, i.e., $u^* = v^*$.

Hence, we proved that \mathcal{T} have a unique fixed point in \mathcal{X} .

Here completes the proof. ■

Example 2.5: Let $\mathcal{X} = [0,1]$ and $(\mathcal{X}, \mathcal{S})$ be a usual S-metric space which is complete, define by

$$\mathcal{S}(u, v, w) = |u - w| + |v - w|.$$

Consider a mapping $\mathcal{T}: X \rightarrow X$ be define as $(u) = \frac{u}{8}$, for all $u, v, w \in X$. Obviously,

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) = 2 |\mathcal{T}u - \mathcal{T}v| = 2 \left| \frac{u}{8} - \frac{v}{8} \right| = \frac{1}{4} |u - v|,$$

$$\mathcal{S}(u, u, v) = 2 |u - v|.$$

Also,

$$\mathcal{S}(u, u, \mathcal{T}u) = 2 |u - \mathcal{T}u| = 2 \left| u - \frac{u}{8} \right| = \frac{7u}{4},$$

$$\mathcal{S}(v, v, \mathcal{T}v) = 2 |v - \mathcal{T}v| = 2 \left| v - \frac{v}{8} \right| = \frac{7v}{4},$$

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) &= \frac{1}{4} |u - v| \\ &= \frac{1}{8} 2 |u - v| \\ &\leq \frac{1}{8} \mathcal{S}(u, u, v) + \frac{1}{4} \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)} + \frac{1}{7} \frac{\mathcal{S}(v, v, \mathcal{T}v)[1 + \mathcal{S}(u, u, \mathcal{T}u)]}{1 + \mathcal{S}(u, u, v)}. \end{aligned}$$

It is clear that, $a_1 + a_2 + a_3 = \frac{1}{8} + \frac{1}{4} + \frac{1}{7} = \frac{29}{56} < 1$. Thus, we conclude that inequality (2.11) of Theorem 2.4 remains valid. Hence, \mathcal{T} has a unique fixed point and the fixed point is 0.

Theorem 2.6: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping satisfying the following condition

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) &\leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)} + a_3 \frac{\mathcal{S}(u, u, \mathcal{T}v)\mathcal{S}(v, v, \mathcal{T}u)}{\mathcal{S}(u, u, v)} \\ &\quad + a_4 [\mathcal{S}(u, u, \mathcal{T}u) + \mathcal{S}(v, v, \mathcal{T}v)] + a_5 [\mathcal{S}(v, v, \mathcal{T}u) + \mathcal{S}(u, u, \mathcal{T}v)], \end{aligned} \quad (2.15)$$

for all $u, v \in \mathcal{X}$ and a_1, a_2, a_3, a_4, a_5 are non-negative constant $a_1 + a_2 + a_3 + 2a_4 + 3a_5 < 1$. Then, \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Choose $u_0 \in \mathcal{X}$. Construct a sequence $\{u_n\}$ in \mathcal{X} by $\mathcal{T}u_n = u_{n+1}$.

For all $n \in \mathbb{N}$, from condition (2.15) with $u = u_n$ and $v = u_{n-1}$, we have

$$\begin{aligned} \mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_n, u_n, \mathcal{T}u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\quad + a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\ &\quad + a_4 [\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1}) + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)] \end{aligned}$$

$$\begin{aligned}
& +a_5[\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1}) + \mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n)] \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_n, u_n, u_{n+1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
& +a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})\mathcal{S}(u_n, u_n, u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} + a_4[\mathcal{S}(u_{n-1}, u_{n-1}, u_n) + \mathcal{S}(u_n, u_n, u_{n+1})] \\
& +a_5[\mathcal{S}(u_n, u_n, u_n) + \mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})] \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_n, u_n, u_{n+1}) \\
& +a_4[\mathcal{S}(u_{n-1}, u_{n-1}, u_n) + \mathcal{S}(u_n, u_n, u_{n+1})] \\
& +a_5[2 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + \mathcal{S}(u_n, u_n, u_{n+1})].
\end{aligned}$$

It follows that

$$(1 - a_2 - a_4 - a_5)\mathcal{S}(u_n, u_n, u_{n+1}) \leq (a_1 + a_4 + 2a_5) \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.16)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1 + a_4 + 2a_5}{1 - a_2 - a_4 - a_5} \right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \frac{a_1 + a_4 + 2a_5}{1 - a_2 - a_4 - a_5}$. In view of $a_1 + a_2 + a_3 + 2a_4 + 3a_5 < 1$, we have $0 \leq \lambda < 1$. Thus, from Lemma 1.13, $\{u_n\}$ is Cauchy sequence in \mathcal{X} . Since, $(\mathcal{X}, \mathcal{S})$ is a complete S-metric space, so there exists some point $u^* \in \mathcal{X}$ such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

Again from (2.15) it is easy to see that

$$\mathcal{S}(u^*, u^*, \mathcal{T}u^*) \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) \quad (2.17)$$

$$\begin{aligned}
& \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*) \\
& \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)} \\
& +a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}u_n)}{\mathcal{S}(u_n, u_n, u^*)} + a_4[\mathcal{S}(u_n, u_n, \mathcal{T}u_n) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)] \\
& +a_5[\mathcal{S}(u^*, u^*, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u^*)] \\
& \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)}
\end{aligned}$$

$$\begin{aligned}
& + a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, u_{n+1})}{1+\mathcal{S}(u_n, u_n, u^*)} + a_4[\mathcal{S}(u_n, u_n, u_{n+1}) + \mathcal{S}(u^*, u^*, \mathcal{T}u^*)] \\
& + a_5[\mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_n, u_n, \mathcal{T}u^*)]
\end{aligned} \tag{2.18}$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.18), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u^*, u^*, \mathcal{T}u^*) = 0$.

Hence, $\mathcal{T}u^* = u^*$ it follows that u^* is a fixed point of \mathcal{T} .

Finally, we prove the uniqueness of fixed point.

Indeed, if there is another fixed point v^* , then by (2.15), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) & = \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
& \leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(v^*, v^*, \mathcal{T}v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_3 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}v^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*)}{\mathcal{S}(u^*, u^*, v^*)} \\
& + a_4 [\mathcal{S}(u^*, u^*, \mathcal{T}u^*) + \mathcal{S}(v^*, v^*, \mathcal{T}v^*)] + a_5 [\mathcal{S}(v^*, v^*, \mathcal{T}u^*) + \mathcal{S}(u^*, u^*, \mathcal{T}v^*)] \\
& \leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(v^*, v^*, v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_3 \frac{\mathcal{S}(u^*, u^*, v^*)\mathcal{S}(v^*, v^*, u^*)}{\mathcal{S}(u^*, u^*, v^*)} \\
& + a_4 [\mathcal{S}(u^*, u^*, u^*) + \mathcal{S}(v^*, v^*, v^*)] + a_5 [\mathcal{S}(v^*, v^*, u^*) + \mathcal{S}(u^*, u^*, v^*)] \\
\mathcal{S}(u^*, u^*, v^*) & \leq (a_1 + a_3 + 2a_5) \mathcal{S}(u^*, u^*, v^*).
\end{aligned} \tag{2.19}$$

Since $0 < a_1 + a_2 + a_3 + 2a_4 + 3a_5 < 1 \Rightarrow a_1 + a_3 + 2a_5 < 1$, thus, we obtain $\mathcal{S}(u^*, u^*, v^*) = 0$, which further implies $u^* = v^*$.

Therefore, \mathcal{T} have a unique fixed point in \mathcal{X} .

Here completes the proof. \blacksquare

Theorem 2.7: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a self map that satisfies the following inequality

$$\begin{aligned}
\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) & \leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)} + a_3 \frac{\mathcal{S}(u, u, \mathcal{T}v)\mathcal{S}(v, v, \mathcal{T}u)}{\mathcal{S}(u, u, v)} \\
& + a_4 \frac{\mathcal{S}(v, v, \mathcal{T}v)[1+\mathcal{S}(u, u, \mathcal{T}u)]}{1+\mathcal{S}(u, u, v)},
\end{aligned} \tag{2.20}$$

for all $u, v \in \mathcal{X}$ and a_1, a_2, a_3, a_4 are non-negative constant $a_1 + a_2 + a_3 + a_4 < 1$. Then, \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Choose $u_0 \in X$ and construct a Picard iterative sequence $\{u_n\}$ as $\mathcal{T}u_n = u_{n+1}$.

For all $n \in \mathbb{N}$, from condition (2.20) with $u = u_n$ and $v = u_{n-1}$, we have

$$\begin{aligned}
\mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\
&\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_n, u_n, \mathcal{T}u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
&\quad + a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
&\quad + a_4 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)[1 + \mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})]}{1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
&\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_n, u_n, u_{n+1})}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
&\quad + a_3 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})\mathcal{S}(u_n, u_n, u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)} + a_4 \frac{\mathcal{S}(u_n, u_n, u_{n+1})[1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)]}{1 + \mathcal{S}(u_{n-1}, u_{n-1}, u_n)} \\
&\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_n, u_n, u_{n+1}) + a_4 \mathcal{S}(u_n, u_n, u_{n+1}),
\end{aligned}$$

which further implies,

$$(1 - a_2 - a_4)\mathcal{S}(u_n, u_n, u_{n+1}) \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.21)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1}{1 - a_2 - a_4}\right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \frac{a_1}{1 - a_2 - a_4}$. In view of $a_1 + a_2 + a_3 + a_4 < 1$, we have $0 \leq \lambda < 1$. Thus, from Lemma

1.13, $\{u_n\}$ is Cauchy sequence in \mathcal{X} . Since, $(\mathcal{X}, \mathcal{S})$ is a complete S-metric space, so there exists some point $u^* \in \mathcal{X}$ such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

Again from (2.20) it is easy to see that

$$\begin{aligned}
\mathcal{S}(u^*, u^*, \mathcal{T}u^*) &\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) \quad (2.22) \\
&= 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*) \\
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)} \\
&\quad + a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}u_n)}{\mathcal{S}(u_n, u_n, u^*)} + a_4 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)[1 + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)]}{1 + \mathcal{S}(u_n, u_n, u^*)}
\end{aligned}$$

$$\begin{aligned}
&\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*) + a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u^*, u^*, \mathcal{T}u^*)}{\mathcal{S}(u_n, u_n, u^*)} \\
&+ a_3 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, u_{n+1})}{1+\mathcal{S}(u_n, u_n, u^*)} + a_4 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)[1+\mathcal{S}(u_n, u_n, u_{n+1})]}{1+\mathcal{S}(u_n, u_n, u^*)}.
\end{aligned} \tag{2.23}$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.23), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u^*, u^*, \mathcal{T}u^*) = 0$.

Hence, $\mathcal{T}u^* = u^*$ it follows that u^* is a fixed point of \mathcal{T} .

Finally, we prove the uniqueness of fixed point.

Indeed, if there is another fixed point v^* , then by (2.20), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) &= \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_1 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(v^*, v^*, \mathcal{T}v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_1 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}v^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*)}{\mathcal{S}(u^*, u^*, v^*)} \\
&+ a_4 \frac{\mathcal{S}(v^*, v^*, \mathcal{T}v^*)[1+\mathcal{S}(u^*, u^*, \mathcal{T}u^*)]}{1+\mathcal{S}(u^*, u^*, v^*)} \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(v^*, v^*, v^*)}{\mathcal{S}(u^*, u^*, v^*)} + a_3 \frac{\mathcal{S}(u^*, u^*, v^*)\mathcal{S}(v^*, v^*, u^*)}{\mathcal{S}(u^*, u^*, v^*)} \\
&+ a_4 \frac{\mathcal{S}(v^*, v^*, v^*)[1+\mathcal{S}(u^*, u^*, u^*)]}{1+\mathcal{S}(u^*, u^*, v^*)}.
\end{aligned}$$

$$\mathcal{S}(u^*, u^*, v^*) \leq (a_1 + a_3) \mathcal{S}(u^*, u^*, v^*)$$

$$\mathcal{S}(u^*, u^*, v^*) \leq (a_1 + a_2 + a_3 + a_4) \mathcal{S}(u^*, u^*, v^*), \tag{2.24}$$

a contradiction.

Thus, we obtain $\mathcal{S}(u^*, u^*, v^*) = 0$, which further implies $u^* = v^*$.

Therefore, \mathcal{T} have a unique fixed point in \mathcal{X} .

Here completes the proof. \blacksquare

Example 2.8: Let $\mathcal{X} = [0,1]$ be equipped with complete S-metric space define by

$$\mathcal{S}(u, v, w) = (|u - v| + |u - w| + |v - w|)^2.$$

Let the mapping $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be defined by

$$\mathcal{T}(u) = \frac{u}{5}.$$

Then, for all $u, v, w \in \mathcal{X}$, we have,

$$\mathcal{S}(u, u, v) = 4|u - v|^2,$$

$$\mathcal{S}(u, u, \mathcal{T}u) = 4|u - \mathcal{T}u|^2 = 4\left|u - \frac{u}{5}\right|^2 = \frac{16}{25}u^2$$

$$\mathcal{S}(v, v, \mathcal{T}v) = 4|v - \mathcal{T}v|^2 = 4\left|v - \frac{v}{5}\right|^2 = \frac{16}{25}v^2$$

Also,

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) &= (|\mathcal{T}u - \mathcal{T}u| + |\mathcal{T}u - \mathcal{T}v| + |\mathcal{T}u - \mathcal{T}v|)^2 \\ &= (2|\mathcal{T}u - \mathcal{T}v|)^2 \\ &= \frac{4}{25}|u - v|^2 \\ &\leq \frac{1}{25}\mathcal{S}(u, v, w) + \frac{4}{25}\frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(v, v, \mathcal{T}v)}{\mathcal{S}(u, u, v)} + \frac{2}{5}\frac{\mathcal{S}(v, v, \mathcal{T}v)[1 + \mathcal{S}(u, u, \mathcal{T}u)]}{1 + \mathcal{S}(u, u, v)} \end{aligned}$$

Clearly, we have $a_1 + a_2 + a_3 = \frac{1}{25} + \frac{4}{25} + \frac{2}{5} = \frac{3}{5} < 1$. Then, from Theorem 2.7 we conclude that, \mathcal{T} has a unique fixed point. Also, 0 is the only fixed point of \mathcal{T} .

Theorem 2.9: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space. Let $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping satisfying

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v) \leq a_1 \mathcal{S}(u, u, v) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}v)\mathcal{S}(v, v, \mathcal{T}u)}{\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u)} + a_3 \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}v), \quad (2.25)$$

for all $u, v \in \mathcal{X}$ and $a_1, a_2, a_3 \geq 0$. $\mathcal{S}(u, u, \mathcal{T}v) + \mathcal{S}(v, v, \mathcal{T}u) \neq 0$ with $a_1 + a_2 + a_3 < 1$. Then \mathcal{T} has a unique fixed point \mathcal{X} .

Proof: Choose u_0 as an arbitrary point in X . We define a sequence $\{u_n\}$ in \mathcal{X} by $\mathcal{T}u_n = u_{n+1}$. Then or all $n \in \mathbb{N}$, from condition (2.25) with $u = u_n$ and $v = u_{n-1}$, we have

$$\begin{aligned} \mathcal{S}(u_n, u_n, u_{n+1}) &= \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\ &\leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \\ &\quad + a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_{n-1})\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})}{\mathcal{S}(u_{n-1}, u_{n-1}, \mathcal{T}u_n) + \mathcal{S}(u_n, u_n, \mathcal{T}u_{n-1})} \end{aligned}$$

$$\begin{aligned}
& +a_3 \mathcal{S}(\mathcal{T}u_{n-1}, \mathcal{T}u_{n-1}, \mathcal{T}u_n) \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) \\
& +a_2 \frac{\mathcal{S}(u_{n-1}, u_{n-1}, u_n)\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})+\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_n, u_n, u_n)}{\mathcal{S}(u_{n-1}, u_{n-1}, u_{n+1})+\mathcal{S}(u_n, u_n, u_n)} \\
& +a_3 \mathcal{S}(u_n, u_n, u_{n+1}) \\
& \leq a_1 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_2 \mathcal{S}(u_{n-1}, u_{n-1}, u_n) + a_3 \mathcal{S}(u_n, u_n, u_{n+1}).
\end{aligned}$$

Which further implies

$$(1 - a_3)\mathcal{S}(u_n, u_n, u_{n+1}) \leq (a_1 + a_2)\mathcal{S}(u_{n-1}, u_{n-1}, u_n) \quad (2.26)$$

$$\mathcal{S}(u_n, u_n, u_{n+1}) \leq \left(\frac{a_1 + a_2}{1 - a_3}\right) \mathcal{S}(u_{n-1}, u_{n-1}, u_n).$$

Put $\lambda = \frac{a_1 + a_2}{1 - a_3}$. In view of $a_1 + a_2 + a_3 < 1$, we have $0 \leq \lambda < 1$. Thus, from Lemma 1.13 $\{u_n\}$ is Cauchy sequence in \mathcal{X} . Since, $(\mathcal{X}, \mathcal{S})$ is a complete S-metric space, so there exists some point $u^* \in \mathcal{X}$ such that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.

Again from (2.25) it is easy to see that

$$\mathcal{S}(u^*, u^*, \mathcal{T}u^*) \leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) \quad (2.27)$$

$$\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*)$$

$$\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*)$$

$$+a_2 \frac{\mathcal{S}(u_n, u_n, \mathcal{T}u_n)\mathcal{S}(u_n, u_n, \mathcal{T}u^*)+\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}u_n)}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)+\mathcal{S}(u^*, u^*, \mathcal{T}u_n)}$$

$$+a_3 \mathcal{S}(\mathcal{T}u_n, \mathcal{T}u_n, \mathcal{T}u^*)$$

$$\leq 2 \mathcal{S}(u^*, u^*, u_{n+1}) + a_1 \mathcal{S}(u_n, u_n, u^*)$$

$$+a_2 \frac{\mathcal{S}(u_n, u_n, u_{n+1})\mathcal{S}(u_n, u_n, \mathcal{T}u^*)+\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, u_{n+1})}{\mathcal{S}(u_n, u_n, \mathcal{T}u^*)+\mathcal{S}(u^*, u^*, u_{n+1})}$$

$$+ a_3 \mathcal{S}(u_{n+1}, u_{n+1}, \mathcal{T}u^*) . \quad (2.28)$$

Taking the limit as $n \rightarrow \infty$ on both side of (2.28), we have $\lim_{n \rightarrow \infty} \mathcal{S}(u^*, u^*, \mathcal{T}u^*) = 0$.

Hence, $\mathcal{T}u^* = u^*$ it follows that u^* is a fixed point of \mathcal{T} .

Finally, we claim the uniqueness of fixed point.

Indeed, if there is another fixed point v^* , then by (2.25), we have

$$\begin{aligned}
\mathcal{S}(u^*, u^*, v^*) &= \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, \mathcal{T}u^*)\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}v^*)\mathcal{S}(v^*, v^*, \mathcal{T}u^*)}{\mathcal{S}(u^*, u^*, \mathcal{T}v^*) + \mathcal{S}(v^*, v^*, \mathcal{T}u^*)} \\
&\quad + a_3 \mathcal{S}(\mathcal{T}u^*, \mathcal{T}u^*, \mathcal{T}v^*) \\
&\leq a_1 \mathcal{S}(u^*, u^*, v^*) + a_2 \frac{\mathcal{S}(u^*, u^*, u^*)\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, v^*)\mathcal{S}(v^*, v^*, u^*)}{\mathcal{S}(u^*, u^*, v^*) + \mathcal{S}(v^*, v^*, u^*)} \\
&\quad + a_3 \mathcal{S}(u^*, u^*, v^*) \\
\mathcal{S}(u^*, u^*, v^*) &\leq (a_1 + a_3) \mathcal{S}(u^*, u^*, v^*). \tag{2.29}
\end{aligned}$$

Since $0 < a_1 + a_2 + a_3 < 1 \Rightarrow a_1 + a_3 < 1$, thus, we obtain $\mathcal{S}(u^*, u^*, v^*) = 0$, i.e., $u^* = v^*$.

Hence, we proved that \mathcal{T} have a unique fixed point in \mathcal{X} .

Here completes the proof. \blacksquare

Remark 2.10.

1. If we put $a_2 = a_3 = a_4 = a_5 = 0$ in Theorem 2.6, we get the Banach Contraction Theorem [2] in S-metric space.
2. If we put $a_1 = a_2 = a_3 = a_5 = 0$ in Theorem 2.6, we get the Kanan Theorem [11] in S-metric space.
3. If we put $a_2 = a_3 = a_5 = 0$ in Theorem 2.6, we get the Fisher Theorem [9] in S-metric space.
4. If we put $a_1 = a_2 = a_3 = a_4 = 0$ in Theorem 2.6, we get the result of Chaterjee Theorem [5] in S-metric space.
5. If we put $a_2 = a_3 = 0$ in Theorem 2.9, we get the result of Dass and Gupta Theorem [8] in S-metric space.

Theorem 2.11: Let $(\mathcal{X}, \mathcal{S})$ be a complete S-metric space and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ be a mapping such that $F(\mathcal{T}) \neq \Phi$ and that

$$\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}^2u) \leq \lambda \mathcal{S}(u, u, \mathcal{T}u), \quad (2.30)$$

for all $u \in \mathcal{X}$, where $0 \leq \lambda < 1$ is a constant. Then \mathcal{T} has the P property.

Proof: We always assume that $n > 1$. Since the statement for $n = 1$ is trivial. Let $w \in F(\mathcal{T}^n)$.

By the hypotheses, we get

$$\begin{aligned} \mathcal{S}(w, w, \mathcal{T}w) &= \mathcal{S}(\mathcal{T}\mathcal{T}^{n-1}w, \mathcal{T}\mathcal{T}^{n-1}w, \mathcal{T}^2\mathcal{T}^{n-1}w) \\ &\leq \lambda \mathcal{S}(\mathcal{T}^{n-1}w, \mathcal{T}^{n-1}w, \mathcal{T}^n w) \\ &\leq \lambda \mathcal{S}(\mathcal{T}\mathcal{T}^{n-2}w, \mathcal{T}\mathcal{T}^{n-2}w, \mathcal{T}^2\mathcal{T}^{n-2}w) \\ &\leq \lambda^2 \mathcal{S}(\mathcal{T}^{n-2}w, \mathcal{T}^{n-2}w, \mathcal{T}^{n-1}w) \\ &\leq \dots \leq \lambda^n \mathcal{S}(w, w, \mathcal{T}w) \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

Hence $\mathcal{S}(w, w, \mathcal{T}w) = 0$, that is $\mathcal{T}w = w$.

Theorem 2.12: Under the condition of Theorem 2.3, \mathcal{T} has the P property.

Proof: We have to prove that the mapping \mathcal{T} satisfies (2.30). In fact, for any $u \in \mathcal{X}$, we have

$$\begin{aligned} \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}^2u) &= \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}\mathcal{T}u) \\ &\leq a_1 \mathcal{S}(u, u, \mathcal{T}u) + a_2 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(u, u, \mathcal{T}\mathcal{T}u) + \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}\mathcal{T}u)\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}u)}{\mathcal{S}(u, u, \mathcal{T}\mathcal{T}u) + \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}u)} \\ &\quad + a_3 \frac{\mathcal{S}(u, u, \mathcal{T}u)\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}u) + \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}\mathcal{T}u)\mathcal{S}(u, u, \mathcal{T}\mathcal{T}u)}{\mathcal{S}(u, u, \mathcal{T}\mathcal{T}u) + \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}u)} \\ &\leq a_1 \mathcal{S}(u, u, \mathcal{T}u) + a_2 \mathcal{S}(u, u, \mathcal{T}u) + a_3 \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}\mathcal{T}u) \\ (1 - a_3)\mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}^2u) &\leq (a_1 + a_2)\mathcal{S}(u, u, \mathcal{T}u) \\ \mathcal{S}(\mathcal{T}u, \mathcal{T}u, \mathcal{T}^2u) &\leq \frac{a_1 + a_2}{1 - a_3} \mathcal{S}(u, u, \mathcal{T}u) \end{aligned}$$

Deduce that $\lambda = \frac{a_1 + a_2}{1 - a_3}$. Note that $a_1 + a_2 + a_3 < 1$, then $\lambda < 1$. Accordingly, (2.30) is satisfied.

Consequently, by Theorem 2.3, \mathcal{T} has the P property.

Conclusion: In this article, we have established the existence of fixed points theorems of rational type contractions mappings in the framework of S-metric spaces and also studied the

P property for some mappings. To further demonstrated the reliability of the findings in the article, we additionally offer a few examples.

References

- [1] Amini-Harandi A., Fixed point theorem for quasi-contraction maps in b-metric spaces, Fixed Point Theory. Vol. 5(2), pp 351-358, (2014).
- [2] Asadi M., Karapınar E., Kumar A., $\alpha - \psi$ -Geraghty contractions on generalized metric spaces, Journal of Inequalities and Applications Vol. (1), pp 423, (2014).
- [3] Asadi M., Discontinuity of control function in the (F, φ, θ) -Contraction in metric spaces Filomat Vol. 31(17), pp 5427-5433.
- [4] Asadi M., Salimi P., Some fixed point and common fixed point theorems on G-metric spaces, Nonlinear Functional Analysis and Applications. Vol. 21(3), pp 523-530.
- [5] Asadi M, Khalesi A., Lower semi-continuity in a generalized metric space, Advances in the Theory of Nonlinear Analysis and its Applications(ATNAA) 1 ...
- [6] Banach S., Sur les opérations dans les ensembles abstraits et leur application aux equations integrals, Fund. Math. Vol. 3, pp 133-181, (1922).
- [7] Bakhtin I. A., The contraction mapping principle in almost metric spaces. Funct. Anal. Vol. 30, pp 26-37, (1989).
- [8] Chatterjee S. Fixed point theorems, Dokladi na Bolgarskata Akademiya na Naukite. Vol. 25(6), pp 727-730, (1989).
- [9] Czerwik S., The contraction mapping in b - metric spaces. Acta Math. Inform. Univ. Ostrav. Vol. 1, pp 5- 11, (1993).
- [10] Czerwik, S. Nonlinear set - valued contraction mapping in b -metric spaces. Atti Semin. Mat. Fis. Dell'Universita Modena Reeggio Emilia. Vol. 46, pp 263–276, (1998).

- [11] Das B. K. and Gupta S., An extension of Banach contraction principle through rational expression, Indian J. Pure Appl. Math. Vol. 6(12), pp 1455-1458, (1975).
- [12] Eshraghisamani M., Vaezpour M., Asadi M., New fixed point results on Branciari metric spaces, Journal of Mathematical Analysis Vol. 8(6), pp 132-141.
- [13] Fisher B., A fixed point theorem for compact metric spaces, Publ. Math. Debrecen. Vol. 25 (1978).
- [14] Hussein M. M., Bin Saad M.G., Al-Sayad A. A., Fixed point results of relation type- Contraction mapping in b-metric spaces with an application, Earthline Journal of Mathematical science. Vol. 12(2), pp 141-164, (2023).
- [15] Jeong G. S. and Rhoades B. E., Maps for which $F(T) = F(T^n)$, Fixed Point Theory and Appl., Vol. 6, pp 87–131, (2005).
- [16] Jeong G. S. and Rhoades B. E., More maps for which $F(T) = F(T^n)$, Demonstratio Math., Vol. 40, pp 671–680, (2007).
- [17] Kannan R., Some results on fixed points, Bull. Cal. Math. Soc. Vol. 60, pp 71-76, (1968).
- [18] Manoj K., Sushma D., Parul S., Fixed point theorems by using altering distance function in S-metric space, Communication in Mathematics and application. Vol. 13(2), pp 553-573, (2022).
- [19] Monfared H., Asadi M., Azhini M., $F(\psi, \theta)$ -contractions for α -admissible mappings on metric spaces and related fixed point results, Communications in Nonlinear Analysis (CNA). Vol. 2(1), pp 86-94.
- [20] Naddler S. B. Jr, Multi-valued contraction mappings. Pac. J. Math. Vol. 30, pp 475-488, (1969).
- [21] Sedghi S., Shobe N., Aliouche A., A generalization of fixed point theorem in S-metric spaces. Mat. Vesnik. Vol. 64, pp 258-266, (2012).
- [22] Sedghi S. and Dung N. V., Fixed point theorems on S-metric space, Mat. Vesnik, Vol.

66(1), pp 113–124, (2014).

[23] Sedghi S., Shobe N., Dosenovic T., Fixed Point Results In S-Metric Spaces, *Nonlinear Funct. Anal. Appl.*, Vol. 20(1), pp 55–67, (2015).

[24] Sedghi S., Kyu Kim J., Gholidahneh A., Mahdi Rezaee M., Fixed point theorems in S-metric Spaces, *East Asian Math. J.* Vol. 32(5), pp 677–684, (2016).

[25] Sedghi S., Gholidahneh A., Dosenovic T., Esfahani J., Radenovic S., Common fixed point of four maps in S-metric spaces, *J. Linear Topol. Algebra.* Vol. 5(2), pp 93–104, (2016).

[26] Younis M., Singh D., Asadi M., Josh, V., Results on contractions of Reich type in Graphical - metric spaces with applications, *Filomat* Vol. 33(17), pp 5723–5735.