

Standardization of softwood grafting in Avocado (*Persea americana* Mill) var. TKD1

ABSTRACT

In recent years, Hass avocados (*Persea americana* Mill) have become more widely cultivated in Colombia. One of the difficulties facing the nation is boosting orchard output by ensuring graft plants have the best agronomic, genetic, and phytosanitary criteria. In this study, the growth and development of avocado rootstocks derived from seeds of various weight ranges will be monitored for 90 days. Four weights of avocado seeds (S₁, S₂, S₃, S₄) (30–40 g, 40–50 g, 50–60 g, and 60–70 g) and four scion maturity of (D₁, D₂, D₃, D₄) (60 days old, 70 days old, 80 days old, 90 days old) were used as treatments in a factorial completely randomized experimental design. The outcome showed that the best germination rate of 75% for S₄ and the highest survival percentage (94.98%) for S₄D₄. After 90th DAG the maximum number of new shoots per plant (5.75), number of leaves/plants (18.15), leaf area (20.83 cm²), and carbohydrate content (6.57%) were reported. Based on the results of the experiment, it was determined that the optimum method for propagating avocados was softwood grafting using 60–70 g seed size (rootstock) and 90-day-old scion. These plants had better vigor and faster germination rates, all of which could reduce the length of time spent in the nursery while still ensuring the quality of the material, which is influenced by improved seedling growth. Thus, there is a need to standardize the time and method of propagation in avocado under the tropical humid conditions of India so that the vegetative propagated plants may be provided to the growers to get better yield and quality in future.

Keywords: seed weight, Germination, scion maturity, carbohydrate, Avocados

Introduction

The Avocado (*Persea americana* Mill) is native to tropical America, mostly Mexico and Central America. One of the challenges facing the country is ensuring graft plants adhere to the highest agronomic, genetic, and phytosanitary criteria in order to increase orchard productivity. It is one of the healthiest fruits, with greater protein levels (up to 4%), higher fat levels (up to 30%), and lower sugar levels (1.0%). Avocado is often consumed raw, as a filling for sandwiches, or in salads. Additionally, the pulp may be frozen and used in milkshakes and ice cream. Although avocados may grow in a range of soils and climates, they are sensitive to poor drainage and are unable to withstand waterlogging. Sri Lankan farmers brought avocados to India in the early 20th century.

It is farmed on a modest scale in several sections of Tamil Nadu, Kerala, Karnataka, and the North Eastern Himalayan states. Many other states, however, have gardens where plants may be observed flourishing. Avocado cultivation is becoming increasingly popular owing to its nutritional benefits and increased health consciousness. Avocado is cultivated as a mixed crop in the coffee-based cropping system in Kodagu (Karnataka), The Palani, Shervaroy Hills (Tamil Nadu), and other regions. In some regions of Tamil Nadu, Kerala, Karnataka, and the North Eastern Himalayan states, it is grown on a small basis. However, many other states in the nation have gardens where plants can be seen growing. Avocado farming is currently becoming more and more popular due to its nutritional benefits and rising health consciousness. Avocado is planted as one of the mixed crops in the coffee-based cropping system in Kodagu (Karnataka), The Palani, Shervaroy Hills (Tamil Nadu), and other locations.

The production of seeds in various regions without fulfilling the minimal quality requirements has led to a large amount of planting material that, when used in the field, starts to pose issues, mostly at the root level. The cultures with a wide range of responses to edaphoclimatic and sanitary circumstances are a natural consequence of significant genetic variety (Ben-Ya'acov and Michelson, 1995). As a result, the rootstock's seed must come from mature, fruitful trees that have produced at least two crops, are well-formed, and have excellent, healthy fruits Bernal et al. (2014).

(There is a significant volume of planting material that, when utilized in the field, starts to pose problems, primarily at the root level, as a result of the manufacturing of seeds in various locations that do not meet the basic quality criteria. The cultures' various reactions to edaphoclimatic and sanitary conditions are a natural result of the high genetic diversity (Ben-Ya'acov and Michelson, 1995). Consequently, the seed for the rootstock must originate from old, fertile trees that have produced at least two crops, are well-formed, and have great, healthy fruits. Bernal and others (2014).)

In terms of the growth and development of avocado fruit, the rootstock gives the tree nutrients, water, and other molecules like hormones and proteins through the root system that are important for healthy fruit development.

Lazare et al. (2020) state that while the scion produces the required photo assimilates through photosynthesis, an appropriate rootstock-scion connection is still required to ensure high yields and good fruit quality.

Yields and fruit quality are affected by abiotic elements (factors) including light intensity, water stress, temperature, and soil fertility, biotic ones like pests and diseases, and genetic factors like the rootstock (Mickelbart, 2012). An excellent root system that can withstand unfavorable conditions and ensures the early growth of the fruit can favor these elements, limiting the fruit's permanence over the tree (Lira et al., 2020). Therefore, seed weight and scion maturity for softwood grafting must be standardized as a result.

Materials and Methods

The experiment on "Standardization of softwood grafting in avocado (*Persea americana* Mill) var. TKD1" was carried out in a mist chamber from February to July 2023 at the Horticultural Research station Thadiyankudisai during the year 2022-2023. The latitude and longitude of the location are 10°S and 77°E with an elevation of 1098 m above Mean Sea Level. The average rainfall per year is about 1400 mm. The experiment had 16 treatments and 3 replications, and it was set up in a factorial completely randomized design (FCRD). In this softwood grafting is followed and scion materials are collected from a 20-year-old avocado tree of variety TKD 1 at the Horticultural Research Station Thadiyankudisai, scion materials for softwood grafting were gathered.

Factor 1: Size of Seed (Rootstock)

S₁ – 30-40g

S₂ – 40-50g

S₃ – 50-60g

S₄ – 60-70g

Factor 2: Age of Scion (Softwood)

D₁ – 60 days old

D₂ – 70 days old

D₃ – 80 days old

D₄ – 90 days old

List 1. List of Treatment combinations used for the study

S*D	D ₁	D ₂	D ₃	D ₄
S ₁	S ₁ D ₁	S ₁ D ₂	S ₁ D ₃	S ₁ D ₄
S ₂	S ₂ D ₁	S ₂ D ₂	S ₂ D ₃	S ₂ D ₄
S ₃	S ₃ D ₁	S ₃ D ₂	S ₃ D ₃	S ₃ D ₄
S ₄	S ₄ D ₁	S ₄ D ₂	S ₄ D ₃	S ₄ D ₄

The biometrical observations were made on five randomly chosen plants from each replication to evaluate the morphological traits, for rootstock such as the number of days taken for sprouting, plant height, leaf area, shoot length, and observations after grafting with scion number of days taken to bud sprouting, number of new shoots/plants, survival percentage of grafting, leaf area (cm²), carbohydrate content. The data for the number of days taken for sprouting for rootstock was recorded after the emergence of 1st sprouting and expressed in numbers. The plant height of rootstock plants was measured on the 30th, 45th, and 60th day after sprouting and expressed in cm. The leaf area was calculated at 60 days after sprouting.

The after-grafting data regarding the number of days taken to sprouting was calculated by observation of plants on alternate days from the days of planting and their mean value was used to calculate the days taken for the first sprout. The number of shoots per plant and number of new leaves per plant were recorded of fully developed plants were recorded 30, 60, 90 days after propagation. The data on the percentage of survival were recorded after 90 days of planting the grafting/budding/cutting, carbohydrate content, and leaf area calculated after 90 days of propagation. The data were then statistically evaluated using the Panse and Sukhtame (1954) approach.

Result and Discussion

1. Germination percentage (%)

(Figure 1) Avocado seed germination rates varied according to seed size, with S₄ (60–70 g) showing the highest rate and S₃ (50–60 g), S₂ (40–50 g), and S₁ (30–40 g) showing the lowest. Seed size and germination percentage were shown to be directly correlated, with 85% for S₄ (60–70 g), 75% for S₃ (50–60 g), 69% for S₂ (40–50 g), and 60% for S₁ (30–40 g). The size and quantity of seed reserves, which should be related to a stronger nutritional reserve in bigger seeds, are related to the success of seed germination and the establishment of seedlings. During the formation of the avocado fruit, significant starch reserves build up in the seed, generating a pool of carbohydrates reserves that are gradually broken down to sucrose, which is then used for the initial growth of the embryo and seedling Tefayetal. (2012). This outcome is related to the avocado studies by Mejia et al. (2022), Ndoro (2018), and Adjei et al. (2011).

2. Number of days taken for sprouting (days)

According to a study of the data in (table 1), different seed sizes and scion maturities, as well as the minimum (19.25) number of days required for sprouting in S₄, were documented. The minimal number of days required for avocados to sprout (20.75) was seen in D₄, indicating that the scion's maturity had a major impact on this factor. The interplay between avocado seed weight and scion age also affected the length of time it took for avocados to sprout, with S₄ (60–70 g) and D₄ (90 days old) requiring the least amount of time 17 days to sprout. In softwood grafting the size of the seed and scion maturity plays a major role in seedling growth, whereas larger seed size and 90 days old scion should be used for grafting. During avocado fruit development, substantial starch reserves build up in the seed, generating a pool of carbohydrate reserves that are gradually broken down to sucrose, which is needed for initial embryo and seedling growth. The results conformed with those of Cordula et al. (2014), Tefay et al. (2012), and Ndoro (2018) in avocado.

3. Number of shoots per plant

(Table 2) The results showed that the highest (5.37) and minimum (2.55) shoots per plant were recorded in S₄ and S₁, depending on the seed size and scion maturity. The maximum (4.46) and minimum (3.78) shoots were seen during D₄ and D₁, indicating that the scion maturity had a substantial impact on the time taken for avocado seeds to sprout. The interaction between avocado seed size and scion maturity also had an impact on the number of shoots, with the greatest (5.75) and minimum (1.95) values seen in S₄ D₄ and S₁ D₁, respectively. The average number of stems on the seedlings with the largest seeds (above 70 g) was three, which is the most stems ever seen Paixo et al. (2016). This is probably because larger seeds contain more nutrients, which encourages the development of multi-stem rods. The number of shoots produced is more influenced by the greater seed size and the scion's age in days. Similar results were obtained in avocados by Souza et al. (2020) and Paixo et al. (2016).

4. Number of leaves per plant

(Table 3) The effect of seed size and scion maturity on the number of leaves per plant showed that the highest value (17.59) and lowest value (12.40) of leaves were recorded in S₄ and S₁. The effect of scion maturity was also significant with respect to the time it took for avocados to sprout, with the highest (15.62) and lowest (14.43) leaves being observed during D₄ and D₁. The interaction effect of seed size and scion maturity in avocados also had an impact on the number of leaves, with the highest (18.15) and lowest (11.98) values found in S₄D₄ and S₁D₁. The rootstock and scion have an impact on how quickly grafts grow by increasing the production of photosynthates and dry matter, improving soil nutrient and water intake, and increasing the amount of synthetic food material. The scion's ideal maturity aids in the production of additional leaves. These findings concur with those in avocados from Mejia et al. (2021) and Olorunmaiye et al. (2011). Nimbalkar (2011) on karonda attained similar outcomes.

5. Survival percentage (%)

(Table 4) According to the study's findings, S₄ and S₁ had the highest and lowest survival rates, respectively (93.98% and 60.83%). Regarding the avocado's taken-to-survival percentage, the effect of scion maturity was also significant; the highest (84.53%) and lowest (74.27%) percentages were seen on the D₄ and D₁, respectively. The interaction between seed size and scion maturity in avocados also had an impact on survival percentage, with S₄D₄ and S₁D₁ showing the highest and lowest survival percentages (94.98 and 52.34%, respectively). The sufficient amount of carbohydrates and other food components in the scion and rootstock, as well as the mobilization of the stored food components for new development, maybe the cause of the survival percentage. This in turn leads to high meristematic activity in the scion. Tanuja and Thippesha (2017) in the sapota study, Maheswari and Nivetha (2015) in the jackfruit study, and Angadi et al. (2011) in the wood apple study all provided support for this conclusion.

6. Leaf area (cm²)

Analysis of the results (Table 5) showed that S₄ and S₁ had the highest and lowest leaf areas among seeds of various sizes and scion maturities, respectively, measuring 19.99 cm² and 16.03 cm². Regarding avocado leaf area, the maximum (18.81) and minimum (17.29) values were noted in D₄ and D₁ as the effect of scion maturity. The

interaction impact of avocado seed size and scion maturity, which was found to have maximum (19.20) and minimum (15.16) values in S₄D₄ and S₁D₁, also affected the time it took for avocado buds to sprout. The early and vigorous union formation, better nutrient uptake, sufficient scion development, and suitable rootstock may have contributed to the increased number of leaves per plant and leaf area. The same outcomes were observed in studies on sapota by Tanuja and Thippesha (2017), mango by Alam et al. (2006), and guava by Rajatiya et al. (2022).

7. Carbohydrate content (%)

(Table 6) Carbohydrate levels were found to vary with seed size and scion age, with S₄ and S₃ having the highest (6.13%) and lowest (5.72%) levels, respectively. About the carbohydrate content of avocados, the effect of scion maturity was likewise significant; the highest (6.08%) and lowest (5.60%) values were seen on days D₄ and D₁. The interplay between avocado seed size and scion maturity also had an impact on the fruit's carbohydrate content, with maximum (6.57%) and minimum (5.28%) levels found in S₄D₄ and S₁D₁. The rootstock's scion maturity and seed size have a significant impact on the development of greater carbohydrate contents. The right seed size and level of maturity leads to an increase in carbohydrate content, which encourages high graft intake and raises the success rate. For mango, this conclusion was in line with Karna et al. (2017).

The buildup of higher carbohydrate contents is highly dependent on the scion maturity and seed size of the rootstock. Increased carbohydrate content, which promotes high graft intake and increases the success rate, is a result of the proper seed size and maturation level. This finding agreed with Karna et al. (2017) for mango.




Figure: 1 Effect of seed size on germination percentage

Table 1: Influence of the size of the seed, scion maturity, and their interaction effect on the number of days taken to bud sprouting of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	35.25	32.72	26.24	24.68	29.72
S ₂	28.61	26.82	24.55	22.74	25.68
S ₃	27.93	25.72	22.85	20.13	24.15
S ₄	22.69	20.74	19.58	17.62	20.15
Mean	28.62	26.50	23.30	21.29	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.208		0.208		0.417
CD(p=0.05)	0.425		0.425		0.850

Table 2: Influence of the size of the seed, scion maturity, and their interaction effect on the number of shoots per plant of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	1.95	2.20	2.90	3.10	2.55
S ₂	3.64	3.75	3.95	4.10	3.86
S ₃	4.50	4.79	4.89	4.90	4.77
S ₄	4.99	5.20	5.55	5.75	5.37
Mean	3.78	3.98	4.32	4.46	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.050		0.050		0.100
CD(p=0.05)	0.137		0.137		0.274

Table 3: Influence of the size of the seed, scion maturity, and their interaction effect on the number of leaves per plant of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	11.98	12.14	12.56	12.95	12.40
S ₂	13.45	13.98	14.32	14.89	14.16
S ₃	15.32	15.85	16.21	16.52	15.97
S ₄	16.98	17.34	17.89	18.15	17.59
Mean	14.43	14.82	15.24	15.62	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.169		0.169		0.338
CD(p=0.05)	0.344		0.344		0.689

Table 4: Influence of the size of the seed, scion maturity, and their interaction effect on survival percentage of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	52.34	60.73	65.07	65.2	60.83
S ₂	65.43	67.53	84.04	85.96	75.74
S ₃	87.08	88.86	89.65	91.99	89.39
S ₄	92.24	93.99	94.72	94.98	93.98
Mean	74.27	77.77	83.37	84.53	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.910		0.910		1.821
CD(p=0.05)	1.855		1.855		3.710

Table 5: Influence of the size of the seed, scion maturity, and their interaction effect on the leaf area of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	15.16	15.68	16.22	17.06	16.03
S ₂	16.91	17.44	17.47	18.15	17.49
S ₃	17.92	18.72	18.92	19.20	18.69
S ₄	19.18	19.88	20.10	20.83	19.99
Mean	17.29	17.93	18.17	18.81	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.188		0.188		0.376
CD(p=0.05)	0.383		0.383		0.767

Table 6: Influence of the size of the seed, scion maturity, and their interaction effect on the carbohydrate content of Avocado

Size of seed	Scion maturity				Mean
	D ₁	D ₂	D ₃	D ₄	
S ₁	5.19	5.28	5.55	5.72	5.43
S ₂	5.72	5.77	5.82	6.05	5.84
S ₃	6.09	6.10	6.12	6.13	6.11
S ₄	6.26	6.29	6.30	6.57	6.35
Mean	5.81	5.86	5.94	6.11	
	Seed size		Scion maturity		Interaction (S X D)
SE(d)	0.055		0.055		0.110
CD(p=0.05)	0.112		0.112		0.225

Conclusion

According to the experiment's findings, for softwood grafting the seeds that are 60–70 g in weight and scions that are 90 days old are optimal for propagating avocados, followed by seeds that are 50–60 g in weight and scions that are 80 days old or older. Farmers have trouble propagating avocados because there aren't any standard rootstock and scion available. Therefore, standardizing rootstock and scion maturities for softwood grafting will enable the large-scale manufacture of high-quality planting material to farmers' advantage. It will also boost growth and yield, which will increase the farmers' revenue.

According to the results of the experiment, seeds with a weight of 60-70 g and scions that are 90 days old are best for propagating avocados, followed by seeds with a weight of 50-60 g and scions that are 80 days old or older. Farmers have difficulty growing avocados because conventional rootstock and scion are unavailable. As a result, standardized rootstock and scion maturities for softwood grafting will enable farmers to benefit from large-scale production of high-quality planting material. It will also promote growth and yield, increasing farmers' earnings.

REFERENCES

- Adjei, P. Y., Banful, B. K., & Idun, I. A. (2011). Seed size and seed cut-length effects on germination behavior and seedling growth of avocado (*Persea americana* Merr.). *International Journal of Agricultural Research*, 6(3), 299-305.
- Alam, M. A., Islam, M. S., Uddin, M. Z., Barman, J. C., & Quamruzzaman, A. K. M. (2006). Effect of age of seedling and variety of scion in stone grafting of mango. *Int. J. Sustain. Crop Prod*, 1(2), 27-32.
- Ben-Ya'acov, A., & Michelson, E. (1995). Avocado rootstocks. *HORTICULTURAL REVIEWS-WESTPORT THEN NEW YORK*, 17, 381-429.

- Córdula, E., Morim, M. P., & Alves, M. (2014). Morfologia de frutos e sementes de Fabaceaeocorrentesemumaáreaprioritária para a conservação da Caatingaem Pernambuco, Brasil. *Rodriguésia*, 65, 505-516.
- Ghosh, S. P. (2000). Avocado production in India. *Avocado production in Asia and the Pacific*, 24-30.
- Karna, A. K., Varu, D. K., Panda, P. A., &Hota, D. (2017). Standardization of grafting time and height on success of softwood grafting in mango (*Mangiferaindica* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(5), 2803-2807.
- Lazare, S., Haberman, A., Yermiyahu, U., Erel, R., Simenski, E., & Dag, A. (2020). Avocado rootstock influences scion leaf mineral content. *Archives of Agronomy and Soil Science*, 66(10), 1399-1409.
- Maheswari, T. U., &Nivetha, K. (2015). Effect of age of the rootstock on the success of softwood grafting in jack (*Artocarpusheterophyllus* Lam.). *Plant Arch*, 15, 823-825.
- Mejía-Jaramillo, L., Barrera-Sánchez, C., & Córdoba-Gaona, O. (2022). Effect of the seed weight on the growth of young avocado rootstock seedlings. *Bioagro*, 34(2), 183-194.
- Mickelbart, M. V., Robinson, P. W., Witney, G., &Arpaia, M. L. (2012). 'Hass' avocado tree growth on four rootstocks in California. II. Shoot and root growth. *Scientia horticulturae*, 143, 205-210.
- Ndoro, L. L. (2018). Effect of seed size on germination and seedling performance on grafted avocado.
- Nimbalkar, S. D., Haldankar, P. M., Pujari, K. H., Dhekale, J. H., &Somavanshi, A. V. (2011). Effect of maturity of scion and polythene bag cover on softwood grafting of karonda (*Carissa carandus* L.). *Indian Journal of Horticulture*, 68(3), 429-432.
- Olalde-Lira, G. G., Raya Montañó, Y. A., Apérez Barrios, P., Vargas-Sandoval, M., Pedraza Santos, M. E., Raymundo, T., ...& Blanca, M. (2020). Caracterización de *Fusarium* spp., fitopatógeno de aguacate (*Perseaamericana* Miller var. *drymifolia* (Schltdl. y Cham.)) en Michoacán, México. *Revista de la Facultad de CienciasAgrarias. Universidad Nacional de Cuyo*, 52(2), 301-316.
- Olorunmaiye, K. S., Fatoba, P. O., Olorunmaiye, P. M., &OreOluwaAdeyemi, C. (2012). Seed weight and Morpho-physiological parameters in Mango(*Mangiferaindica*) seedlings. *Pomologiacroatica*, 17(3-4), 77-86.
- PAIXÃO, M. V. S., Lopes, J. C., Schmildt, E. R., Alexandre, R. S., &Meneghelli, C. M. (2016). Avocado seedlings multiple stems production. *RevistaBrasileira de Fruticultura*, 38, e-221.
- Panase, V. G., &Sukhatme, P. V. (1954). Statistical methods for agricultural workers. *Statistical methods for agricultural workers*.
- Raghavendra, V. N., Angadi, S. G., Mokashi, A. N., Allolli, T. B., Venugopal, C. K., &Mummigatti, U. V. (2011). Studies on Softwood Grafting in Wood Apple (*Feronia limonio* L.). *Actahorticulturae*, (890), 165.
- Rajatiya, P. H., Kanzaria, D. R., DishaThanki, P. J., Patel, H. N., Senjaliya, H. J., BhadarkaChandni, R., &Parmar, V. M. (2022). Success of guava (*Psidiumguajava* L.) Grafts affected by rootstock height and scion stick length.
- Souza, J. A. C., Prates, G. C., Ferreira, J. T., Castro, G. D. M. D., &Salomão, L. C. C. (2020). Avocado: Is it possible to produce two seedlings with one seed?. *Revista Ceres*, 67, 152-155.
- TAMAYO-VÉLEZ, Á. D. J., Bernal-Estrada, J. A., Díaz-Díez, C. A., Cano-Gallego, L. E., & CÓRDOBA-GAONA, O. D. J. (2022). Effect of rootstock/scion compatibility on fruit and foliar nutrient composition in avocado (*Perseaamericana* Mill.) cv. Hass in Colombia. *RevistaColombiana de CienciasHortícolas*, 16(3).
- Tanuja, P., &Thippesha, D. (2017). Effect of different age groups of scion on success rate of softwood grafting in sapota. *Journal of Pharmacognosy and Phytochemistry*, 6(4), 1886-1889.
- Tesfay, S. Z., Bertling, I., Bower, J. P., &Lovatt, C. (2012). The quest for the function of 'Hass' avocado carbohydrates: clues from fruit and seed development as well as seed germination. *Australian Journal of Botany*, 60(1), 79-86.