

A study of Phytochemical analysis and Pharmacological activities of *Withania somnifera*

Abstract

Ashwagandha roots (*Withania somnifera* L. Dunal) have a history of utilization in Ayurveda to address conditions such as fever, asthma, arthritis, rheumatism, inflammation, tuberculosis, mental disorders, and male sexual issues. Through phytochemical analyses, it has been determined that the plant predominantly contains alkaloids (such as *withanine*, *sominiferine*, *tropine*, *isopelletierine*, and *anaferine*) and steroidal lactones (including withanolides A-Y, withaferine A, *withasominiferols* A-C, *withanone*, and *sitoindosides*) as the active compounds. These constituents and extracts exhibit various pharmacological effects, encompassing antioxidant, antitumor, antimicrobial, antivenom, and anti-parkinsonian properties. This review provides an up-to-date overview of *ashwagandha*, focusing on its phytochemistry and pharmacology. The plant and numerous secondary metabolites have displayed efficacy in ameliorating diverse human ailments. Nonetheless, additional research is imperative to ascertain the precise mechanisms underlying their actions.

Keywords

Withania somnifera, *Ashwagandha*, Phytochemical analysis, Pharmacological activities, Medicinal plants, Traditional medicine, Ecological significance.

Introduction

Ashwagandha (*Withania somnifera* L. Dunal), also known as Indian ginseng or winter cherry, originates from northwestern India [1]. Its name translates to "horse's smell" in Sanskrit, referring to the root's fragrance. It is found in India, Pakistan, Sri Lanka, Afghanistan, Egypt, South Africa, Morocco, and Jordan. In India, it's cultivated mainly in Uttar Pradesh, Madhya Pradesh, Punjab, Gujarat, and Rajasthan [2]. The Latin name "*somnifera*" means "sleep-inducing" [3]. Used in Asian medicine for 3000+ years, *ashwagandha* treats various health issues [5]. It's called Amukkira, Amukkiram, Punir, Asgandh, Akshan, Tilli in India [4]. Its powdered root, consumed daily, aims to delay aging, rejuvenate organs, and enhance fertility [6]. *Ashwagandha* leaves are used in Ayurveda and Unani for tumors [5]. It's part of 200+ Indian folk formulations, treating issues like asthma, inflammation, insomnia, anxiety, and more [7-13]. The root extracts (withanolides) possess antioxidant, immunomodulatory, anti-aging, and other properties [7-13]. Traditionally, boiled *ashwagandha* root powder with milk was used for female sterility [14]. It enhances vitality, strength, and bodily fluids [15]. *Ashwagandha* fruits aid digestion, and infant growth, and have sedative qualities [16]. Leaves address swelling, fever, and ophthalmitis [17]. The plant is useful for geriatric problems, stress, and arthritis [18, 19]. In modern medicine, *W. somnifera* extracts enhance brain health, counteract aging, and help with conditions like weakness, epilepsy, memory loss, and neurodegenerative diseases [20-22]. *Ashwagandha* extracts are dietary supplements globally [21]. Studies confirm its aphrodisiac potential and impact on testosterone [23-28]. This review highlights *Ashwagandha's* phytochemical and therapeutic benefits against various diseases.

Phytochemical Analysis

Ashwagandha roots contain diverse phytochemicals like amino acids, alkaloids, ergostane steroids, terpenes, and flavonoids, offering disease-mitigating potential [19, 26]. Essential bioactive compounds include withaferin A, withanone, and withanolides, with therapeutic effects such as antioxidant, antidiabetic, antimicrobial, anticancer, and more [29-41]. Withaferin A inhibits angiogenesis and counters cancer [42]. Sitoindoside-IX and -X enhance memory and reduce stress [43]. Nutritional composition (roots, leaves, fruits) is shown in Table 1, including minerals like calcium, copper, zinc, carotenoids, and vitamin C [44-46]. Geographic factors affect phytochemical composition [44]. Relevant metabolites include flavonoids [47]. Indian-grown *ashwagandha* roots contain alkaloids (0.13–0.31%) [48]. Alkaloids and steroidal lactones are major constituents. Alkaloids: *withanine*, *pseudowithanine*, etc. Steroidal lactones: *withanolides* A-Y, *withaferin* A, and more [49-52]. Molecular structures are in Table 2.

Pharmacological Activities





The pharmacological effects of *W. somnifera* have been comprehensively evaluated in preceding articles [2, 5, 11, 50, 53-57], encompassing research conducted up until 2015. In recent times, two additional reports [58, 59] have also addressed specific facets of *ashwagandha's* pharmacological activities, spanning up to 2020. To provide an updated perspective on this pivotal

Ayurvedic herb, we have undertaken a review of pharmacological studies conducted between 2015 and 2020, which were not encompassed in earlier reviews. This review aims to analyze the advancements made during these years compared to the preceding ones. The extracts and bioactive components derived from *ashwagandha* exhibit a diverse array of pharmacological effects, including antioxidative, anticancer, and immunomodulatory activities. A compilation of the significant pharmacological activities attributed to *ashwagandha* is presented in Table 3.

Compositions	Proximate[%]			Reference
	Fruit	Roots	Leaves	
Protein	11.0	1.6	5.3	[44]
Ash	9.1	3.7	8.6	
Carbohydrates	55.9	64.4	51.5	
Crude Fiber	4.0	5.0	2.3	
Fat	2.9	2.4	1.1	
Minerals(mg/100g)				
Copper	4.2	0.8-3.3	3.5	[45,46]
Calcium	-	23	-	
Iron	60.2	74.0	94.5	
Carotenoids	-	7.6	-	
Vitamin C	-	3.7	-	
Manganese	3.7	1.2-5.9	3.4	
Zinc	4.0	1.6-4.4	3.6	
Metabolites(mg/100g)				
Alkaloids				
Berberine	-	0.41	-	[47]
Harmine	-	0.26	-	
Caffeine	-	1.22	-	
Papaverine	-	0.16	-	
Noscapine	-	0.32	-	
Theobromine	-	0.26	-	
Flavonoids				
Kaempferol	-	0.78	-	
Myricetin	-	0.22	-	

Rutin	-	4.21	-	[47]
Quercetin	-	7.21	-	
Quercitrin	-	5.22	-	
Rhamnetin	-	1.15	-	
Phenolic acids				
Coumaric acid	-	0.67	-	[47]
Caecic acid	-	1.99	-	
Chlorogenic acid	-	1.03	-	
Ferulic acid	-	0.55	-	
Gallic acid	-	4.02	-	

Table 1
Proximate, mineral and metabolites concentration of *Withania somnifera*

Bioactive constituents	Chemical structures	Bioactive constituents	Chemical structures
Cuscohygrine		Withanone	
Anahygrine		Withanolide A	







Trophine		Sominone	
Steroidal Lactone		Withanoside IV	
Withaferin A		Ashwagandhaide	

Table 2

Chemical structures of the major bioactive constituents of *Withania somnifera*

Activity	Extract Type	Methods Used	IC ₅₀ /dosage	References
Adaptogenic and anxiolytic activity	Root extract	Fiy-eight male and female participants with a baseline perceived stress scale (PSS) score >20	250 and 600 mg/twice/day, 8 weeks	[69]
Anticancer	Methonol ectract	HepG2 hepatoma cell line	IC ₅₀ :1.89 µg/ml	[65]
Anticancer	withaferin A	Transgenic adenocarcinoma of mouse prostate(TRAMP) model	5 mg/kg/day; 8 week	[64]

Activity	Extract Type	Methods Used	IC ₅₀ /dosage	References
Adaptogenic and anxiolytic activity	Root extract	Fiy-eight male and female participants with a baseline perceived stress scale (PSS) score >20	250 and 600 mg/twice/day, 8 weeks	[69]
Anticancer	Methonol ectract	HepG2 hepatoma cell line	IC ₅₀ :1.89 µg/ml	[65]
Anticancer	Methonol ectract	HCT 116 human colorectal cell line	IC ₅₀ :2.19 µg/ml	[65]
Anticancer	<i>W. somnifera</i> prot fraction	MDA-MB-231 human breast cancer cells	IC ₅₀ :92 µg/ml	[63]
Anti-neuro inflammatory	Leaf extract	<i>Ashwagandha</i> leaf water extract using β-amyloid and lipopolysaccharide (LPS)- stimulated primary microglial cells and BV-2 microglial cell line	IC ₅₀ :0.2%; 48h	[61]
Anti-neuro inflammatory	Chloroform fractio	Chloroform fraction IV (FIV) using β-amyloid an lipopolysaccharide (LPS)- stimulated primary microglial cells and BV-2 microglial cell line	IC ₅₀ :15 µg/ml; 48h	[61]
Antioxidant activity	Extract type root extract	Lipid peroxides (LPO), superoxide dismutase (SC and glutathione (GSH) were tested in adult male Wistar rats	500 mg/kg, p.o once a day from day 14 to day 21	[60]
Antiarthritic activity	Root extract	Assessed the level of inflammatory cytokines such as Tumor Necrosis Factor (TNF)-α, IL-1β, IL-6 and IL-10 in collagen-induced arthritic (CIA) rats	300 mg/kg/day; 20 day	[62]
Body strength	aqueous extract (Sensoril®)	19 recreationally active men were randomized in a double-blind fashion to placebo or Sensoril®	500 mg/day; 12 week	[67]
Cytotoxic activity	Root extract	The crude extract of <i>Withania</i> was tested for cytotoxicity against A375 cells by MTT assay	350, 250 and 200 µg/m for 24, 48 and 72 h, respectively	[68]
Insomnia and anxie	Root extract	60 patients were randomly divided into two group ashwagandha (n=40) and placebo (n=20)	300 mg, 2 doses/ day; weeks	[72]
Increased testosterone level	withanolide glycosides	43 people completed the 16-week period of trial	21 mg/day for 8 weeks	[71]
Insomnia	Root extract	150 subjects divided into two groups: ashwagandh (n=75) and placebo (n=75)	600 mg/once/ day; 42 days	[73]
Nephroprotective	Root extract	Gentamycin-induced nephrotoxicity in adult male Wistar rats	500 mg/kg, p.o once a day from day 14 to day 21	[60]

Table 3

Pharmacological activities of *W. somnifera* root extracts and alkaloids

LC50 – lethal concentration; IC50 – inhibitory concentration

Neuroprotective activity

Elhadidy et al. [74] found *ashwagandha* countered aluminum-induced neurotoxicity (200 mg/kg p.o.). Dutta et al. [75] noted *ashwagandha* improved motor function in ALS mice and inhibited glial activation. *Ashwagandha* co-treatment with toxins mitigated Parkinson's markers [12, 13]. Limited animal models studied *ashwagandha's* nootropic potential; human trials are needed.

Ashwagandha inhibits various cancers [76-81]. Lee et al. [82] noted apoptotic effects on HNSCC cells. Daily *ashwagandha* (200 mg/kg) reduced lung adenomas [83]. Withaferin A impacted NSCLC cells [84]. It reduced Helicobacter pylori-induced IL-1 [85]. *Ashwagandha* protein fraction induced apoptosis in breast cancer cells [63]. Few studies explored *ashwagandha's* anticancer potential. Future research should rigorously evaluate its efficacy through clinical trials with appropriate controls and dosages.

Antimicrobial activity

Ashwagandha roots display potent antimicrobial properties. Methanolic leaf extracts inhibit gram-positive bacteria like Staphylococcus aureus, and Enterococcus sp. (zones of inhibition: 20.6 mm, 19.4 mm) [86]. A recent study found *ashwagandha* extract inhibiting S. aureus, and E. coli (17 mm to 24 mm zones) [87]. Leaf extracts (6.25 mg/ml, 12.5 mg/ml) hindered the growth of gram-negative bacteria (E. coli, S. typhi, K. pneumonia, P. aeruginosa, C. freundii) [88]. Extract (100 mg/ml) suppressed Staphylococcus aureus, Streptococcus sobrinus, Streptococcus mutans, Salmonella paratyphi B [89, 90]. Most studies use disc diffusion, but limitations exist. Further Minimum Inhibitory Concentration (MIC) assays are necessary to enhance comprehension [91].

Anti-depressant activity

In a six-week trial, 120 mg/day *ashwagandha* root extract improved sleep quality significantly in 150 participants compared to placebo (p<0.001) [92]. The treatment group showed better sleep efficiency, total sleep time, latency, and wake after sleep onset. Salve et al. [69] gave 250 mg/day, and 600 mg/day root extract for eight weeks. Perceived stress and cortisol levels were reduced in 58 participants vs. placebo. Another study using 240 mg/day for 60 days showed stress reduction [28]. Earlier studies demonstrated *ashwagandha* (20, 40, 50 mg/kg/once daily for five days) reduced anxiety in animal models [34, 93].

Anti-inflammatory activity:

A 47% methanolic extract curbed LPS-induced nitric oxide and TNF- α production [94]. Aqueous *ashwagandha* root extract (300 mg/kg/wt.) enhanced IL-10 secretion and suppressed NF- κ B [62]. Doses of 600, and 800 mg/kg mitigated arthritis severity, inhibiting inflammatory mediators [95]. Lower doses (100 mg/kg, 15 days) alleviated arthritis, surpassing hydrocortisone's effect [96]. *Ashwagandha* inhibited granuloma formation [97].

Spermatogenic Activity:

Ashwagandha enhances semen quality by reducing oxidative stress, and elevating hormone levels [23, 26, 71, 98]. 300 mg/kg for 8 weeks improved sperm count, and motility [99]. 100 mg/kg for 30 days increased sperm count, and motility [100]. 200 mg/kg reduced sperm abnormalities [101].

Cultivation of *Withania somnifera*:

Field cultivation yields more fruits, and seeds than greenhouse. Early seedling preparation is crucial for ample fruit production. Withaferin A, and withanolide D were detected in both conditions [102]. Soil salinity affects growth [103]. Enhanced ZnSO₄ leads to increased Withaferin A, and glutathione [104].

Limitations and Future Directions:

This review acknowledges limitations in some studies' experimental designs, emphasizing the need for proper control groups and clear dosages for accurate interpretation. Both organic and aqueous *ashwagandha* extracts deserve exploration to understand their potential fully. While animal and cellular models have shown *ashwagandha's* therapeutic effects, more human clinical trials are essential for translation. Research should expand to human trials involving extracts, alkaloids, and steroidal lactones. Further investigation into bioactive constituents, their bioavailability, and pharmacokinetics is needed to understand their

roles. *Ashwagandha's* alkaloids and steroidal lactones hold pharmaceutical promise. Ethnobotanical and modern medicinal uses should be explored comprehensively to maximize their potential benefits for human health.

Conclusion:

In conclusion, this review emphasizes *Ashwagandha's* ethnopharmacological applications, phytochemical composition, and potential therapeutic benefits for various human conditions. The literature supports its efficacy against ulcers, insomnia, memory impairment, anxiety, bronchitis, and neurological disorders. Bioactive compounds like *withaferin* and *sitoinosides* exhibit cellular protection and counteract illnesses. Delving into ashwagandha extracts and constituents reveals positive impacts. Yet, existing gaps in research need attention for validation and further exploration.

References

1. Andallu B, Radhika B. "Hypoglycemic diuretic and hypocholesterolemic effect of winter cherry (*Withania somnifera* Dunal) root." *Indian J Exp Biol*, 2000; 38:607-609.
2. Kulkarni SK, Dhir A. "*Withania somnifera*: an Indian ginseng." *Prog Neuro-Psychopharmacol Biol Psychiatry*, 2008; 32(5):1093-1105. [DOI: 10.1016/j.pnpbp.2007.09.011](https://doi.org/10.1016/j.pnpbp.2007.09.011)
3. Stearn WT. "Botanical Latin: History, Grammar, Syntax, Terminology and Vocabulary." 4th edition. Portland, Ore, USA: Timber Press, 1995.
4. Pandian A, Ashokkumar K, Sekar S, Sivakumar P, Selvaraj KSV, Karthik M, Hariprasath L. "Botany and ethnopharmacological potential of *ashwagandha*." *J Curr Opin Crop Sci*, 2020; 1(1):35–40.
5. Singh G, Sharma PK, Dudhe R, Singh S. "Biological activities of *Withania somnifera*." *Ann Biol Res*, 2010; 1(3):56–63.
6. Durg S, Shivaram SB, Bavage S. "*Withania somnifera* (Indian ginseng) in male infertility: An evidence-based systematic review and meta-analysis." *Phytomedicine*, 2018; 50:247-256. [DOI: 10.1016/j.phymed.2017.11.011](https://doi.org/10.1016/j.phymed.2017.11.011)
7. Ziauddin M, Phansalkar N, Patki P, Diwanay S, Patwardhan B. "Studies on the immunomodulatory effects of *ashwagandha*." *J Ethnopharmacol*, 1996; 50:69-76. [DOI: 10.1016/0378-8741\(95\)01318-08](https://doi.org/10.1016/0378-8741(95)01318-08)
8. Rasool M, Varalakshmi P. "Immunomodulatory role of *Withania somnifera* root powder on experimental induced inflammation: An in vivo and in vitro study." *Vascul Pharmacol*, 2006; 44:406-410. [DOI: 10.1016/j.vph.2006.01.015](https://doi.org/10.1016/j.vph.2006.01.015)
9. Gupta M, Bisht D, Pandey MM, Ojha SK, Khatoon S, Rastogi S, Rawat AKS. "Standardization of Ashwagandhilehya – An important ayurvedic formulation of *Withania somnifera*." *India J Tradit Know*, 2011; 10:594-598.
10. Dhuley JN. "Nootropic-like effect of *ashwagandha* (*Withania somnifera* L.) in mice." *Phytother Res*, 2001; 15:524-528. [DOI: 10.1002/ptr.874](https://doi.org/10.1002/ptr.874)
11. Mishra LC, Singh BB, Dagenais S. "Scientific basis for the therapeutic use of *Withania somnifera* (*ashwagandha*): A review." *Altern Med Rev*, 2000; 5:334-346.
12. Prakash J, Yadav SK, Chouhan S, Singh SP. "Neuroprotective role of *Withania somnifera* root extract in maneb-paraquat induced mouse model of parkinsonism." *Neurochem Res*, 2013; 38:972-980. [DOI: 10.1007/s11064-013-1005-4](https://doi.org/10.1007/s11064-013-1005-4)
13. Prakash J, Chouhan S, Yadav SK, Westfall S, Rai SN, Singh SP. "*Withania somnifera* alleviates parkinsonian phenotypes by inhibiting apoptotic pathways in dopaminergic neurons." *Neurochem Res*, 2014; 39:2527-2536. [DOI: 10.1007/s11064-014-1443-7](https://doi.org/10.1007/s11064-014-1443-7)
14. Kirtikar KR, Basu BD. "Indian Medicinal Plants." Vol. 3, International Book Distributors Book Sellers and Publishers, Dehradun, India, 1999.
15. Williamson EM. "Major herbs of Ayurveda." Churchill Livingstone, London, UK, 2002, pp. 322-323.
16. Watt GA. "Dictionary of the Economic Products of India." Cosmo Publication, Delhi, India, 1972.
17. Patwardhan B, Panse GT, Kulkarni PH. "*Ashwagandha*: a review." *J Natl Integr Med Assoc*, 1998; 30:7–11.
18. Mirjalili MH, Moyano E, Bonfill M, Cusido RM, Palazon J. "Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine." *Molecules*, 2009; 14:2373–2393. [DOI: 10.3390/molecules14072373](https://doi.org/10.3390/molecules14072373)
19. Alam N, Hossain M, Khalil MI, Moniruzzaman M, Sulaiman SA, Gan SH. "High catechin concentrations detected in *Withania somnifera* (*ashwagandha*) by high-performance liquid chromatography analysis." *BMC Complem Altern Med*, 2011; 11:65. [DOI: 10.1186/1472-6882-11-65](https://doi.org/10.1186/1472-6882-11-65)
20. Singh RH, Narsimhamurthy K, Singh G. "Neuronutrient impact of Ayurvedic Rasayana therapy in brain aging." *BioGerontology*, 2008; 9:369-374. [DOI: 10.1007/s10522-008-9185-z](https://doi.org/10.1007/s10522-008-9185-z)
21. Rajasankar S, Manivasagam T, Sankar V, Prakash S, Muthusamy R, Krishnamurti A, Surendran S. "*Withania somnifera* root extract improves catecholamines and physiological abnormalities seen in a Parkinson's disease model mouse." *J Ethnopharmacol*, 2009; 125:369-373. [DOI: 10.1016/j.jep.2009.08.003](https://doi.org/10.1016/j.jep.2009.08.003)

22. Kuboyama T, Tohda C, Komatsu K. "Effects of *ashwagandha* (roots of *Withania somnifera*) on neurodegenerative diseases." *Biol Pharm Bull*, 2014; 37:892-897. [DOI: 10.1248/bpb.b14-00022](https://doi.org/10.1248/bpb.b14-00022)
23. Ahmad MK, Mahdi AA, Shukla KK, Islam N, Rajender S, Madhukar D, Shankhwar SN, Ahmad S. "*Withania somnifera* improves semen quality by regulating reproductive hormone levels and oxidative stress in seminal plasma of infertile males." *Fertil Steril*, 2010; 94:989-996. [DOI: 10.1016/j.fertnstert.2009.04.046](https://doi.org/10.1016/j.fertnstert.2009.04.046)
24. Mahdi AA, Shukla KK, Ahmad MK, Rajender S, Shankhwar SN, Singh V, Dalela D. "*Withania somnifera* improves semen quality in stress-related male fertility." *Evid Based Complement Alternat Med*, 2011; 576962. [DOI: 10.1093/ecam/nep138](https://doi.org/10.1093/ecam/nep138)
25. Mamidi P, Thakar AB. "Efficacy of *ashwagandha* (*Withania somnifera* Dunal. Linn.) in the management of psychogenic erectile dysfunction." *Ayu*, 2011; 32:322-328. [DOI: 10.4103/0974-8520.93907](https://doi.org/10.4103/0974-8520.93907)
26. Ambiye VR, Langade D, Dongre S, Aptikar P, Kulkarni M, Dongre A. "Clinical evaluation of the spermatogenic activity of the root extract of *ashwagandha* (*Withania somnifera*) in oligospermic males: a pilot study." *Evid Based Complement Alternat Med*, 2013; 2013:571420. [DOI: 10.1155/2013/571420](https://doi.org/10.1155/2013/571420)
27. Nipanikar SU, Nagore DH, Chitlange SS. "Evaluation of aphrodisiac activity of AHPL/AYCAP/0114 capsule in sexually sluggish male rats." *Phcog Mag*, 2018; 14:264-267. [DOI: 10.4103/pm.pm_363_17](https://doi.org/10.4103/pm.pm_363_17)
28. Lopresti AL, Smith SJ, Malvi H, Kodgule R. "An investigation into the stress-relieving and pharmacological actions of an *ashwagandha* (*Withania somnifera*) extract: A randomized, double-blind, placebo-controlled study." *Medicine*, 2019; 98(37):e17186. [DOI: 10.1097/MD.00000000000017186](https://doi.org/10.1097/MD.00000000000017186)
29. Sharma S, Dahunukar S, Karandikar SM. "Effects of long term administration of the roots of *ashwagandha* and shatavari in rats." *Indian Drugs*, 1985; 23:133-139.
30. Grandhi A, Mujumdar AM, Patwardhan B. "A comparative pharmacological investigation of *ashwagandha* and ginseng." *J Ethnopharmacol*, 1994; 44:131-135. [DOI: 10.1016/0378-8741\(94\)01119-2](https://doi.org/10.1016/0378-8741(94)01119-2)
31. Bhattacharya SK, Satyan KS, Chakrabarti A. "Effect of Trasina, an Ayurvedic herbal formulation, on pancreatic islet superoxide dismutase activity in hyperglycaemic rats." *Indian J Exp Biol*, 1997; 35:297-299.
32. Dhuley JN. "Adaptogenic and cardioprotective action of *ashwagandha* in rats and frogs." *J Ethnopharmacol*, 2000; 70:57-63. [DOI: 10.1016/S0378-8741\(99\)00177-4](https://doi.org/10.1016/S0378-8741(99)00177-4)
33. Davis L, Kuttan G. "Immunomodulatory activity of *Withania somnifera*." *J Ethnopharmacol*, 2000; 71:193-200. [DOI: 10.1016/S0378-8741\(99\)00206-8](https://doi.org/10.1016/S0378-8741(99)00206-8)
34. Bhattacharya SK, Bhattacharya A, Sairam K, Ghosal S. "Anxiolytic antidepressant activity of *Withania somnifera* glycowithanolides: an experimental study." *Phytomedicine*, 2000; 7:463-469. [DOI: 10.1016/S0944-7113\(00\)80030-6](https://doi.org/10.1016/S0944-7113(00)80030-6)
35. Bhattacharya A, Ghosal S, Bhattacharya SK. "Anti-oxidant effect of *Withania somnifera* glycowithanolides in chronic footshock stress-induced perturbations of oxidative free radical scavenging enzymes and lipid peroxidation in rat frontal cortex and striatum." *J Ethnopharmacol*, 2001; 74:1-6. [DOI: 10.1016/S0378-8741\(00\)00309-3](https://doi.org/10.1016/S0378-8741(00)00309-3)
36. Girish KS, Machiah KD, Ushanandini S, Kumar HK, Nagaraju S, Govindappa M, Vedavathi M, Kemparaju K. "Antimicrobial properties of a non-toxic glycoprotein (WSG) from *Withania somnifera* (*ashwagandha*)." *J Basic Microbiol*, 2006; 46:365-374. [DOI: 10.1002/jobm.200510108](https://doi.org/10.1002/jobm.200510108)
37. Fong MY, Jin S, Rane M, Singh RK, Gupta R, Kakar SS. "Withaferin A synergizes the therapeutic effect of doxorubicin through ROS-mediated autophagy in ovarian cancer." *PloS One*, 2012; 7(7):e42265. [DOI: 10.1371/journal.pone.0042265](https://doi.org/10.1371/journal.pone.0042265)
38. Sun GY, Li R, Cui J, Hannink M, Gu Z, Fritsche KL, Lubahn DB, Simonyi A. "*Withania somnifera* and its withanolides attenuate oxidative and inflammatory responses and up-regulate antioxidant responses in BV-2 microglial cells." *Neuromolecular Med*, 2016; 18:241-252. [DOI: 10.1007/s12017-016-8411-0](https://doi.org/10.1007/s12017-016-8411-0)
39. Ahmed W, Mofed D, Zekri AR, El-Sayed N, Rahouma M, Sabet S. "Antioxidant activity and apoptotic induction as mechanisms of action of *Withania somnifera* (*ashwagandha*) against a hepatocellular carcinoma cell line." *J Int Med Res*, 2018; 46:1358-1369. [DOI: 10.1177/0300060517752022](https://doi.org/10.1177/0300060517752022)
40. Chukwuma CI, Matsabisa MG, Ibrahim MA, Erukainure OL, Chabalala MH, Islam MS. "Medicinal plants with concomitant anti-diabetic and anti-hypertensive effects as potential sources of dual-acting therapies against diabetes and hypertension: A review." *J Ethnopharmacol*, 2019; 235:329-360. [DOI: 10.1016/j.jep.2019.02.024](https://doi.org/10.1016/j.jep.2019.02.024)
41. Duda-Chodak A, Tarko T, Rus M. "Antioxidant activity and total polyphenol content of selected herbal medicinal products used in Poland." *Herba Pol*, 2011; 57(1):48-57.
42. Mohan R, Hammers HJ, Bargagna-Mohan P, Zhan XH, Herbstritt CJ, Ruiz A, Zhang L, et al. "Withaferin A is a potent inhibitor of angiogenesis." *Angiogenesis*, 2004; 7:115-122. [DOI: 10.1023/B:AGEN.0000037331.82501.2e](https://doi.org/10.1023/B:AGEN.0000037331.82501.2e)
43. Ghosal S, Lal J, Srivastava R, Bhattacharya SK, Upadhyay SN. "Immunomodulatory and CNS effects of sitoindosides IX and X, two new glycowithanolides from *Withania somnifera*." *Phytother Res*, 1989; 3:201-206. [DOI: 10.1002/ptr.2650030510](https://doi.org/10.1002/ptr.2650030510)

44. Krishnamurthy SR, Sarala P. "Proximate nutritive values and mineral components of *Withania somnifera* (Linn.) Dunal." J Chem, 2010; 7:616-851. [DOI: 10.1155/2010/616851](https://doi.org/10.1155/2010/616851)
45. Gulati S, Madan VK, Singh S, Singh I, Dusyant. "Chemical and phytochemical composition of *ashwagandha* (*Withania somnifera* L.) roots." Asian J Chem, 2017; 29:1683-1686. [DOI: 10.14233/ajchem.2017.20536](https://doi.org/10.14233/ajchem.2017.20536)
46. Kumari S, Gupta A. "Nutritional composition of dehydrated *ashwagandha*, shatavari, and ginger root powder." Int J Home Sci, 2016; 2:68-70.
47. Filipiak-Szok A, Kurzawa M, Szlyk E, Twarużek M, Błajet-Kosicka A, Grajewski J. "Determination of mycotoxins, alkaloids, phytochemicals, antioxidants and cytotoxicity in Asiatic ginseng (*ashwagandha*, Dong quai, Panax ginseng)." Chem Pap, 2017; 71:1073-1082. [DOI: 10.1007/s11696-016-0028-0](https://doi.org/10.1007/s11696-016-0028-0)
48. Mirjalili MH, Moyano E, Bonfill M, Cusido RM, Palazon J. "Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine." Molecules, 2009; 14(7):2373-2393. [DOI: 10.3390/molecules14072373](https://doi.org/10.3390/molecules14072373)
49. Ganzera M, Choudhary MI, Khan IA. "Quantitative HPLC analysis of withanolides in *Withania somnifera*." Fitoterapia, 2003; 74:68-76. [DOI: 10.1016/S0367-326X\(02\)00325-8](https://doi.org/10.1016/S0367-326X(02)00325-8)
50. Gupta GL, Rana AC. "*Withania somnifera* (*ashwagandha*) a review." Pharmacogn Rev, 2007; 1:129-136. [DOI: 10.4103/0973-7847.33883](https://doi.org/10.4103/0973-7847.33883)
51. Matsuda H, Murakami T, Kishi A, Yoshikawa M. "Structures of withanosides I, II, III, IV, V, VI, and VII, new withanolide glycosides from the roots of Indian *Withania somnifera* DUNAL and inhibitory activity for tachyphylaxis to clonidine in isolated guinea-pig ileum." Bioorg Med Chem, 2001; 9:1499-1507. [DOI: 10.1016/S0968-0896\(01\)00024-4](https://doi.org/10.1016/S0968-0896(01)00024-4)
52. Kalra R, Kaushik N. "*Withania somnifera* (Linn.) Dunal: a review of chemical and pharmacological diversity." Phytochem Rev, 2017; 16:953-987. [DOI: 10.1007/s11101-017-9504-6](https://doi.org/10.1007/s11101-017-9504-6)
53. Winters M. "Ancient medicine, modern use: *Withania somnifera* and its potential role in integrative oncology." Altern Med Rev, 2006; 11(4):269-277.
54. Verma SK, Kumar A. "Therapeutic uses of *Withania somnifera* (*ashwagandha*) with a note on withanolides and its pharmacological actions." Asian J Pharm Clin Res, 2012; 4:1-4.
55. Uddin Q, Samiulla L, Singh VK, Jamil SS. "Phytochemical and pharmacological profile of *Withania somnifera* Dunai: A review." J Appl Pharm Sci, 2012; 2:170-175.
56. Kumar V, Dey A, Hadimani MB, Emerald M. "Chemistry and Pharmacology of *Withania somnifera*: An Update." Tang 2015; 5(1):1.1-1.13. [DOI: 10.5667/tang.2014.00305](https://doi.org/10.5667/tang.2014.00305)
57. Dar NJ, Hamid A, Ahmad M. "Pharmacological Overview of *Withania somnifera*, the Indian Ginseng." Cell Mol Life Sci 2015; 72:4445-4460. [DOI: 10.1007/s00018-015-2012-1](https://doi.org/10.1007/s00018-015-2012-1)
58. Tripathi N, Shrivastava D, Ahmad Mir B, Kumar S, Govil S, Vahedi M, Bisen PS. "Metabolomic and Biotechnological Approaches to Determine Therapeutic Potential of *Withania somnifera* (L.) Dunal: A Review." Phytomedicine 2018; 50:127-136. [DOI: 10.1016/j.phymed.2017.08.020](https://doi.org/10.1016/j.phymed.2017.08.020)
59. Pérez-Gómez J, Villafaina S, Adsuar JC, Merellano-Navarro E, Collado-Mateo D. "Effects of *Ashwagandha* (*Withania somnifera*) on VO₂max: A Systematic Review and Meta-Analysis." Nutrients 2020; 12(4):1119. [DOI: 10.3390/nu12041119](https://doi.org/10.3390/nu12041119)
60. Govindappa PK, Gautam V, Tripathi SM, Sahni YP, Raghavendra HLS. "Effect of *Withania somnifera* on Gentamicin-Induced Renal Lesions in Rats." Rev Bras Farmacogn 2019; 29:234-240. [DOI: 10.1016/j.bjp.2018.12.005](https://doi.org/10.1016/j.bjp.2018.12.005)
61. Gupta M, Kaur G. "Aqueous Extract from the *Withania somnifera* Leaves as a Potential Anti-Neuroinflammatory Agent: A Mechanistic Study." J Neuroinflammation 2016; 13(1):193. [DOI: 10.1186/s12974-016-0650-3](https://doi.org/10.1186/s12974-016-0650-3)
62. Khan MA, Ahmed RS, Chandra N, Arora VK, Ali A. "In Vivo, Extract from *Withania somnifera* Root Ameliorates Arthritis via Regulation of Key Immune Mediators of Inflammation in Experimental Model of Arthritis." Antiinflamm Antiallergy Agents Med Chem 2019; 18:55-70. [DOI: 10.2174/1871523017666181116092934](https://doi.org/10.2174/1871523017666181116092934)
63. Dar PA, Mir SA, Bhat JA, Hamid A, Singh LR, Malik F, Dar TA. "An Anti-Cancerous Protein Fraction from *Withania somnifera* Induces ROS-Dependent Mitochondria-Mediated Apoptosis in Human MDA-MB-231 Breast Cancer Cells." Int J Biol Macromol 2019; 135:77-87. [DOI: 10.1016/j.ijbiomac.2019.05.120](https://doi.org/10.1016/j.ijbiomac.2019.05.120)
64. Suman S, Das TP, Moselhy J, Pal D, Kolluru V, Alatassi H, Ankem MK, Damodaran C. "Oral Administration of Withaferin A Inhibits Carcinogenesis of Prostate in TRAMP Model." Oncotarget 2016; 7(33):53751-53761. [DOI: 10.18632/oncotarget.10733](https://doi.org/10.18632/oncotarget.10733)
65. Alfaiz MY, Saleh KA, El-Boushnak MA, Elbehairi SE, Alshehri MA, Shati AA. "Antiproliferative Activity of the Methanolic Extract of *Withania somnifera* Leaves from Faifa Mountains, Southwest Saudi Arabia, Against Several Human Cancer Cell Lines." Asian Pac J Cancer Prev 2016; 17:2723-2726.

66. Wankhede S, Langade D, Joshi K, Sinha SR, Bhattacharyya S. "Examining the Effect of *Withania somnifera* Supplementation on Muscle Strength and Recovery: A Randomized Controlled Trial." J Int Soc Sports Nutr 2015; 12:43. DOI: [10.1186/s12970-015-0104-9](https://doi.org/10.1186/s12970-015-0104-9)
67. Ziegenfuss TN, Kedia AW, Sandrock JE, Raub BJ, Kerksick CM, Lopez HL. "Effects of an Aqueous Extract of *Withania somnifera* on Strength Training Adaptations and Recovery: The STAR Trial." Nutrients 2018; 10(11):1807. DOI: [10.3390/nu10111807](https://doi.org/10.3390/nu10111807)
68. Halder B, Singh S, Thakur SS. "*Withania somnifera* Root Extract Has Potent Cytotoxic Effect Against Human Malignant Melanoma Cells." PLoS One 2015; 10(9):e0137498. DOI: [10.1371/journal.pone.0137498](https://doi.org/10.1371/journal.pone.0137498)
69. Salve J, Pate S, Debnath K, Langade D. "Adaptogenic and Anxiolytic Effects of *Ashwagandha* Root Extract in Healthy Adults: A Double-Blind, Randomized, Placebo-Controlled Clinical Study." Cureus 2019; 11(12):e6466. DOI: [10.7759/cureus.6466](https://doi.org/10.7759/cureus.6466)
70. Dey A, Chatterjee SS, Kumar V. "Triethylene Glycol-Like Effects of *Ashwagandha* (*Withania somnifera* (L.) Dunal) Root Extract Devoid of Withanolides in Stressed Mice." Ayu 2018; 39:230-238. DOI: [10.4103/avj.AYU.219.16](https://doi.org/10.4103/avj.AYU.219.16)
71. Lopresti AL, Drummond PD, Smith SJ. "A Randomized, Double-Blind, Placebo-Controlled, Crossover Study Examining the Hormonal and Vitality Effects of *Ashwagandha* (*Withania somnifera*) in Aging, Overweight Males." Am J Mens Health 2019; 13(2):1557988319835985. DOI: [10.1177/1557988319835985](https://doi.org/10.1177/1557988319835985)
72. Langade D, Kanchi S, Salve J, Debnath K, Ambegaokar D. "Efficacy and Safety of *Ashwagandha* (*Withania somnifera*) Root Extract in Insomnia and Anxiety: A Double-Blind, Randomized, Placebo-Controlled Study." Cureus 2019; 11(9):e5797. DOI: [10.7759/cureus.5797](https://doi.org/10.7759/cureus.5797)
73. Deshpande A, Irani N, Balakrishnan R. "Study Protocol and Rationale for a Prospective, Randomized, Double-Blind, Placebo-Controlled Study to Evaluate the Effects of *Ashwagandha* (*Withania somnifera*) Extract on Nonrestorative Sleep." Medicine 2018; 97(26):e11299. DOI: [10.1097/MD.00000000000011299](https://doi.org/10.1097/MD.00000000000011299)
74. Elhadidy ME, Sawie HG, Meguid NA, Khadrawy YA. "Protective Effect of *Ashwagandha* (*Withania somnifera*) Against Neurotoxicity Induced by Aluminum Chloride in Rats." Asian Pac J Trop Biomed 2018; 8:59-66. DOI: [10.4103/2221-1691.221139](https://doi.org/10.4103/2221-1691.221139)
75. Dutta K, Patel P, Julien JP. "Protective Effects of *Withania somnifera* Extract in SOD1G93A Mouse Model of Amyotrophic Lateral Sclerosis." Exp Neurol 2018; 309:193-204. DOI: [10.1016/j.expneurol.2018.08.008](https://doi.org/10.1016/j.expneurol.2018.08.008)
76. Prakash J, Gupta SK, Dinda AK. "*Withania somnifera* Root Extract Prevents DMBA-Induced Squamous Cell Carcinoma of Skin in Swiss Albino Mice." Nutr Cancer 2002; 42:91-97. DOI: [10.1207/S15327914NC421_1](https://doi.org/10.1207/S15327914NC421_1)
77. Jayaprakasam B, Zhang Y, Seeram N, Nair M. "Growth Inhibition of Tumor Cell Lines by Withanolides from *Withania somnifera* Leaves." Life Sci 2003; 74:125-132. DOI: [10.1016/j.lfs.2003.07.007](https://doi.org/10.1016/j.lfs.2003.07.007)
78. Senthilnathan P, Padmavathi R, Banu SM, Sakthisekaran D. "Enhancement of Antitumor Effect of Paclitaxel in Combination with Immunomodulatory *Withania somnifera* on Benzo(a)pyrene Induced Experimental Lung Cancer." Chem Biol Interact 2006; 159:180-185. DOI: [10.1016/j.cbi.2005.11.003](https://doi.org/10.1016/j.cbi.2005.11.003)
79. Singh G, Kumar P. "Evaluation of Antimicrobial Efficacy of Flavonoids of *Withania somnifera* L." Indian J Pharm Sci 2011; 73:473-478.
80. Yadav B, Bajaj A, Saxena M, Saxena AK. "In Vitro Anticancer Activity of the Root, Stem, and Leaves of *Withania somnifera* Against Various Human Cancer Cell Lines." Indian J Pharm Sci 2010; 72:659-663. DOI: [10.4103/0250-474X.78543](https://doi.org/10.4103/0250-474X.78543)
81. Nema R, Khare S, Jain P, Pradhan A. "Anticancer Activity of *Withania somnifera* Leaves Flavonoids Compound." Int J Pharm Sci Rev Res 2013; 19:103-106.
82. Lee HE, Shin JA, Jeong JH, Jeon JG, Lee MH, Cho SD. "Anticancer Activity of *Ashwagandha* Against Human Head and Neck Cancer Cell Lines." J Oral Pathol Med 2016; 45(3):193-201. DOI: [10.1111/jop.12353](https://doi.org/10.1111/jop.12353)
83. Dutta R, Khalil R, Green R, Mohapatra SS, Mohapatra S. "*Withania somnifera* (*Ashwagandha*) and Withaferin A: Potential in Integrative Oncology." Int J Mol Sci 2019; 20(21):5310. DOI: [10.3390/ijms20215310](https://doi.org/10.3390/ijms20215310)
84. Liu X, Chen L, Liang T, Tian XD, Liu Y, Zhang T. "Withaferin A Induces Mitochondrial-Dependent Apoptosis in Non-Small Cell Lung Cancer Cells via Generation of Reactive Oxygen Species." J Buon 2017; 22:244-250.
85. Kim JE, Lee JY, Kang MJ, Jeong YJ, Choi JA, Oh SM, Lee KB, Park JH. "Withaferin A Inhibits Helicobacter pylori-Induced Production of IL-1 β in Dendritic Cells by Regulating NF- κ B and NLRP3 Inflammasome Activation." Immune Netw 2015; 15:269-277.
86. Bisht P, Rawat V. "Antibacterial Activity of *Withania somnifera* Against Gram-Positive Isolates from Pus Samples." Ayu 2014; 35(3):330-332.
87. Abduljalil JM, AL-Rakham AA, AL-Haj TM, AL-Rrimy AM, AL-Wheabi AS. "Preliminary Phytochemical Analysis and Antibacterial Activity of Methanol Extracts from *Origanum majorana*, *Rumex nervosus*, and *Withania somnifera*." Int J Pharma Res Health Sci 2018; 6:2844-2850.

88. Alam N, Hossain M, Mottalib Md A, Sulaiman SA, Gan SH, Khalil Md I. "Methanolic Extracts of *Withania somnifera* Leaves, Fruits, and Roots Possess Antioxidant Properties and Antibacterial Activities." *BMC Complement Altern Med* 2012; 12:175.
89. Pandit S, Chang KW, Jeon JG. "Effects of *Withania somnifera* on the Growth and Virulence Properties of *Streptococcus mutans* and *Streptococcus sobrinus* at Sub-MIC Levels." *Anaerobe* 2013; 19:1-8.
90. Furmanova M, Gajdzis KD, Starościak B, Stefan SJ. "In Vitro Cultivation of *Withania somnifera* (L.) Dun. Organs and Their Antibacterial Activity." *Herba Pol* 1999; 44(4):265–269. DOI: [10.31763/hp.4439265](https://doi.org/10.31763/hp.4439265)
91. Ashokkumar K, Murugan M, Dhanya MK, Warkentin TD. "Botany, Traditional Uses, Phytochemistry, and Biological Activities of Cardamom [*Elettaria cardamomum* (L.) Maton] – A Critical Review." *J Ethnopharmacol* 2020; 246:112244. DOI: [10.1016/j.jep.2019.112244](https://doi.org/10.1016/j.jep.2019.112244)
92. Deshpande A, Irani N, Balkrishnan R, Benny IR. "Effects of *Ashwagandha* (*Withania somnifera*) Extract on Sleep Quality in Healthy Adults: A Randomized, Double-Blind, Placebo-Controlled Study." *Sleep Med* 2020; 72:28-36. DOI: [10.1016/j.sleep.2020.03.012](https://doi.org/10.1016/j.sleep.2020.03.012)
93. Jayanthi MK, Prathima C, Huralikuppi JC, Suresha RN, Dhar M. "Anti-Depressant Effects of *Withania somnifera* Extract in Experimental Mice." *Int J Pharm Bio Sci* 2012; 3:33–42.
94. Baek SC, Lee S, Kim S, Jo MS, Yu JS, Ko YJ, et al. "Withaninsams A and B: Phenylpropanoid Esters from the Roots of Indian Ginseng (*Withania somnifera*)." *Plants* 2019; 8(12):527. DOI: [10.3390/plants8120527](https://doi.org/10.3390/plants8120527)
95. Gupta A, Singh S. "Evaluation of Anti-Inflammatory Effects of *Withania somnifera* Root on Collagen-Induced Arthritis in Rats." *Pharm Biol* 2014; 52:308-320. DOI: [10.3109/13880209.2013.835325](https://doi.org/10.3109/13880209.2013.835325)
96. Begum VH, Sadique J. "Long-Term Effect of *Withania somnifera* Herbal Drug on Adjuvant-Induced Arthritis in Rats." *Indian J Exp Biol* 1988; 26:877–882.
97. Al-Hindawi MK, Al-Khafaji SH, Abdul-Nabi MH. "Anti-Granuloma Activity of Iraqi *Withania somnifera*." *J Ethnopharmacol* 1992; 37(2):113–116. DOI: [10.1016/0378-8741\(92\)90069-4](https://doi.org/10.1016/0378-8741(92)90069-4)
98. Shukla KK, Mahdi AA, Mishra V, Rajender S, Sankhwar SN, Patel D, Das M. "*Withania somnifera* Improves Semen Quality by Combating Oxidative Stress and Cell Death and Improving Essential Metal Concentrations." *Reprod Biomed Online* 2011; 22:421–427. DOI: [10.1016/j.rbmo.2011.01.010](https://doi.org/10.1016/j.rbmo.2011.01.010)
99. Sahin K, Orhan C, Akdemir F, Tuzcu M, Gencoglu H, Sahin N, et al. "Comparative Evaluation of Sexual Functions and NF-κB and Nrf2 Pathways of Some Aphrodisiac Herbal Extracts in Male Rat." *BMC Compl Alt Med* 2016; 16:318. DOI: [10.1186/s12906-016-1303-x](https://doi.org/10.1186/s12906-016-1303-x)
100. Kumar A, Kumar R, Rahman MS, Iqbal MA, Anand G, Niraj PK, Ali M. Phytoremedial eect of *Withania somnifera* against arsenic-induced testicular toxicity in Charles foster rats. *Avicenna J Phytomed* 2015; 5:355–364.
101. Bhargavan D, Deepa B, Shetty H, Krishna AP. "Protective Effect of *Withania somnifera* against Oxidative Damage Caused by Ethanol in the Testes of Adult Male Rats." *Int J Basic Clin Pharmacol* 2015; 4:1104-1108.
102. Obidosaka G, Sadowska A, Rumowska M. "Generative Propagation of *Withania somnifera* L." *Herba Pol* 1998; 44:258–264.
103. Said-Al Ahl HAH, Omer EA. "Medicinal and Aromatic Plants Production Under Salt Stress: A Review." *Herba Pol* 2011; 57(1):72–87.
104. Furmanowa M, Gajdzis-Kuls D, Pukacka S, Pukacki P, Zobel A, Malanowski M. "Increase of Withaferin and Glutathione Contents of *Withania somnifera* (L.) Dun. Shoots Cultivated In Vitro on Medium Supplemented with Zn²⁺." *Herba Pol* 2001; 47(4):275–279.
105. A Dar P, R Singh L, A Kamal M, A Dar T. Unique medicinal properties of *Withania somnifera*: Phytochemical constituents and protein component. *Current pharmaceutical design*. 2016 Feb 1;22(5):535-40.
106. Sharifi-Rad J, Quispe C, Ayatollahi SA, Kobarfard F, Staniak M, Stępień A, Czopek K, Sen S, Acharya K, Matthews KR, Sener B. Chemical Composition, Biological Activity, and Health-Promoting Effects of *Withania somnifera* for Pharma-Food Industry Applications. *Journal of Food Quality*. 2021 Dec 29;2021:1-4.