

The effect of mutagenesis on the production of mycelia growth of *Auricularia* species in Nigeria

Abstract

Auricularia species is a jelly-like edible mushroom belonging to the class Basidiomycete, Family Auriculariaceae. It has many medicinal properties and is widely consumed in Nigeria. Not much research effort appears to have been conducted on its domestication and commercial production in Nigeria. This study evaluated the use of sawdust and rice bran as potential substrates for its cultivation. Two strains of *Auricularia* species were collected from the wild and identified as *Auricularia polytricha* and *Auricularia subglabra* by genetic DNA extraction. Pure cultures of the species were screened for strain improvement by mutagenesis using UV light and ethyl methyl sulfonate mutagens. Spawn of the wild species and the mutants were produced using sorghum grain. The produced spawn was used for the production of the mushroom fruit bodies using sawdust supplemented with rice bran. The biological efficiencies of both wild and mutant were 197% and 90% respectively. A combination of sawdust and rice bran supported the growth of both species of *Auricularia*. The Rate of spawn running during spawn production was also negatively affected by mutagenesis in mutants.

KEYWORDS: Mutagenesis, *Auricularia polytrichia*, *Auricularia subglabra*, Genetic DNA, Substrates

Introduction

The strong demand for mushrooms has stimulated the increase in world commercial production. Interestingly Africa contributes only about 1% of total world production of edible mushrooms (Shivute, 2020; Ofodile and Yusuf, 2010). Forests and woodlands that are mushrooms natural habitats have been degraded due to human activity like clearing for agriculture, conversion to other land uses, charcoal burning and logging (Jew, *et al.*, 2016). Edible mushrooms found in many Africa countries have been used in folk medicine (Thomas, *et al.*, 2019). Therefore there is need to protect the mushrooms from being extinct there is need to domesticate as Africans depend on hunting mushrooms in their wild habitat.

The wood ear mushroom (*Auricularia* species) belongs to the Kingdom Fungi and Division Basidiomycota. There are many species of this mushroom and they are widely distributed in the temperate, tropic and semi tropic regions of the world (Kick *et al.*, 2008; Mukerji and Manoharachary, 2010). In nature, *Auricularia* species grow in different habitats such as on dead hardwood trees (*Abies balsamea*) (Conte and Læssøe, 2008), *Sycamore maple* also known as Sycamore (Harding, 2008), on dead or dying branches of trees or decaying log etc. They grow into different sizes, shapes and color in the wet evergreen and forest habitats (Sterry and Barry, 2009; Mohanan, 2011).

Auricularia species has a soft, jelly-like texture, with a mild flavor which makes it edible (Priya, *et al.*, 2016; Conte and Læssøe, 2008). The whole fruit body is edible but not eaten raw and must be washed and cooked (Tomblin, 2007). The fruit body can be smooth, as is typical of younger specimens or undulating with folds and wrinkles and the colour becomes darker with age (Conte and Laessoe, 2008). The inner surface is lighter grey-brown in colour and smooth. It is sometimes wrinkled with folds and may have "veins", making it appear even more ear-like when dry (Sterry and Hughes, 2009; Mohanan, 2011). The internal characteristic has hyphal stratification and abhymenial hair with medulla in where medulla is surrounded by large zona laxa superioris and narrower zonz laxa inferioris (Onyango, *et al.*, 2016). *Auricularia* mushroom is the fourth commercially cultivated in the world after *Agaricus*, *Lentinus*, and *Pleurotus* species (Priya *et al.*, 2016). In Nigeria, commercial production of mushroom has been limited to *Pleurotus* species which are only available to the rich and hotel.

The most common method of cultivation of *Auricularia* species is the log or bag culture methods on hardwood chips and/or sawdust (Mcintosh, 2009; Irawati *et al.*, 2012). The mushroom has been used in the treatment of hemorrhoids, hemoptysis, angina, diarrhea, gastrointestinal upsets, healing of colds and fevers in China (Chen *et al.*, 2008; Harding, 2008). Research reveals that *Auricularia* extract prevents blood clotting (Harding, 2008), stroke, and heart attack (Elkhateeb *et al.*, 2019) and effective in treating diabetes (Lu *et al.*, 2018b).

Mycelia mating, protoplast fusion, induced mutation, cross breeding and molecular genetics transformation has been employed in mushroom breeding (Lee *et al.*, 2011). Molecular breeding by gene transformation has been used to produce new strain with specific property (Lim *et al.*, 2009). Mutagenesis has been reported to improvement yield and nutritional quality in some mushroom species. Improved colour quality and sporelessness have also been reported in mutant strains of some mushroom species (Sharma and Sharma, 2014). Prabu *et al.* (2011) has used protoplast fusion to improve desired mushroom traits. This method has been used to produce mushroom hybrids when conventional methods cannot be achieved. Mushroom produces billions of spores into the air, and this causes health problems like allergies and fever attacks (Chakravarty, 2011).The basidiospore and mycelia of two strains of *Pleurotus florida* and *P. sajor-caju* have been exposed to UV radiation in order to develop low spore strain (Ravishankar *et al.*, 2006; Bahram, *et al.*, 2014). Mutagenesis of two chemical treatment and UV irradiation has been applied for induction of sporelessness in mutant strains of *Coprinus cineris*, *Pleurotus ostreatus* and *Pleurotus pulmonarius* (Sharma and Sharma, 2014).

The goals of mushroom breeding are to produce high yield and good qualities and reduced production cost (Prabu *et al.*, 2011). Liu *et al.* (2011) generated cold tolerant strain of *Volvariella volvacea* by random mutagenesis using alkylating mutagen ethyl methyl sulfonate (EMS). Chemical mutagens are more effective than physical mutagens but chemical mutagens are less suitable due to its disadvantages like uneven penetration into cell wall of mushroom mycelia and target cells, low reproducibility and health risks (Kang *et al.*, 2011).

To the best of my knowledge and up to date, *Auricularia* species production has not been documented in Nigeria. Africa, particularly Nigeria is still far behind in this regard and cultivation of edible mushroom has experienced unprecedented advancement (Osemwegie and Dania, 2016). There is increasing demand for continuous supply of quality mushrooms, *Auricularia* inclusive, hence the need for this research work.

MATERIAL AND METHODS

Strain Collection

Two strains of *Auricularia* species (EW 001 and EW 002) were collected from the wild in the southern part of Nigeria (University of Benin city) and used for this work. The isolates were identified by a Mycologist (Professor Omon Isikhenen) at University of Benin.

Isolation of pure culture:

The collected *Auricularia* species were hydrated, washed several times (six to seven times) with normal saline water, dipped in 70% alcohol for 2 minutes and rinsed with normal saline water. The species were cut into tiny pieces of about 2 mm by 2 mm and inoculated into sterilised potato dextrose agar (PDA). 0.04 g of streptomycin sulphate antibiotic was added to the medium to prevent the growth of bacteria contaminants. Inoculated plates were sealed and incubated at 25°C for mycelia growth. The isolated pure cultures were maintained in PDA slants at 4°C.

Mutagenesis:

Physical mutagen (UV light): The physical mutagen was carried out using Sharma and Sharma, 2014 method:

Actively growing cultures (3 days old) of collected *Auricularia* species (EW 001 and EW 002) on PDA plates were exposed to UV light (244nm, Millipore xx63 70000) for 30, 60,90, and 120 minutes intervals to induce mutation. The generated mutants were subculture on PDA plates, incubated at 25 °C for 7 days. The diameter of mycelia growth was measured. The isolated wild species were used as control.

Chemical mutagen (ethyl methyl sulfonate)

The isolated pure cultures of *Auricularia* species (EW 001 and EW 002) were carried out using Sharma and Sharma, 2014 method by inoculating on sterile Potato Dextrose Agar medium that contained different concentrations of ethyl methyl sulfonate (EMS) ranging from 0.001, 0.002, 0.003, 0.004 and 0.005 % and incubated at 25±2 °C for 7 days to generate mutants. The rate of mycelia ramification in the mutants and the wild strains, determined by the diameter of mycelia growth was monitored.

Cultivation of wild and mutant strains of isolated *Auricularia* species

The substrate for the cultivation was composted as follows; sawdust (79%), calcium carbonate (2%) and rice bran (19%) by Onyango *et al.*, 2011 methods. Sawdust and calcium

carbonate were mixed in the proportion as stated above and the moisture content adjusted to 60-65%. The properly mixed substrates were piled up in cone-shape heap to about 1 meter high for composting. The heap was turned every day to allow even distribution of the generated heat to enable lignin decomposition and the heap left to stand for 30 days. After 30 days of composting, rice bran was added in the required proportion and mixed thoroughly. The compounded substrate was packed inside heat resistant polyethylene bags (1000 g/bag) and sterilized at 121 °C for 30 minutes. The sterilized bags were inoculated with the produced spawn and incubated in the dark at 25±3 °C, until full mycelia colonization was achieved. Rate of mycelia colonization was monitored. The colonized bags were transferred to fruiting house for fruit body growth. The date of initiation of primordia was noted and biological efficiency calculated.

RESULTS AND DISCUSSION

The isolation of *Auricularia* species

The isolated species (EW001, EW002) were identified by Abikoye *et al.* (2016) as *Auricularia polytricha* and *Auricularia subglabra* respectively. Physical mutagenesis (UV light exposure) generated two mutants from each of the identified species; EW1M1 and EW1M2, and EW2M1 and EW2M2 from EW001 and EW002 respectively. Mutagenesis by physical irradiation, retarded the rate of mycelia running (Figure 1a). The mycelia growth of mutant and wild of the *Auricularia* were observed after incubation for five days at 25-26 °C. The *Auricularia* EW001 produced higher mycelia growth than the *Auricularia* EW002 in the physical and chemical mutagens. The four different UV light treatments supported mycelia growth with variations except the 90 min that was not fully ramified. EW 001 (30 min, 60 min and 120 min) showed mycelia growth retarded of 35 mm, 35 mm and 30 mm respectively when compared with the control with the highest mycelia growth of 55 mm. EW 002 (60 min, 90 min and 120 min) showed retarded mycelia growth of 35 mm, 40 mm and 45 mm respectively when compared with the control of 70 mm mycelia growth (Fig 1a).

Chemical mutagenesis using ethyl methyl sulfonate (EMS) generated 5 mutants each; E1M1, E1M2, E1M3, E1M4, E1M5 from EW001 and E2M1, E2M2, E2M3, E2M4, E2M5 from EW002. The EW 001 and EW002 were further used because of the better yield of mycelia and faster rate of their mycelia growth. The 90 min and 120 mins of the EW 001 and EW 002 formed the EW1M1, EW1M2, EW2M1 and EW2M2 mutants that were used in this study. Concentrations with 0.001- 0.004% showed mycelia growth for the two selected species and no growth was observed for EW 001 (0.005%) whereas EW 002 (0.005%) showed slow growth of 40 mm. A retarded mycelia growth of 35 mm, 65 mm and 60 mm were observed in EW 001 (0.003%), EW 005 (0.002% and 0.004%) respectively. Both *Auricularia* EW001 and EW002 were further produced into spawn and then cultivated on sawdust.

Variation of exposure time of the isolates to physical mutagen (UV light) did not affect rate of mycelia ramification of the generated mutants of EW001. However, there was an observed increase in the rate of retardation of mycelia running of mutants of EW002 with increase in exposure time (Figure 1a). Similar result has been recorded by Ravishanker (2006). This

result is also in agreement with Sharma and Sharma (2014) who observed a retardation of mycelia growth of some mushroom species they subjected to physical mutagenesis using UV light.




Figure 1a: The mycelia growth of two *Auricularia* species after exposure to irradiation.

Mutagenesis by chemical mutagen also retarded the rate of mycelia running of the mutants. Similar results have been reported by Ravishanker *et al.* (2006) and Sharma and Sharma (2014) who also used ethyl methyl sulfonate to induced mutation in some species of *Pleurotus*. The retardation of the mycaelia growth rate was more pronounced in isolate EW002 than in EW001 (Figure1b). The retardatory effect of the mutagen generally increased with increase in the concentration of the mutagen (Figure1b).




Figure 1b: The mycelia growth of two *Auricularia* species after different treatment with Ethyl Methyl Sulfonate (EMS).

A combination of sawdust and rice bran supported the growth of both species of *Auricularia* and their mutants. Colonization of the production substrates was faster in the wild strains than the mutants. Full colonization was achieved in 30 days in the wild strains, while it took 50 days for the mutants to fully colonize the substrate. The pin head of the wild strains appeared after the 3rd day of exposure while the mutants pin head appeared on the 10th day. Similar result has been reported by Irawati *et al.* (2012). The sporophores of the wild

strains were bigger and wider than the mutants' strain that appears smaller and never opened (Plates 1a and 1b).

In this report, the *Auricularia* species fruit bodies were very low in both wild and mutant strains when grown on sawdust and rice bran substrate. The biological efficiencies of wild is 197% and mutant species 90%. Though both wild species and mutant strains utilise the nutrients in the substrates, the wild species gene was not altered this makes it easier to utilise the nutrient in the substrate unlike the mutant strain genes that has been altered by mutagens. Biological efficiencies are highly affected by the mutagens. Razaki *et al* (2012) reported that sawdust mixed with oil palm frond and sawdust mixed with empty fruit bunch and spent grain gave the highest yield of sporophore with 288.9% and 260.7% biological efficiencies almost three times the control (sawdust) when *Auricularia polytricha* was grown on these substrates. This is an indication that *Auricularia* species uses the chemical components of the wood especially cellulose and lignin for mycelia growth and produce fruiting body. Onyango *et al* (2011) observed wheat straw and maize cob to be the best substrates for *Auricularia* species with higher yield of mushroom compared to the other substrates they used.



Plate 1a: Matured sporophores from the EW01 grown on sawdust and rice bran.



Plate 1b: Matured sporophores from EW1M1.

Conclusion

A combination of sawdust and rice bran in the right proportion could be a suitable substrate for the cultivation of *Auricularia* species. Mutagenesis, either by physical means (UV light) or chemical means (EMS) negatively affects the mycelia growth of this mushroom and by extension its production. The negative effect of mutagenesis on the mycelia of the *Auricularia* species (mutant strains) may have resulted in aberrant, impaired or loss of function for the gene responsible for yield and the accumulation of the mutation may have lead to the sporophore folding that refused to open.

REFERENCES

- Abikoye, E. T., Oloke, J. K., Dike, E. N., Bora, S. S., Das, S. and Barooah, M. (2016). *Journal of Science and Industrial Research*, **75**(12): 740-746.
- Bahram, B. T; Hamid, R. P; Matin, J. M and Jami, J (2014). Chemical and physical induced mutagenesis in basidiospores of oyster mushroom *Pleurotus ostreatus* var. *florida*. *International Journal of advanced Research*. **2**(11): 915-921.
- Chakraverty, B. (2011). Trends in mushroom cultivation and breeding. *Australian Journal of Agricultural Engineer*. **2**:102-109. -
- Chen G., Luo Y. C., Li B., Guo Y., Li Y., Su W and Xiao Z. L (2008). 'Effect of polysaccharide from *Auricularia auricular* on blood lipid metabolism and lipoprotein lipase activity of ICR mice fed a cholesterol-enriched diet. *Journal of Food Science* **73**(6):103-108.
- Conte, A. D; Læssøe, T (2008). *The Edible Mushroom Book*. Dorling Kindersley. p. 91. [ISBN 978-1-4053-3213-2](https://doi.org/10.1002/9781405332132).
- De Silver, D. D., Raphior, S., Fans, F., Bahkali, A. H (2012a). Medicinal mushrooms in supportive cancer therapies: An approach to anti cancer effects and putative mechanisms of action. *Fungal Diversity* **55**: 1-35.
- Elhateeb, W. A, Daba, G. M, Thomas, P.W, Wen, T.C (2019). Medicinal mushrooms as a new source of natural therapeutic bioactive compound. *Egyptian Pharmaceutical Journal*. **8**(2):88-101
- Harding, P (2008). *Mushroom Miscellany*. Harper Collins Publisher. London 2008p
- Irawati, D. C., Hayashi, Y., Takashima, S., Wedatama, F., Ishiguri, K., Lizuka, N., Yoshizawa and Yokoto, S (2012). "Cultivation of the edible mushroom *Auricularia polytricha* using sawdust based substrate made of three Indonesian commercial plantation species, *Falcataria moluccana*, *Shorea* spp and *Tectona grandis*." *Micologia Aplicada International*. **24**(2): 33- 41.
- Kang, S. Y., Ha, B. K. and Kim, D. S. (2011). Mutation breeding pp150-161. *In: Radiation Technology and its Applications* (Yoo BD ed.), Nuclear Training and Education Center, Korea.
- Kirk P. M; Cannon P. F; Minter D. W; Stalpers J. A. (eds) 2008 – *Dictionary of the Fungi*, 10th edition. CABI Publishing, UK.
- Jew, E. K. K, Dougill, A. J, Sallu, S. M, O'Connell, J, Benton, T. G (2016). Miombo woodland under threat: consequences for tree diversity and carbon storage. *Ecological Management*. **361**:144-153.

- Lee, J., Kang, H., Kim, S., Chang-Yun, L. And Hyeon-Su, R. (2011). Breeding new strains of mushroom by basidiospore chemical mutagenesis. *Mycobiology*. **39(4)**: 272-277.
- Lim, H. and Choi, H. T (2009). Enhanced expression of Chitinase during the autolysis of mushroom in *Coprinellus congregates*. *Journal of Microbiology*. **47**:225-228.
- Liu, N, Chen, M, Song, J, Zhag, Y, Gong, P, Chen, X (2022). Effects of *Auricularia auricular* polysaccharide on gut Microbiota composition in Type 2 Diabetic Mice. *Molecules*. **27**(180): 6061.
- Liu, Z., Zhang, K., Lin, J. F. And Gao, L. Q. (2011). Breeding cold tolerance strain by chemical mutagenesis in *Volvariella volvacea*. *Science Horticulture*. **130**:18-24.
- Lu, A., Meng, S., Zhiyu, F., Yaoyan, X., Mengen, Y., Shoang, W., Yongiun, Z and Wermin, W (2018b). Antidiabetic effects of the *Auricularia auricular* polysaccharides simulated hydrolysates in experimental Type-2 Diabetic Rats. *Natural Produced Communication*. **13**(2): 195-200.
- Ma, Y., Wang, C., Zhang, Q., Peng, X (2018). The effect of polysaccharides from *Auricularia auricular* (Huaier) in adjuvant anti gastrointestinal cancer therapy: A systematic review and network meta analysis. *Pharmacological Research* **132**: 80-89.
- Mohanan, C. (2011). *Macrofungi of Kerala*. Kerala Forest Research Institute. pp. 90–3. [ISBN 81-85041-73-3](#).
- Mukerji and Manoharachary (2010). Taxonomy and ecology of Indian fungi.
- Ofodile, L .N and Yusuf, I .O. (2010) Mushrooms and their development in Africa. Agriculture: Africa's "engine for growth" - Plant science and biotechnology hold the key. *Aspects of Applied Biology* **96**: 1-6
- Onyango, B. O, Mbaluto, C. M, Mutuku, C. S, Otieno, D. O (2016). Molecular characterization of wood ear mushrooms (*Auricularia* sp) from Kakmega Forest in Western Kenya. *Current Research in Environment and Applied Mycology* **6**(1):51-60.
- Onyango, B. O, Palapala, P.F, Arama, S. O, Wagai, Gichimu, B. M (2011). Suitability of selected supplemented substrates for cultivation of Kenya native Wood Ear Mushroom (*Auricularia auricu*). *American Journal of Food Technology*. <http://dx.doi.org/10.4236/ajps.2012.312202>.
- Osemwegie, O. O and Dania, A. T (2016). Production of Mushrooms as Complimentary Agroincome in Nigeria. *Annals of Science and Technology*. **1**(1): 19-27.
- Prabu, C. S. and Lee, Y. K. (2011). Mutation breeding of mushroom by radiation. *Journal of Radiation Industry*. **5**(4):285-295.

- Priya, R. U, Geetha, D, Darshan, S (2016). Biology and cultivation of Black Ear Mushroom - *Auricularia*. *Advances in Life Sciences*. 5(22).
- Razak, D. L; Noorlidah, A; Norjuliza, M. K. J; Vikineswary, S (2012). Comparative study of mycelia growth and sporophore yield of *Auricularia polytricha* (Mont) Sacc on selected palm oil wastes as fruiting substrate. *Applied Microbiological of Biotechnology* **97**:3207-3213.
- Ravishanker, S; Pandey, M; Tewari, R. P; Krishna, V (2006). "Development of sporeless/low sporing strains of *Pleurotus* through mutation." *World Journal of Microbiologic Biotechnology* **22**:1021-1025.
- Sharma, R and Sharma, B. M (2014). "Strain improvement in *Pleurotus ostraetus* using UV light and Ethyl methyl sulfonate as mutagens." *Academic Journals*. **8**(5): 432-436.
- Shivute, F. N. (2020). Cultivation of edible mushrooms in Namibia: Prospects and challenges of small scale farmers. *African Journal of Agricultural Research*. **16**(11):1582-1586.
- Sterry, P; Barry, H (2009). *Complete Guide to British Mushrooms and Toadstools*. HarperCollins. p. 290. [ISBN 978-0-00-723224-6](#).
- Thomas, P. W, Elkahateeb, W. A, Daba, G, (2019). Truffle and Truffle-like fungi from continental Africa. *Acta Mycologica*. 54(2): 1132.
- Tomblin, Gill (2007). *How to Identify Edible Mushrooms*. Harper Collins UK. p. 146. [ISBN 978-0007259618](#)