

Evaluation of soil fertility status of research farm, College of Agriculture and Research Station, Katghora, Korba, Chhattisgarh

Formatted: English (United States)

ABSTRACT

The present investigation was conducted to evaluate the soil fertility status of soils in the research farm, College of Agriculture, and Research Station Katghora, Korba, (C.G.) for soil available macro and micronutrients status. A total of 102 soil surface samples (0-15) were collected from each field and analysed with respect to concerning pH, EC, OC, macronutrients (N, P, K, S) and micronutrients (Fe, Mn, Cu, Zn, and B) by standard procedures. Results were classified under various fertility ratings classes of available nutrients. The fertility status of the study area was reported on the basis of based on nutrient index value. The results showed that soil pH varied from 4.31 to 5.42 with a mean of 4.79 and indicated that the soils were found to be moderately acidic to strongly acidic in soil reaction. The electrical conductivity of soil water suspensions ranged from 0.04 to 0.11 with a mean 0.07 indicated that soils of study area were suitable for all crops. The OC varied from 2.8 to 6.4, with mean value 4.3 g kg⁻¹ and found that about 80 percent soils are under low organic carbon content. Most of the soil samples were found low in available N content, it ranged from 150.53 to 263.16, with mean 224.28 kg ha⁻¹. The available phosphorus varied from 20.51 to 93.32 with mean 49.66 kg ha⁻¹. The available potassium varied from 103.86 to 233.55 with mean 148.81 kg ha⁻¹. The available sulphur varied from 5.99 to 51.55 with mean 22.25 kg ha⁻¹. Micronutrients Fe, Mn, Cu and Zn varied from 6.01 to 44.40 mg kg⁻¹ with mean 20.87 mg kg⁻¹, 4.10 to 16.95 with mean 10.82 mg kg⁻¹, 0.38 to 3.58 with mean 1.82 mg kg⁻¹ and 0.48 to 1.29 with mean 0.82 mg kg⁻¹, respectively. The available B varied from 0.14 to 0.91 with mean 0.48 mg kg⁻¹. All the soil samples were found to be high with respect to Fe, Cu and Mn whereas Zn and B were found deficient as 12.75 and 41.19 percent, respectively. According to NIV, the soils of study area were classified in low fertility class for nitrogen, Potassium, Boron, and sulphur, medium fertility class for phosphorus and zinc, and high fertility class for Fe, Mn, Cu.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Keywords: Physico-chemical properties, available nutrients, Nutrient index Value

INTRODUCTION

Soil fertility refers to the inherent capacity of the soil to provide macro and micronutrients in the soil. The physical, chemical and biological tests provide information about the capacity of soil to supply mineral nutrients (Ganorkar and Chinchmalatpure, 2013). Soil productivity is the capacity of the soil to produce crops with a specific system of management and is expressed in terms of yields. Both soil fertility and productivity are the key pillars for food production and soil quality is of equal significance in the background of soil degradation caused by many factors. Balanced fertilizer use refers to the application of essential plant nutrients in optimum quantities and in the right proportional through appropriate method and right time of application suited for a specific crop and agronomic situation. Soil test-based fertility management is an effective tool for increasing the productivity of agricultural soils that have high degree of spatial variability resulting from the combined effects of physical, chemical, or biological processes

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

(Goovaerts,1998). ~~Nutrient~~The nutrient index value method and fertility indicators can be used to evaluate the fertility status of the soil. The soil fertility status under different cropping ~~sequence~~sequences can also be assessed by using nutrient index approach (Singh *et al.*, 2016).

Formatted: English (United States)

Formatted: English (United States)

Material and Methods

The study was conducted on “Evaluation of soil fertility status of research farm, College of Agriculture, And Research Station Katghora, Korba (C.G.)”. The conducted study involved the collection and chemical analysis of 102 soil samples ~~of from~~ college farm. The analysis of collected soil samples was carried out in the laboratory of Department of Soil Science and Agricultural Chemistry, College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur. The study area was located in 22°35'N longitude 82°75' E latitude. The soils of the study area, in general red yellow colour with light texture was found to be sandy loam. These soils were found to be acidic in nature soil reaction. Kharif is the main cropping season of study area, the total geographical area paddy is the main crop followed by Maize and pigeon pea pulses crops. Wheat, Maize and linseed crops are also grown in Rabi season. The surface soil samples (0-15 cm) were collected from each field of the study area. The collected soil samples were air dried and soil samples were crushed by wooden hammer and sieved through a 2 mm sieve. Soil pH was determined by glass electrode pH (Piper 1967), EC with Solubridge method (Black 1965), OC by wet digestion method (Walkley and Black’s rapid titration method 1934) Available potassium was extracted by shaking with neutral normal ammonium acetate for 5 minutes (Hanway and Heidal 1952), DTPA- extractable micro-nutrients with 0.005 N Di-ethylene Triamine Penta Acetic acid (DTPA), 0.01M calcium chloride dehydrates and 0.1M Tri ethanol amine buffered at adjusted pH 7.3 using an atomic absorption spectrophotometer Lindsay and Norvell (1978)and Available Boron through ammonium acetate and EDTA used as buffer masking solution and azomethine-H (Berger and Trong 1939).

Formatted: English (United States)

The Nutrient Index values and Fertility Classes

According to Ramamoorthy and Bajaj (1969), the nutrient index values (NIV) for various soil parameters were determined from the amount or proportion of samples with low, medium, or high usable nutrient status and classified into different fertility groups.

$$NIV = \frac{(1 \times PL) + (2 \times PM) + (3 \times PH)}{100}$$

Where,

NIV = nutrient index value

PL= % samples fall under low category.

PM= % sample es fall under medium category.

PH = % sample as fall under high category.

Table 1 Nutrient Index Values for Nutrients

NIVfortheNutrient	Fertilityclass(basedonNIV)		
	Low	Medium	High

Macronutrients(N,P,K andS)and Micronutrients(Fe,Mn,Cu,ZnandB)	<1.33	1.33-2.33	>2.33
--	-------	-----------	-------

Formatted: English (United States)

Result and Discussion

Soil pH

A study on the soil reaction (pH) of the study area revealed that the soils were moderately acidic to strongly acidic and the pH varied from 4.31 to 5.42 with a mean value of 4.79 ± 0.25 . Out of the total soil samples (102 samples), 82.35% of the soils were found moderately acidic and 17.64 % as strongly acidic in soil reaction. The soil pH shows are acidic which may be due to [the](#) leaching loss of basic cations from the soil surface because of high rainfall, parent material i.e., igneous rocks are found in the study area they produce acidity in soil after decomposition by weathering and vegetation also leads to acidic conditions. Similar results were reported by Chandrakaret *et al.*, (2017) while working on the soils pH status of Janjgir-Champa they found that pH of the research area ranged from 4.73 to 5.98 with an average value of 5.83 ± 0.38 .

Formatted: English (United States)

EC (dSm^{-1})

The electrical conductivity of the soil water suspension ranged from 0.04 to 0.11 dSm^{-1} with a mean value of $0.07 \pm 0.01 \text{ dSm}^{-1}$. All the soil samples fall under the normal range ($< 1.0 \text{ dSm}^{-1}$). Table 3 indicated that the soils of [the](#) study area safe for growing ~~of~~ all types of crops. The very less EC value of area may be ascribed to extensive leaching of all soluble salt to deeper layer and runoff losses from surface soil layer. The results were supported by the research work done previously by Meher *et al.*, (2020), in the soils of Pahanda Durg district in Chhattisgarh state. He concluded that the EC ranged between $0.01\text{-}0.32 \text{ dSm}^{-1}$ with an average of 0.11 dSm^{-1} . All samples were under good class i.e., $< 1.0 \text{ dSm}^{-1}$.

Formatted: English (United States)

Organic Carbon (%)

The variations in the soil organic carbon content were 2.8 to 6.4 g kg^{-1} with an average of $0.43 \pm 0.98 \text{ g kg}^{-1}$. Out of all the soil samples, majority of the soil samples i.e., 79.41% were found to be in low in OC and 20.58% samples in medium organic carbon status. The soil C content was found to be low in the study area which may be due to high temperature that enhanced oxidation of native organic matter and carbon ~~and also~~ low rates of organic matter additions in soils. The obtained results were in line with the results obtained by research work of Meher *et al.* (2020) reported that the OC ranged from $2.7 \text{ to } 7.1 \text{ g kg}^{-1}$ with a mean of 4.8 g kg^{-1} of KVK farm Pahanda in Durg District, Chhattisgarh. Similar results were also supported by Singh *et al.*, (2018), Kumar *et al.*, (2017), Dadsena *et al.* (2021).

Formatted: English (United States)

Available Nitrogen (kg ha^{-1})

The available Nitrogen content in the soil samples ranged from 150.53 to $263.16 \text{ kg ha}^{-1}$ with a mean value of $224.28 \pm 32.70 \text{ kg ha}^{-1}$. All the soil samples were found in low nitrogen status. Asija (1956) is 70-76 % only i.e., it recovers only oxidizable N fraction of dry soil N pool. It can be noted that the whole area seems N deficient which might be due to the fact that these soils were very poor organic C content which is biggest source of N. It can also be due to the extensive leaching and runoff losses of various form of N a tropical environment is its high

temperature which leads to rapid loss of soil organic matter due to volatilization. The similar results were supported by Singh *et al.* (2017) investigated the soils of Baloda-Bazar District from 55 different villages and found that soil available nitrogen content ranged from 165.3 to 263.25 kg ha⁻¹ in different soil orders, with a mean value of 197.3 kg ha⁻¹, falling it's in the low fertility index group. The similar results were also supported by Tirkey *et al.*, (2017).

Available Phosphorus (kg ha⁻¹)

The status of the available phosphorus varied from 20.51 to 93.32 kg ha⁻¹ with an average value of 49.66±15.80 kg ha⁻¹. Majority of the soil samples i.e., 78.43 % was found to be in medium and 8.82 % in high categories, whereas only 12.74 % in the low categories. The deficient P status can be attributed to low organic carbon content and as maximum phosphorus is fixed kaolinite clay mineral and Al and Fe oxide present in the acidic soil of study area. So, its availability is low to medium. The similar results were reported by Tigga *et al.*, (2022), reported that available P content in KVK farm varied from 36.45 to 63.91 kg ha⁻¹, with a mean value of 52.32 kg ha⁻¹. Patil *et al.*, (2016) in the soils of the Dindur sub-watershed of Karnataka where it was observed that the status of P was low to medium in range. The results were also supported by Singh *et al.*, (2018). Jatav (2010) also conclude that soil P deficiency may also be due to improper fertilization in [the](#) past, porous texture, soil reaction or other nutrient interaction and fixation.

Formatted: English (United States)

Available potassium (kg ha⁻¹)

The available potassium of the study area varied from 103.86 to 223.55 kg ha⁻¹ with a mean value of 148.81±32.26 kg ha⁻¹. It was found that 39.21 % and 60.78% of samples were under low and medium category, respectively. It was found that Inceptisol of study area had low to medium the status with respect to available potassium. That may be attributed to presence of kaolinite and micaceous clay minerals that inherently have low to medium K in their crystal lattices. The similar results were supported by Dadsena *et al.*, (2021) assessed the soil fertility status of Bamhanidih village, Janjgir–Champa district of Chhattisgarh. They reported that the available K ranges from 202.5-293.7 kg ha⁻¹, with a mean value of 255.3 kg ha⁻¹. The similar result showed that the Palani *et al.* (2021).

Available Sulphur (kg ha⁻¹)

The sulphur status varied from 5.99 to 51.55 kg ha⁻¹ with a mean value of 22.25±10.58 kg ha⁻¹. It was found that sulphur had 59.80%, 28.43 % and 11.76 % of samples were found in low, medium and high rating, respectively. The study area had S status within low to medium ranged which may be due to the poor organic C reserve in soil along with the leaching and runoff losses of sulphate ions from surface layer of study area. The findings were supported by Mehar *et al.* (2020) who assessed the fertility status of the KVK farm in Pahanda. The similar results were also supported by Dadsena *et al.*, (2021). Goswami *et al.* (2014).

Available Manganese (mg kg⁻¹)

The status of the available Mn ranged from 4.10 to 16.95 mg kg⁻¹ with a mean value of 10.82±2.71 mg ha⁻¹. Almost soil samples were found to be in high level of Mn were collected soil samples from different villages of Majhwa Block of Mirzapur district. The results showed that the DTPA extractable Mn 4.6 to 16.2 with a mean value 5.13 mg kg⁻¹. The Mn bearing minerals in the parent material might be the reason for higher Mn content in the soils and due to better

Formatted: English (United States)

supply of Mn to rice in flooded soil as Mn is soluble in relatively acidic and reduced soil condition. The similar result showed the Mehar *et al.*, (2019), (Mandal and Haldar, 1980). Singh *et al.*, (2017)

Available Iron (mg kg⁻¹)

The status of the available Fe varied from 6.01 to 44.40 mg kg⁻¹ with a mean value of 20.87 ± 10.10 mg ha⁻¹. Most of the soil samples were found to be in high levels of Fe. During study area in the research farm, high available Fe content in Katghora, Korba, might be due to its topography and cultivation of rice, induced by prolonged submergence coupled with reducing conditions. This order of the soils in study area is not deficient in Fe as the amount of Fe required by crops is being released by Fe bearing minerals in these soils. The results were supported by Rajeswar *et al.* (2009) Krishna *et al.* (2018). Manda *et al.*, (2018), Dadsena *et al.*, (2021).

Formatted: English (United States)

Available Copper (mg kg⁻¹)

The status of the available Cu in the study area ranged from 0.38 to 3.58 mg kg⁻¹ with a mean value of 1.82 ± 0.99 mg ha⁻¹. All the samples were found to be high in Cu. Evaluated soils of the Baloda Bazar district Chhattisgarh. The soil analysis showed that the available Cu content ranged from 0.20 to 3.00 mg kg⁻¹ with an average value of 1.81 mg kg⁻¹. The similar result showed the Singh *et al.* (2018), Dadsena *et al.* (2021) and Palani *et al.* (2021).

Formatted: English (United States)

Available Zinc (mg kg⁻¹)

The status of the available Zn varied from 0.48 to 1.29 mg kg⁻¹ with a mean value of 0.82 ± 0.25 mg ha⁻¹. The results were classified into 3 different rating categories as 12.75, 74.50 and 12.75 % samples were under deficient, sufficient and high categories for available Zn respectively. Zn was related to the important soil characteristics. The Zn deficiency increased with increase in pH and decrease with increase in organic C. Here it can be noted that the dominant portion of the area seems Zn sufficient which might be due to the low soil pH which renders Zn in soil solution and makes it available for crops. The similar result showed the Similar observation was made by Rai *et al.* (1970), Mehar *et al.*, (2019), Barooah *et al.* (2020), Mandal (2018).

Formatted: English (United States)

Available Boron (mg kg⁻¹)

The status of the available B varied from 0.14 to 0.91 mg kg⁻¹ with a mean value of 0.48 ± 0.17 mg ha⁻¹ (Table 3). The percentage distribution of the soil samples was found to be sufficient (59.80 %) and deficient (40.20%) categories. This may be attributed to continuous cereal-cereal cropping without supporting B fertilizers which resulted in mining of from soil reserve. These results were in line with the finding of Kumar (2014), Mehar *et al.*, (2019).

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Table 2: Salient properties of soil in the study area of katghora korba

S. No.	Parameters	Range	Average	SD ±
1	Soil pH	4.31-5.42	4.79	0.25
2	EC (dS/m)	0.04-0.11	0.07	0.01

regarding fertilization and other management practices (Olivares, 2016; Araya-Alman et al. 2020).

Formatted: Font: Not Bold

The study's focus on the research farm in Korba, Chhattisgarh, acknowledges the importance of region-specific research. Soil characteristics and fertility can vary widely based on factors such as climate (Parra et al. 2012; Olivares and Hernandez, 2019), vegetation (Lopez et al. 2019; Lopez and Olivares, 2019), and geological conditions (Hernandez and Olivares, 2019; Hernandez et al. 2020). A study in this specific region helps provide targeted recommendations for local farmers (Hernandez et al. 2018a).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

~~Sustainable Agriculture:~~ Analyzing soil quality is a fundamental aspect of promoting sustainable agricultural practices (Hernandez et al. 2018b; Lobo et al. 2023). This study can shed light on whether the current farming methods in the Korba region deplete soil nutrients or maintaining a balanced nutrient cycle (Vega et al. 2022). Comparing these findings with soil studies in Latin American tropical areas can provide valuable insights into best practices for sustaining soil health (Rey et al. 2022; Rodriguez-Yzquierdo et al. 2023).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

The comparison between soil quality in Chhattisgarh and Latin American tropical areas offers a unique opportunity to identify commonalities and differences in soil fertility management across different parts of the world (Zingaretti et al. 2016; Olivares, 2023). This comparison can lead to the exchange of knowledge and ideas, potentially resulting in innovative strategies to improve soil quality globally (Olivares et al. 2020; Pitti et al. 2021).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Collaborative research that spans continents enhances the global understanding of soil science and agricultural practices (Montenegro et al. 2021). Such collaborations foster knowledge sharing and the adoption of successful strategies from one region to another (Zingaretti and Olivares, 2018).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Soil quality is closely linked to environmental health. By examining soil fertility status and comparing it with studies from Latin America (Hernandez and Olivares, 2020), the researchers can assess the potential impact of agricultural practices on soil degradation (Olivares et al. 2018), nutrient loss (Paredes et al. 2021), and erosion in both regions (Olivares et al. 2015). This information can contribute to more informed land management decisions.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

In conclusion, the study on evaluating soil fertility status in Korba, Chhattisgarh, and its comparison with studies from tropical areas in Latin America holds immense relevance for promoting sustainable agriculture, global knowledge exchange, environmental protection, and informed policy-making. It contributes to the broader understanding of soil science and its practical applications in diverse agricultural landscapes.

CONCLUSION

Formatted: English (United States)

The soils of research farm were moderately to strongly acidic with normal electrical conductivity and organic carbon status was found in lower categories. The mean values of macronutrients were found as 224.28, 49.66, 148.80 and 22.25 kg ha⁻¹ soil available N, P, K and S, respectively. The average values of micronutrients were observed as 20.87, 10.82, 1.82, 0.82 and 0.48 mg kg⁻¹. As per NIV criteria, the available nitrogen, potassium and sulphur were observed in low fertility whereas available phosphorus classified as medium fertility class. In case of micronutrients, available Fe, Mn and Cu were found in high, Zn in medium and B in low fertility classes.

The soils of study area were classified as moderately to strongly acidic which indicated that lime application is beneficial to crop production. Soil N status was evaluated as low status, 25% extra of General recommended dose (GRD) can be suggested for nitrogenous fertilizer. 60% area was low for sulphur status hence, GRD dose of S in the form of either S powder or bentonite S, should be applied and also recommended use of single super phosphate in place of DAP or complex fertilizers. In case of micronutrients, only boron status was recorded as low category hence borax powder at the rate of 2.5 kg ha⁻¹ can be applied. ZnSO₄·2H₂O should be applied in case of Zn level below critical point @ 25 kg ha⁻¹.

REFERENCES

Formatted: Spanish (Spain, International)

[Araya-Alman, M., Olivares, BO., Acevedo-Opazo, C. et al. Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in Venezuela. J Soil Sci Plant Nutr. 2020, 20 \(3\): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>](#)

Formatted: Font: (Default) Times New Roman, 12 pt, Spanish (Spain, International Sort)

Chandrakar, K., Srivastava, L.K., Raghav, R.S. and Mahant, A.D. 2017. Estimation of available nitrogen, organic carbon, pH and correlation study of the sampled soil of Inceptisols under Janjgir district of Chhattisgarh. A Journal of Multidisciplinary Advance Research, 6(1):80-82.

[Dadsena, R., Senger, A., Singh, C., Pradhan, A.K., Rao, I.P., Sahu, K.K., Porte, S.S., Khalkho, D. and Sahu, R.K., 2021. Soil fertility status of available nitrogen in soil through soil fertility mapping using GPS and GIS techniques of Bamhanidih village and block, Janjgir-Champa district of Chhattisgarh state.](#)

Formatted: English (United States)

Ganorkar and Chinchmalpure. (2013). Physiochemical assessment of soil in Rajura Bazar in Amravati district of Maharashtra (India). International Journal of Chemical, Environment and Pharmaceutical Research, 4(2&3):46-49.

[Goswami, R.G., Mishra, V.N., Singh, A.K. and Kurre, P.K. 2014. Sulphur Status and their spatial Variability in soil of four blocks of Raipur District. An Asian Journal of Soil Science, 9\(2\):330-333.](#)

Formatted: English (United States)

Goswami L., Sarkar S., Mukherjee S., Das S., Barman S., Raul P., Bhattacharyya P., Mandal

NC, Bhattacharya S., Bhattacharya SS., Vermicomposting of tea factory coal ash: metal accumulation and metallothionein response in *Eisenia fetida* (Savigny) and *Lampitoma auritii* (Kinberg). *Bioresource technology*, 1;166: 96-102.

Gomez, K.A., and Gomez, A. A. 1984. Duncan's multiple Range test Statistical procedures for Agricultural Research. 2: 540-544.

Formatted: English (United States)

Hanway, J.J., Heidal H. Soil analysis methods as used in Iowa State College Soil Testing Laboratory. Iowa State College of Agriculture Bulletin. 1952; 57:1-31.

[Hernandez, R., Olivares, B., Arias, A, Molina, JC., Pereira, Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia* 2020. 38\(2\):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>](http://dx.doi.org/10.4067/S0718-34292020000200095)

[Hernández, R; Olivares, B. Arias, A; Molina, JC., Pereira, Y. Agroclimatic zoning of corn crop for sustainable agricultural production in Carabobo, Venezuela. *Revista Universitaria de Geografía*. 2018a, 27 \(2\): 139-159. <https://n9.cl/12m83>](https://n9.cl/12m83)

[Hernández, R; Olivares, B., Arias, A; Molina, JC., Pereira, Y. Identification of potential agroclimatic zones for the production of onion \(*Allium cepa* L.\) in Carabobo, Venezuela. *Journal of the Selva Andina Biosphere*. 2018b, 6 \(2\): 70-82. \[http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf\]\(http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf\)](http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf)

Formatted: Font: Italic

[Hernández, R. Olivares, B., Ecoterritorial sectorization for the sustainable agricultural production of potato \(*Solanum tuberosum* L.\) in Carabobo, Venezuela. *Agricultural Science and Technology*. 2019. 20\(2\): 339-354. \[https://doi.org/10.21930/rcta.vol20_num2_art:1462\]\(https://doi.org/10.21930/rcta.vol20_num2_art:1462\)](https://doi.org/10.21930/rcta.vol20_num2_art:1462)

Formatted: Font: Italic

[Hernández, R., Olivares, B. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. *Tropical and Subtropical Agroecosystems*, 2020. 23\(2\):1-12. <https://n9.cl/zeedh>](https://n9.cl/zeedh)

Formatted: English (United States)

Jatav, G. K. Evaluation of Soil Fertility Status in Inceptisol of Baloda Block in Janjgir District of Chhattisgarh. M.Sc. Thesis, I.G.K.V., Raipur, 2010, 67-78.

Jackson, M. L. Soil chemical analysis, Prentice Hall of India Pvt. Ltd. N. Delhi, 1967, 205.

Lindsay, WL, Norvell WA. Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Sci. Soc of Amer. J.* 1978; 42:421-428.

[Lobo, D; Olivares, B; Rey, J.C; Vega, A; Rueda-Calderón, A. Relationships between the Visual Evaluation of Soil Structure \(VESS\) and soil properties in agriculture: A meta-analysis. *Scientia agropecuaria*, 2023. 14 \(1\): 67– 78. <http://dx.doi.org/10.17268/sci.agropecu.2023.007>](http://dx.doi.org/10.17268/sci.agropecu.2023.007)

Formatted: Spanish (Spain, International Sort)

[López-Beltrán, M., Olivares, B., Lobo-Luján, D. Cambios de usos de suelo y vegetación en la comunidad agraria Kashaama, Anzoátegui, Venezuela: 2001-2013. *Revista Geográfica De América Central*. 2019, 2\(63\):269-291. <https://doi.org/10.15359/rgac.63-2.10>](https://doi.org/10.15359/rgac.63-2.10)

Formatted: Spanish (Spain, International Sort)

[López, M. Olivares, B., Normalized Difference Vegetation Index \(NDVI\) applied to the agricultural indigenous territory of Kashaama, Venezuela. *UNED Research Journal*. 2019, 11\(2\): 112-121. <https://doi.org/10.22458/urj.v11i2.2299>](https://doi.org/10.22458/urj.v11i2.2299)

Formatted: English (United States)

Mandal, S. Characterization and Delineation of Farming Situations of Chhattisgarh Plains, M. Tech. (Agril. Engg.) Thesis, I.G.K.V. Raipur (Chhattisgarh), 2018, 34-50. 10.

Meher, S.K., Singh, V., Bala, J., Samuel, S., Motghare, R. and Tedia, K. 2020. Assessment of spatial variability of soil fertility status in KVK farm of Pahanda in Durg district, Chhattisgarh using GIS-GPS. International Journal of Chemical Studies, 8(2): 304-309.

[Montenegro, E., Pitti-Rodríguez, J., Olivares-Campos, B. Identification of the main subsistence crops of Teribe: a case study based on multivariate techniques. Idesia \(Arica\). 2021. 39\(3\), 83-94. <https://dx.doi.org/10.4067/S0718-34292021000300083>](#)

[Olivares, B. Description of soil management in agricultural production systems of sector hammock in Anzoátegui, Venezuela. La Granja: Revista de Ciencias de la Vida. 2016, 23\(1\): 14-24. <https://doi.org/10.17163/lgr.n23.2016.02>](#)

[Olivares, B., Hernández, R. Regional analysis of homogeneous precipitation areas in Carabobo, Venezuela. Revista Lasallista de Investigación. 2019, 16\(2\):90-105. <https://doi.org/10.22507/rli.v16n2a9>](#)

[Olivares, B., Pitti, J., Montenegro, E. Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. Revista Brasileira de Gestao e Desenvolvimento Regional. 2020. 16\(3\):59-71. <https://doi.org/10.54399/rbgdr.v16i3.5871>](#)

[Olivares Campos, B. O. Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health. Springer Nature. 2023. <https://doi.org/10.1007/978-3-031-34475-6>](#)

[Olivares, B., Hernández, R; Coelho, R., Molina, J.C., Pereira, Y. Analysis of climate types: Main strategies for sustainable decisions in agricultural areas of Carabobo, Venezuela. Scientia Agropecuaria. 2018. 9\(3\): 359 – 369. \[10.17268/sci.agropecu.2018.03.07\]\(https://doi.org/10.17268/sci.agropecu.2018.03.07\)](#)

[Olivares, B., Lobo, D. y Verbist, K. Application of the USLE model in erosion plots under soil and water conservation practices in San Pedro de Melipilla, Chile. Revista Ciencia e Ingeniería. 2015. 36 \(1\):3-10.](#)

[Paredes, F., Olivares, B., Rey, J., Lobo, D., Galvis-Causil, S. The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. SAINS TANAH - Journal of Soil Science and Agroclimatology, 2021. 18\(1\), 58-64. <http://dx.doi.org/10.20961/stjssa.v18i1.50379>](#)

[Parra, R., Olivares, B., Cortez, A., Rodríguez, M.F. Pluviometric homogeneity patterns in climatic stations of Anzoátegui state, Venezuela. Revista Multiciencias. 2012. 12 \(Extraordinario\): 11-17. <https://n9.cl/xbslq>](#)

[Pitti, J; Olivares, B; Montenegro, E. The role of agriculture in the Changuinola District: a case of applied economics in Panama. Tropical and Subtropical Agroecosystems. 2021. 25 – 1: 1 - 11. <http://dx.doi.org/10.56369/tsaes.3815>](#)

[Piper, CS. Soil and plant analysis. Asian publishing House, Bombay, N. Delhi, 1967, 85-102.](#)

[Ramamoorthy, B.& Bajaj, J.C. 1969. Available nitrogen, phosphorus and potassium status of Indian soils. Fertilizer News 14\(8\): 25-36.](#)

Formatted: Spanish (Spain, International)

Field Code Changed

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, Spanish (Spain, International Sort)

Formatted: English (United States)

Formatted: English (United States)

Field Code Changed

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, English (United States)

Formatted: Portuguese (Brazil)

Field Code Changed

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, Portuguese (Brazil)

Formatted: English (United States)

Field Code Changed

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Field Code Changed

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

[Rey, J.C.; Olivares, B.O.; Perichi, G.; Lobo, D. Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. Sustainability, 2022, 14, 13531. <https://doi.org/10.3390/su142013531>](#)

[Rodríguez-Yzquierdo, G.; Olivares, B.O.; Silva-Escobar, O.; González-Ulloa, A.; Soto-Suarez, M.; Betancourt-Vásquez, M. Mapping of the Susceptibility of Colombian Musaceae Lands to a Deadly Disease: Fusarium oxysporum f. sp. cubense Tropical Race 4. Horticulturae 2023, 9, 757. <https://doi.org/10.3390/horticulturae9070757>](#)

[Singh, G., Sharma, M., Manan, J., Singh, G. \(2016\). Assessment of Soil Fertility Status under Different Cropping Sequences in District Kapurthala. J Krishi Vigyan. 5\(1\): 1-9](#)

[Singh, C., Bajpai, R.K., Tiwari, A., Chandra, M., and Krishna, B. 2018. Evaluations of soil fertility status of available major nutrients \(N, P & K\) and micronutrients \(Fe, Mn, Cu & Zn\) in Vertisols of Balodabazar block in Balodabazar district of Chhattisgarh. J Pharmacogn Phytochem, 2, pp.10-12.](#)

[Singh, S. N., Latore A.M. and Singh. S.K., Soil Fertility Status of Majhwa Block of Mirzapur District of Eastern UP, India. Int. J. Curr. Microbiol. App. Sci \(2017\) 6\(9\): 2019-2026.](#)

[Subbiah, V., Asija GL. A. rapid procedure for estimation of available nitrogen in soil Current Science. 1956; 25:259- 260.](#)

[Tirkey, D.E., Thomas. T. Assessment of Soil Sample by Analysing Chemical Properties of Soil in Korba District of Chhattisgarh, India. Research Journal of Chemical and Environmental Sciences Res J. Chem. Environ. Sci. Vol 5 \[4\] August 2017: 76-81.](#)

[Vega A, Olivares B, Calderón MAR, Rey JC, Lobo D, Gómez JA, Landa BB. Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. Plants, 2022, 11\(15\):2070. <https://doi.org/10.3390/plants11152070>](#)

[Walkley, A., Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil science. 1934 Jan 1;37\(1\):29-38.](#)

[Zingaretti, M.L., Olivares, B. Analysis of the meteorological drought in four agricultural locations of Venezuela by the combination of multivariate methods. UNED Research Journal. 2018. 10 \(1\):181-192. <http://dx.doi.org/10.22458/urj.v10i1.2026>](#)

[Zingaretti, M.L., Olivares, B., Demey Zambrano, J.A., Demey, J.R. Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. Revista FAVE - Ciencias Agrarias 2016. 15 \(2\): 39-50. <https://doi.org/10.14409/fa.v15i2.6587>](#)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International

Formatted: No underline, Font color: Auto, Spanish (Spain, International Sort)

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)