

Review Article

A REVIEW ON: COMPREHENSIVE CONSEQUENCES OF GALL STONES AS CONCOMITANT DISEASE IN HYPOTHYROIDISM (CCCCH)

ABSTRACT

This review investigates the potential correlation between hypothyroidism and gallstone disease, with a particular focus on sub clinical hypothyroidism. The study explores the physiological factors that may contribute to gall stone formation in hypothyroid patients, including changes in bile flow and sphincter of Oddi dysfunction. The review also highlights the role of thyroid hormone in lipid metabolism and their potential impact on the composition of bile. The findings suggest a significant association between gallstone disorders and thyroid malfunction, particularly subclinical hypothyroidism. The review concludes that early identification and treatment of subclinical hypothyroidism could significantly reduce biliary stone disease patients. regular monitoring of thyroid levels in hypothyroid individuals is recommended to prevent complications due to abnormal thyroid hormone levels.

KEYWORDS:

Hypothyroidism, Gallstone Disease, Subclinical Hypothyroidism, Thyroid Hormone Level, Biliary Stone Disease

INTRODUCTION

Thyroid hormones have profound variation during the life span and are associated with severe adverse health impacts ^(1,2). The most prevalent biliary

disease is gallstones, which affect 10-15% of adults in the United States (20 million people). Cholecystectomy was performed on 3% of these adults, and 85% of them remain asymptomatic. Females are affected more frequently than men by a ratio of 3:1. ^(3,4) Several variables, such as age, gender, and ethnic origin, are connected to the occurrence of gallstones (33%) in the population. Gallstones can form as a result of specific medical conditions. Gallstones are more likely to form in people who are overweight, pregnant, have certain dietary restrictions, have Crohn's disease, have had stomach surgery, had their terminal ileum removed, have genetic spherocytosis, have sickle cell disease, or have thalassemia ⁽⁵⁾. There has been debate about whether thyroid conditions may lead to gallstone disease for many years. There are several theories, in particular, suggesting a potential connection between gallstone disease and hypothyroidism. These factors include the established connection between thyroid dysfunction and lipid metabolic issues, which may afterwards result in a change in the bile's composition. ⁽⁶⁾ Subsequent investigations ⁽⁷⁾ also showed that hypothyroid patients had decreased biliary flow. Thyroid hormone receptors are expressed in the sphincter of Oddi, and thyroxine directly relaxes the sphincter. ⁽⁸⁾ Gallstone development is thought to be influenced by fundamental physiological factors such as poor bile flow and sphincter of Oddi dysfunction ⁽⁹⁾. The delayed biliary tract emptying caused by the reduced pro-relaxing effect of T4 on the sphincter of Oddi (SO) in hypothyroidism suggested by experimental studies, along with a potential cholesterol load in the bile and a decreased hepatocytic excretion rate, may provide an important explanation for the rising prevalence of common bile duct (CBD) stones ⁽¹⁰⁾. This review depicts the correlation between hypothyroid and gallstone

METHOD

The purpose of this review was to determine whether thyroid function and gallstone disease are related in any way. Choledocholithiasis instances with documentation served as the subject of this investigation. The information was gathered using the terms cholelithiasis, thyroid profile and hypothyroidism connection, common bile duct stone, risk factors, and sphincter of Oddi dysfunction to search via multiple databases like google scholar, pubmed, and medline. All of the publications we found were cross-sectional, case-control, and prospective studies. ⁽¹⁵⁾

Individuals were classified as normal if their serum TSH levels were between 0.5 and 4.7mIU/L. Subclinical hypothyroidism is defined as a serum TSH level of 4.7–10mIU/L with normal T3, T4 levels. TSH concentrations more than 10mIU/L are regarded as clinical hypothyroidism. ⁽¹⁸⁾

RESULTS

In conclusion, a number of recent research have found a link between CBD stones and hypothyroidism, or subclinical hypothyroidism. It is possible that changes in the function of the SO, rather than changes in the metabolism of cholesterol or bile excretion rate, are what are responsible for the association between CBD stones and hypothyroidism, as evidenced by the higher prevalence of hypothyroidism in patients with CBD stones compared to patients with gallbladder stones. It is yet unknown if hypothyroid people who have had their gallbladder removed are more likely to get CBD stones than euthyroid people in the same circumstances.

In many respects, it is probable that the absence of thyroxine in hypothyroidism leads to a decrease in bile flow. The lack of the pro relaxant action of thyroxine on the SO appears to be a major element contributing to the lower bile flow in

hypothyroidism, in addition to the increased cholesterol burden in bile and the decreased bile secretion rate. ⁽¹⁷⁾

- Research by Sun et al. in China, Figueredo et al. in California, and Sharma et al. in India all showed that cholelithiasis affects females more frequently than men, which is consistent with the majority of patients in our study being female. ⁽²¹⁾
- 53.7% of the hypothyroid participants in this research had subclinical hypothyroidism. This is consistent with the Laukkarinen et al. research, which found that individuals with cholelithiasis had a considerably high frequency of subclinical hypothyroidism. ^(16,17)
- When compared to the general population, individuals with undiagnosed or subclinical hypothyroidism were 12 times more likely to develop choledocholithiasis, according to case-control research by BJ et al. to determine the relationship between hypothyroidism and gallstone disease. ⁽²⁰⁾
- Prospective cross-sectional research by Ghimire et al. in West Nepal revealed that 43 out of 160 individuals with gallstone disease (26.7%) had hypothyroidism, with 21.1% of these patients having subclinical hypothyroidism and the other 5.6% having clinical hypothyroidism. ⁽²⁴⁾
- Inkinen et al. came to the conclusion that there is a link between CBD stone sufferers and hypothyroidism that has been addressed in their study. According to the study's findings, primary hypothyroidism affected 8% of the study group compared to 1% of the control group, with a P value of less than 0.01. ⁽¹⁰⁾
- A study done in Germany by Völzke et al. revealed a link between gallstone disease and thyroid function based on gender. Comparatively to

women, men were discovered to have an independent risk factor for developing cholelithiasis called thyroid dysfunction. However, in our study, the prevalence of hypothyroidism was higher in girls than in males, perhaps as a result of the lesser size of the sample size that we recruited than in the aforementioned German study. ⁽²³⁾

- According to research by Milionis et al., 70.9% of hypothyroid patients had a BMI of more than 25 kg/m² and were obese [18]. This matches our study's findings that the majority of hypothyroid individuals (73.7%) were fat. [22]

DISCUSSION

Table 1. Data of the studies published reporting an association between hypothyroidism and gall stones.

S N O	AUTHORS	JOURNAL	STUDY TYPE	patient population	gender	age (years)	RESULTS	CONCLUSION
1	Inkinen et al	Hepato gastroenterol	Retrospective	168	65% F	> 60 years	In Group I the prevalence of previously diagnosed hypothyroidism was 7/86 (8%) compared with 1/86 (1%) in Group II (P = 0.01). Hypothyroidism was previously diagnosed only in the common bile duct stone patients (Group I) of over 60 years of age, where the prevalence was 7/66 (11%).	There is a significant association between the common bile duct stones and previously diagnosed hypothyroidism. Their stronger association between the common bile duct stones and hypothyroidism compared to gallbladder stones and hypothyroidism suggests a mechanism other than merely the cholesterol metabolism mediated mechanism. The 11% prevalence of previously diagnosed hypothyroidism in the common bile duct stone

								patients of over 60 years of age suggests all patients with common bile duct stones be screened for current thyroid dysfunction.
2	Dr. Sharath et al.	International Journal of Surgery Science	A case control study	140	Both (male and female)	>18	A case control study was done where 140 cases and 140 controls were sampled. A significant proportion of the cases were found to have hypothyroidism in comparison to controls with odds ratio of 12.21. Other variables included female sex, age greater than 60 years, non-vegetarian diet, increased LDL, VLDL and Triglyceride levels.	This study proves that hypothyroidism is a significant independent risk factor for choledocholithiasis, and this can give validation to inculcate change in the approach and management of a case of choledocholithiasis
3	Dr. pradeep ghirmire et al.	journal of dental and medical science	prospective study	160	Both (male and female)	range of 16 - 83 years	This study shows 75% female and 25% male, age wise minimum age was 16 and maximum age was 83. The mean age was 42.68 years. All 160 patients had gallstone in ultrasound of the abdomen without complication. The prevalence of hypothyroidism in gallstone patient in our study was 26.7%	Higher association of hypothyroidism was seen (26.7%) in gallstone patients
4	Hui Sun, Hong Tang et al	world journal of gastroenterology	observational study	3573	Both (male and female)	range of 18 - 70 years	A total of 3573 people were included, 10.7% (384/3573) of whom had gallstone diseases. Multiple logistic regression analysis indicated that the incidence of	We assume that age and sex are profoundly associated with the incidence of gallstone disease; the metabolic risk factors for gallstone disease were different between men and women

							gallstone disease in subjects aged 40-64 or ≥ 65 years was significantly different from that in those aged 18-39 years ($P < 0.05$); the incidence was higher in women than in men ($P < 0.05$). In men, a high level of fasting plasma glucose was obvious in gallstone disease ($P < 0.05$), and in women, hypertriglyceridemia or obesity were significant in gallstone disease ($P < 0.05$).	
5	Anastasios Milionis et al	international scholarly research notes	correlation study	736	616 females and 118 males	range of 54 - 73 years	The correlation between BMI and the various thyroid hormones differed in both sexes. In women, there was a statistically significant association between BMI and TT3 and TT4, while in men BMI was positively associated with FT3 levels and negatively with TSH. BMI depends on fluctuations in body weight and height.	In conclusion, we have shown that variations of normal thyroid function are accompanied by differences in BMI perhaps due to the changes in the resting energy consumption. The high incidence of the pathological disorders in thyroid function combined with the strong influence of various environmental factors (diet, exercise, etc.) can increase weight with an unknown biological mechanism and lead to obesity. Further studies are required for a general assumption of the existence or nonexistence of a correlation between obesity and variations of normal thyroid function

CONCLUSION:

According to the findings of this study, gallstone disorders and thyroid malfunction, particularly subclinical hypothyroidism, are related. Biliary stone disease is linked to an increased frequency of subclinical hypothyroidism. Subclinical hypothyroidism can greatly reduce biliary stone disease patients if it is identified and treated in a timely manner. In order to prevent complications brought on by an abnormally high rise in thyroid hormone, we suggest that hypothyroid individuals should monitor their thyroid levels every three to six months.

REFERENCES

1. Ramezani Tehrani F, Aghaee M, Asefzadeh S. The comparison of thyroid function tests in cord blood following cesarean section or vaginal delivery. *Int J Endocrinol Metab.* 2003; 1:22–26.
2. Zadeh-Vakili A, Ramezani Tehrani F, Hashemi S, Amouzegar A, Azizi F. Relationship between Sex Hormone Binding Globulin, Thyroid Stimulating Hormone, Prolactin and Serum Androgens with Metabolic Syndrome Parameters in Iranian Women of Reproductive Age. *Diabetes Metabolism.* 2012:S:2.
3. I.J. Beckingham, B.J. Rowlands. Post cholecystectomy problems, H. Blumgart (Ed.), *Surgery of Liver and Biliary Tract* (third ed.), WB Saunders, London (2000), p. 186 PMID: PMC1119388
4. A. Nakeeb, A.G. Comuzzie, L. Martin, *et al.* Gallstones: genetics versus environment *Ann. Surg.*, 235 (2002), p. 842
5. Al-Jiffry B, Shaffer E, Saccone G. Changes in gallbladder motility and gallstone formation following laparoscopic gastric banding for morbid obesity. *Can J Gastroenterol* 2003; 17:169.
6. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med.* 2000; **160**:526–534.
7. Laukkarinen J, Sand J, Saaristo R, Salmi J, Turjanmaa V, Vehkalahti P, Nordback I. Is bile flow reduced in patients with hypothyroidism? *Surgery.* 2003; **133**:288–293.
8. Inkinen J, Sand J, Arvola P, Pörsti I, Nordback I. Direct effect of thyroxine on pig sphincter of Oddi contractility. *DigDisSci.* 2001; **46**:182–186.
9. Cicala M, Habib FI, Fiocca F, Pallotta N, Corazziari E. Increased sphincter of Oddi basal pressure in patients affected by gallstone disease: a role for biliary stasis and colicky pain? *Gut.* 2001; **48**:414–417.

10. Inkinen J, Sand J, Nordback I: Association between common bile duct stones and treated hypothyroidism. *Hepatogastroenterology*. 2000, 47:919-21.

11 Yawar Zahoor Watali, Rahul Jain, Rajandeep Singh Bali, Ankul Mittal; Is hypothyroidism a risk for gallstone disease? a study to assess the association; *International Surgery Journal* 4 (8); Vol 4; 2665-2669

12 Henry Völzke, Daniel M Robinson, and Ulrich John; Association between thyroid function and gallstone disease; *World Journal of Gastroenterology*; 2005; 11(35); 5530-5534

13 Hassan H. Zaini, Kussay M. Zwain; Prevalence of hypothyroidism in patients with gallstone disease; *AL-Qadisiyah Medical Journal*; Vol 6 No.10; 108-117

14 Rana ranjit ,Arun gupta, sukrit shah, avreen singh , Amrit pal singh bhatia;prevalence of hypothyroidism in patients with biliary stone: A prospective study *Int surg J* 2016,vol3 ,2022_2024 .

15 Kharishma P.nair ,siddharth gowthaman ;thyroid function and gallstone disease ;is there an association ? *international surgery journal int surgJ*. 2021 may;89(5);1670-1673.

16. Johanna Laukkarinen, Gediminas Kiudelis, Marko Lempinen, Sari Ratu, Hanna Pelli, Juhani Sand, Esko Kemppainen, Caj Haglund, and Isto Nordback Increased Prevalence of Subclinical Hypothyroidism in Common Bile Duct Stone Patients *The Journal of Clinical Endocrinology & Metabolism* 92(11):4260 – 4264

17 Johanna Laukkarinen, Juhani Sand, and Isto Nordback *The Underlying Mechanisms: How Hypothyroidism Affects the Formation of Common Bile Duct Stones—A Review* Hindawi Publishing Corporation *HPB Surgery* Volume 2012, Article ID 102825, 7 pages

18. Mir Mujtaba Ahmad , Mir Irfan Nazir , Haneef Mohamed Dar1, Ubaid Ali , Moonis Mirza , Khalid Kawoosa ,Evaluation of thyroid profile in biliary tract stones *International Surgery Journal* Ahmad MM et al. *Int Surg J*. 2015 Aug;2(3):344-347.

19. Suaad L. Ibrahim *The Impact Of Thyroid Dysfunction And TSH On The Pathogenesis Of Gallstone And Its Composition* *Journal of Kufa for Nursing Science* Vol. (4) No.(1) 2014

20 Bj SC, Kb N, Goutham M. Hypothyroidism as a risk factor for choledocholithiasis: A case-control study in an Indian population. *Int J Surg Sci.*

2019;3(4):439-44.

21.Sun H, Tang H, Jiang S, Zeng L, Chen E-Q, Zhou T-Y, . Gender and metabolic differences of gallstone diseases. *World J Gastroenterol WJG.* 2009; 15(15):1886–91.

22 .Milionis A, Milionis C. Correlation between Body Mass Index and Thyroid Function in Euthyroid Individuals in Greece. Vol. 2013, *ISRN Biomarkers.* Hindawi; 2013. pe651494.

23 Völzke H, Robinson DM, John U. Association between thyroid function and gallstone disease. *World J Gastroenterol WJG.* 2005; 11(35):5530–4.

24 Pradeep Ghimire MS, Frcs ED. Association of Gallstone Disease with Hypothyroidism in Western Region of Nepal. *IOSR J Dental and Med Sci.* 2017;16(8):40-2.

UNDL