

# **Crop Health Monitoring Through Remote Sensing: A Review**

---

## **ABSTRACT**

Agriculture is basis of livelihood for a major portion of world population. It provides food to humans. With the increasing population and climate change there is need to enhance production to fulfil the demand of growing population. Remote sensing technology has potential to predict nutrient requirement by providing various information related to plant and soil in quantitative terms thereby increasing productivity. It plays important role in monitoring crop health, crop growth and development, nutrient management, pest and disease management, water management, etc. Evaluation of crop canopy provide various information regarding agronomic parameters. The data provides a better alternative for natural management than traditional methods and this kind of management enhances efficiency of various resources by avoiding their overuse. This paper reviews remote sensing technology for crop health monitoring, highlighting its importance with new ideas for agriculture.

*Keywords: Remote sensing, Drought indices, Crop health monitoring, Nutrient management, Water management, Weed management.*

## **1. INTRODUCTION**

Remote sensing is a technology that enables to acquire information about any object without coming into direct contact with the object under study. Its application is transforming management and planning in the field of agriculture. With the introduction of new technologies, it seems a very bright future to move farm management to a next level of production and productivity. Agriculture production can be enhanced by proper use of major agricultural resources like water, fertilizer and pesticides, as when and where required. Fertilizer is one of the major resources that can affect crop health, thereby reducing yield and quality. Improper use of fertilizer leads to imbalance between organic content and nutrients in the crop. Hence, it is important to precisely apply nutrient to enhance the fertilizer use efficiency and reduce the loss from the farming system. This can be achieved by monitoring crop growth and development. Monitoring agricultural crop production during the growing season and estimating the potential crop yields are both important for the assessment of seasonal production [1]. Unfavourable climatic condition and growing conditions may result in variation of crop productivity. Dong et al., [2] observed effect of climatic variability on cropland productivity variation based on remote sensing observations. Remote sensing helps to timely monitor these conditions and provide a model of crop with precision. The monitoring of crop health follows seasonal patterns in relation to biological life cycle of the crop. This technology enables to collect data timely with precision without destructive sampling of the crop. Remote sensing uses various sensors for recording data.

These sensors sense the electromagnetic radiations. These sensors are available in two forms: passive sensors and active sensors. Passive sensors record the radiations that is reflected by the object or emitted from the earth surface. On other hand, active sensors (e.g., LIDAR, RADAR) are those which emits their own electromagnetic radiation [3]. In agricultural remote sensing, most of the sensors are designed to record a specific portion of the electromagnetic radiation. Interaction between the electromagnetic spectrum with any of the material can be used for the qualitative and quantitative analyses of various materials [4]. Remote sensing using space-borne sensors are tools for taking repetitive and synoptic observations. These data can be used for various assessment and management of field [3]. This technology has importance for the efficient utilization of available resources and to take appropriate measures to lower the appropriate loss caused by the climatic variation or any other stress factor. This paper reviews remote sensing and its application for crop health monitoring, its utility and future perspective in the field of agriculture.

### 1. Crop health monitoring

Remote sensing is a tool for monitoring crop growth and development, nutrient deficiency, diseases, moisture stress, etc. It can be used to track the growth of crop at different time intervals. Timely information about crop can help to identify the problem by various vegetative factors. It is important to understand crop production response to agronomic management and environmental stress [5]. Vegetative indices like Normalised difference vegetative index (NDVI) measures the greenness over the time. It has been observed that NDVI is highly correlated with crop growth and health, and can be used for monitoring crop condition [6]. The crop condition can be monitored by (i) instantaneous monitoring method in which NDVI values of cropland is compared with those of the same period in the previous year and (ii) the crop growth process monitoring method that forms the crop growing profiles with time series NDVI images and assesses the crop condition by comparing inter-annual crop growth profile [7]. Leaf area index (LAI) and crop biomass are two crucial pointers of crop health and development [8]. Remote sensing data on these pointers can assist to obtain significant information on site specific properties (e.g., soils, topography), management (e.g., water, nutrients and other inputs), and various stress factors (e.g., diseases, weeds, water, and nutrient stress) [9]. The data obtained used to estimate LAI and biomass for various crops. Several studies showed that LAI and biomass were highly correlated with several Optical Spectral Vegetation Index (OSVIs) and Radar Polarimetric Parameters (RPPs) [10], the LAI can be assessed from both hyperspectral and the 3D canopy models [11], the red edge position (REP) extracted from ground hyperspectral reflectance can accurately estimate the kinnow mandarin LAI and Chlorophyll content and this can be effectively used to assess crop health status in a wide range for real-time nutrition management in the orchard [12]. Crop health monitoring and quantifying crop stress due to biotic and abiotic stress can be done by using various indices for mapping and monitoring drought. Certain factors like evapotranspiration, soil moisture and vegetation conditions can be used to assess and monitor drought characteristics [13]. The indices for soil moisture status in rooting zone are given in table 1.

**Table 1. Various indices for soil moisture status in rooting zone**

S. no.	Drought indices	Reference
1	PDSI (Palmer Drought Severity Index)	[14,15]
2	Drought Severity Index (DSI)	[16]
3	Evapotranspiration Deficit Index (ETDI)	[17]
4	Standardised Precipitation and Evaporation Index (SPEI)	[18]

From remote sensing, several drought indices can be obtained viz., Normalised Difference Water Index [19], Crop Water Stress Index [20], Water Deficit Index [21]. It can provide a

good estimate of evaporative fraction, the ratio of ET and available radiant energy, with the AVHRR and MODIS data [22].

**Table 2. Some vegetative indices used in agriculture:**

Index	Application	References
Normalised difference vegetative (NDVI)	Biomass, breeding, phenotyping, yield, disease, nitrogen management, soil moisture, water stress.	[23,25,26,27,28,29]
Green NDVI (GNDVI)	Water stress, biomass, diseases	[30,8,31,32,33,34]
Red edge normalised difference vegetation index (RENDVI)	Yield irrigation management, N-status/application, diseases	[35,36,37,38]
Soil adjusted vegetation index (SAVI)	Yield, biomass, diseases, N-concentration and uptake, water stress	[39,8,33,40,41,42]
Ratio vegetative index	Crop yield biomass	[43,8]
Normalised pigment chlorophyll ratio index (NCPI)	Water stress	[44]
Chlorophyll absorption ratio index (CARI)	Chlorophyll content	[45]
Chlorophyll vegetation index (CVI)	Crop yield, biomass, N-uptake, soil moisture, water stress	[46,47,36]
Water balance index	Irrigation scheduling	[48]
Normalised difference water content (NDWI)	Vegetation water content	[49]
Normalised water index (NWI)	Soil moisture and crop yield	[28]

These indices can be used to assess the crop health and stress condition; hence these data can be used to analyse the quality of the crop.

## 2. Nutrient management

For proper growth and development of crop, sufficient amount of nutrients is required at right stage. Nutrient deficiency like in case of nitrogen, it reduces leaf chlorophyll content that results in low light absorption. Nutrient requirement of crop plant can be estimated by studying leaf optic properties such as fluorescence, reflectance and transmittance. Chlorophyll fluorescence gives quick and precise information related to stress based on the fluorescence emission pattern of leaves, tissues and even the whole plant [50]. This emission is captured when part of light energy absorbed by chlorophyll for photosynthesis is re-emitted when excited with UV- a near 340- 360 nm or blue-green light [51]. The fluorescence emission at different level of plant stress was successfully detected and imaged on different crops for deficiency of nitrogen and zinc on maize (*Zea mays*), as well as heat and water stress on *Zalea (Rhododendron sp.)* [52]. It was observed found that green chlorophyll index based on NIR (800 nm) and green (550 nm) wavelength were strongly related to Chlorophyll Concentration Index (CCI) as a measure of chlorophyll content [53]. Plant dry matter accumulation and grain yield were observed to be pointedly influenced by the absorption of Photosynthetically Active Radiation. It was positively related to yield production at tillering and panicle initiation stage [54]. Thermography is also used for nutrient deficiency. It can visualize stomatal movement without presence of an illuminizing source [55]. A high temperature for under fertilized barley (*Hordeum vulgare L.*) than well

fertilized barley with nitrogen as reference nutrient was observed [56]. Reflectance in the red and near infrared region of the electromagnetic spectrum for estimating the nitrogen requirement of the crop using early season estimates of nitrogen uptake and potential yield have been developed [57]. Sivagnanam et al., [58] found a positive correlation of NDVI inflection point with Nitrogen content were found positive at maximum tillering stage followed by flowering stage, milky stage and least in tillering initiation. The results provide nitrogen estimation through hyperspectral instrument easily with less time consuming. The NDVI increase with increasing leaf greenness and green leaf area, and can be used as a guide for in season nitrogen application [57]. Xu et al. [59] suggested that soil moisture, vegetation and soil crusts can contribute to the conservation of soil total nitrogen.

### **3. Water management**

Application of remote sensing for water management helps for precise application of irrigation water, estimation of soil moisture availability, water requirement at different growth stages and in mitigating water stress and hence, achieving optimum crop growth and yield. Remote sensing data can help to detect variation in the field and to apply variable irrigation with commonly used irrigation systems. This can help to overcome water stress resulting from extremely wet and dry condition, also to get maximum uniform yield in all the parts of the field while reducing water and nutrient losses [60,61]. Spectral reflectance in the visible region was observed higher in water stressed condition than the non-stressed condition. Vegetation indices like NDVI, RVI (Ratio vegetation index), PVI (Perpendicular Vegetative Index) and GI (Green Index) were found lower for stressed and higher for non-stressed crop [62]. Das *et al.* [63] developed a high-resolution soil moisture soil temperature service that can be used for real time decision support system in precision agriculture. Thus, this technology plays major role in efficient use of water and it can be further enhanced by the development of hyperspectral sensors.

### **4. Weed management**

Remote sensing is an efficient way for mapping weed patches in crop for site specific weed management [64]. Remote sensing with precision agriculture helps for better weed management practices [62]. Spectral signature helps to identify and differentiate between weed and crop plant related to their phenological and morphological attributes that are different from the crop [65]. Kaur *et al.*, [66] observed that radiance ratio and NDVI values were maximum in solid stand and minimum in solid weed plots. It was also found that pure stand can be easily distinguished from pure weed stand of *R. spinosus* after 30 DAS. Kaur and Jaidka [67] noted that pure wheat can be distinguished from pure population of *Malvaneglecta* after 30 DAS and remain distinguished upto 120 DAS and different levels of weed population can be differentiated amongst themselves after 60 DAS. Remote sensing technology thus can be used to identify weeds of different species and their infestation in field crops. Weed prescription maps can be prepared with Geographical Information system (GIS), on the basis of which farmers can be recommended for preventive control measures for weed control.

### **5. Disease and pest infestation**

Remote sensing is a tool for disease and pest identification, spectral reflectance for chlorosis, yellowing of leaves and foliage reduction can be used for making correlation and interpretation [68]. Hyperspectral remote sensing technology is helpful in rapid assessment of physico-chemical response of crops to biotic stress like disease infestation without destructive sampling [69]. Riedell et al. [70] found remote sensing as an effective and inexpensive method for pest and disease affected plant in oat and concluded that the difference can be evaluated by canopy characteristics and spectral references. William et al. [71] worked on different types of vegetation indices on Landsat imagery before and after defoliation for differentiating between healthy and unhealthy vegetation cover. It monitors the disease efficiently in the early stages of disease development, when it is difficult to

distinguish the symptoms with field monitoring. Various techniques using RGB, multi-spectral, hyperspectral, thermal, and fluorescence imaging have been used to identify diseases in a wide range of crops [72]. Specific disease indices will amend disease detection, identification and monitoring in precision agriculture applications [73]. Spectral disease indices have possibility to improve disease detection, identification and monitoring in precision agriculture applications [74]. Diseases can cause considerable loss of crop production and thus their detection at the beginning and its spatial extent can help to contain the disease spread and lower production losses [65].

## 6. Conclusion

On the basis of findings of different research workers, it can be concluded that remote sensing can improve crop assessment and crop monitoring which can help in crop growth, site specific nutrient management, water management, weed management and monitoring pest and disease. This technology collects real time data with accurate position that leads to an effective analysis of data. Thus, we can identify different problems and solutions. Farmers can easily apply the solutions for the identified problems. Further, there is need to develop more accurate and new methodologies to crop health monitoring, disease detection under diverse climatic condition and field condition. Hence, remote sensing plays an important role in agriculture specially in the improvement of economy through crop health monitoring.

## REFERENCES

1. Doraswamy PC, Moulin S, Cook PW, and Stern A. 2003. Crop Yield Assessment from Remote Sensing. *PE & RS*. 69: 665-674.
2. Dong T, Liu J, Shang J, Quain B, Huffman T, Zhang Y, Champagne C and Daneshfar. 2016. Productivity in the Canadian Prairies Using Time Series MODIS FAPAR. *Remote Sens*. 2016. 8(4).281
3. Navalgund RR, Jayaraman and Roy PS. 2007. Remote Sensing Applications: An Overview. *Current Science*. 93: 12.
4. Omania E, Bae E, Park E, Kim MS, Baek I Kabenge I, Cho BK. 2023. Remote sensing in field crop Monitoring: A comprehensive review of sensor Systems, Data Analyses and Resent Advances. *Remote Sens*. 15, 354.<http://doi.org/10.3390/rs15020254>.
5. Peng Y, Dai YL, Fang S, Gong Y, Wu X, Zhu R, Liu K.2019. Remote prediction of yield based on LAI estimation in oilseed rape under different planting methods and nitrogen fertilizer applications. *Agric. For. Meteorol*. 271 :116- 125.
6. Badhwar GD. 1980. Crop emergence date determination from spectral data. *Photogram. Eng. And Remote Sensing*. 46: 369-377.
7. Meng JH and Wu BF. 2008. Study on the crop condition monitoring methods with remote sensing. *Intl. Archives Photogram, Remote sensing, and Spatial Info. Sci. (Beijing, China)* 37 (part B8): 945-950.
8. Zhou L, Chen N, Chen Z, Xing C. 2016. ROSSC: An efficient remote sensing observation-sharing method based on cloud computing for soil moisture mapping in precision agriculture. *IEEE J. Sel. Top Appl. Earth Obs. Remote Sensing*. 5588-5598.
9. Compos I, Gonzalez-Gomez, L Villodre J, Calera M, Campoy J, Jimenez N, Plaza C, Sanchez-Prieto S, Carlera A. 2019. Mapping within-field variability in wheat yield and biomass using remote sensing vegetation indices. *Precis. Agric*. 20: 214-236.

10. Jin X, Yang G, Xu X, Yang H, Feng H, Li Z, Shen J, Lan Y, Zhao. 2015. Combined multi-temporal optical and radar parameters for estimating LAI and biomass in winter wheat using HJ and RADARSAR-2 data. *Remote sens.* 7: 13251-13272.
11. Kalisperakis I, StentoumisCh, Grammatikopoulos, Karantzas. 2015. Leaf area index estimation in vineyard from UAV hyperspectral data, 2 D image Mosaics and 3 D canopy surface models. *The International Archives of the photogrammetry, Remote Sensing and Spatial Information Sciences*, Vol. XL-1/W4,2015. International Conference on Unmanned Aerial Vehicles in Geomatics, 30 Aug-20 Sep 2015, Toronto, Canada.
12. Ali A, Imran MM, 2020. Evaluating the potential of red edge position (REP) of hyperspectral remote sensing data for real time estimation LAI & chlorophyll content of Kinnow Mandarin (*Citrus reticulata*) fruit orchard. *Sci. Hortic. Amst.* 2020. 267: 109326.
13. Su B, Dong X. 2017. Drought monitoring and assessment using Remote sensing. *Remote Sensing of Hydrological Extremes*, Springer Remote Sensing/Photogrammetry, DOI 10.101007/978-3-319-43744-6\_8.
14. Palmer WC. 1965. Meteorological drought, Research paper 45. US Department of Commerce, Weather Bureau, Washington, DC.
15. Palmer WC, 1968. Keeping track of crop moisture conditions, nationwide: the new crop moisture index. *Weatherwise* 21(4), 156-161.
16. Su Z, Yacob A, He Y, Boogard H, Wen J, Gao B, Roerink G, Van Diepen K. 2003. Assessing relative soil moisture with remote sensing data: theory and experimental validation. *Phys Chem Earth* 28(1-3)89-101.
17. Narasimhan B, Srinivasan R. 2005. Development and evaluation of soil moisture deficit index (SMDI) and Evapotranspiration Deficit Index (ETDI) for agricultural drought monitoring. *Agricultural and Forest Meteorology.* 133(2005) 69-88.
18. Vitente-Serrano SM, Beguera S, Lopez-Moreno JI. 2010. A multiscale drought index sensitive to global warming: The standardized precipitation evaporation index. *J Climate.* 23: 1696-1718.
19. Rouse JW, Haas RH, Schell JA and Deering DW. 1973. Monitoring vegetation systems in the great plains with ERTS. *Third ERTS symposium, NASA,* 351(1): 309-317.
20. Jackson RD, Idso SB, Reginato RJ and Pinter PJ 1981. Canopy temperature as a crop water stress indicator. *Water Resources Research* 17: 1133-1138.
21. Moran MS, Clarke TR, Inoue Y. and Vidal A. 1994. Estimating crop water deficit using the relation between surface-air temperature and vegetation index. *Remote sens Environ.*, 61(3): 319-246.
22. Batra N, Islam S Venturi V, Bisht G and Jiang L. 2006. Estimation and comparison of evapotranspiration from MODIS and AVHRR sensors for clear sky days over the southern Great Plains. *Remote Sens Environ.* 103: 1-15.
23. Schaefer MT, Lamb DW. 2016. A combination of plant NDVI LiDAR measurements improve the estimation of pasture biomass in tall fescue (*Festuca arundinacea* var. Fletcher). *Remote Sens.* 8: 109.
24. Duan T, Chapman SC, Guo Y, Zheng B. 2017. Dynamic monitoring of NDVI in wheat agronomy and breeding trials using an unmanned aerial vehicle. *Field crops Res.* 210: 71-80.
25. Hassan MA, Yang M, Rasheed A, Yang G, Renolds M, Xia X, Xiao Y, He Z. 2019. A rapid monitoring of NDVI across the wheat growth cycle for grain yield prediction using a multispectral UAV platform. *Plant Sci.* 282: 95-103.

26. Yuan L, Pu R, Zhang J, Yang H. 2016. Using high spatial resolution satellite imagery for mapping powdery mildew at a regional scale. *Precis. Agric.* 17: 332-348.
27. Amaral LR, Molin JP, Portz G, Finazzi FB, Cortinov L. 2015. Comparison of crop canopy reflectance sensors used to identify sugarcane biomass and nitrogen status. *Precis Agric.* 16:15-28.
28. Ihuoma SO, Madramootoo CA. 2019. Sensitivity of spectral vegetation indices for monitoring water stress in tomato plants. *Comput. Electron. Agric.* 163: 104860.
29. Ballester C, Zarco-Tejada PJ, Nicolas E, Alarcon JJ, Fereres E, Intrigliolo DS, Gonzalez-Dugo, VJPA. 2018. Evaluating the performance of xanthophyll, chlorophyll and structure-sensitive spectral indices to detect water stress in five fruit tree species. *Precis. Agric.* 19: 178-193.
30. Zhou J, Khot LR, Boydston RA, Miklas PN, Porter L. 2018. Low altitude remote sensing Technologies for crop stress monitoring: A case study on spatial and temporal monitoring of irrigated pinto bean. *Precis. Agric.* 19: 555-569.
31. Cao Q, Miao Y, Shen J Yu W, Yuan F, Cheng S, Huang S, Wang H, Yang W, Liu F. 2016. Improving in-season estimation of rice yield potential and responsiveness to top dressing nitrogen application with crop circle active crop canopy sensor. *Precis. Agric.* 2016. 17: 136-154.
32. Lukas V, Novac J, Neudret L Svobodova I, Rodriguez-Moreno F, Edrees M, Kren J. 2016. The combination UAV survey and landsat imagery for monitoring of crop vigor in precision Agriculture. In *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Science, Proceedings of the 2016 XXIII ISPRS Congress, Prague, Czech Republic, 12-19 July 2016*. Available online: <http://www.int-arch-photogramm-remote-sens-spatial-inf-sci.net/XLI-B8/953/2016/>.
33. Khan MS, Semwal M, Sharma A, Verma RK. 2020. An Artificial neural network model for estimating Mentha Biomass Yield using Landsat 8 OLI. *Precis. Agric* 21: 18-33.
34. Pourazar H, Samadzadegan F, Javan FD. 2019. Aerial Multipectral imagery for plant disease detection: Radiometric calibration necessity assessment. *Eur. J. Remote Sens.* 52: 17-31.
35. De Lara A, Longchamps L, Khosla R. 2019. Soil water content and high-resolution imagery for precision irrigation: Maize Yield. *Agron J.* 9:174.
36. Martinez-Casasnovas JA, Uribeetxebarria A Escola A, Arno J. 2019. Santinel-2 vegetation indices and apparent electrical conductivity to predict barley (*Hordeum vulgare* L.) yield. In *precision Agriculture*; Wageningen Academic Publishers: Wageningen, The Netherlands. pp 415-421.
37. Shaver TM, Kruger GR, Rudnick DR. Crop canopy sensor orientation for late season nitrogen determination in corn. *J. Plant Nutri.* 2017. 40: 2217-2223.
38. DadrasJavan F, Samadzadegan F, Pourazar SHS, Fazeli H. 2019. UAV-based multispectral imagery for fast citrus greening detection. *J. Plant Dis. Protect.* 126: 307-318.
39. Venancio LP, Mantovani EC, Do Amaral CH, Neale CMU, Goncalves IZ, Filgueirans R, Compos I. 2019. Forecasting corn yield at the farm level in the Brazil based on the FAO-66 approach and soil-Adjusted vegetation index (SAVI). *Agric-water Manag.* 2019. 225: 105779.
40. Phadikar S and Goswami J. 2016. Vegetation indices based segmentation for automatic classification of brown spot and blast diseases of rice. In *Proceedings of the 3<sup>rd</sup> International Conference on recent Advances in information technology (RAIT), Dhanbad, India, 3 March 2016.*; pp. 284-289.

41. Lu J, Miao Y, Huang Y, Shi W, Hu X, Wang X, Wan J. 2015. Evaluating an unmanned Aerial Vehicle-based Remote Sensing System for Estimation of Rice Nitrogen status. In Proceedings of the Forth International Conference on Agro-Geoinformatics (Agro-geoinformatics), Istanbul, Turkey, 20 July 2015. pp 198-203.
42. Wolters S, Solderstrom M, Piikki K, Stenberg M. 2019. Near-real time winter N uptake from a combination of proximal and remote optical measurements: How to refine Sentinel-2 satellite image images for use in a precision agriculture decision support system. In Proceedings of the 12<sup>th</sup> European Conference on Precision Agriculture, Montpellier, France, 8-11 July 2019; Wageningen Academic Publishers: Wageningen, The Netherlands, 2019; pp. 415- 421.
43. Ranjan R, Chandel AK, Khot LR, Bahlol HY, Zhou J, Boydston RA, Miklas PN. Irrigated pinto bean crop stress and yield assessment using ground based low altitude remote sensing technology. *Inf. Process. Agric.* 2019. 6: 502-514.
44. Klem K, Zahora J, Zemek F, Trunda P, Tuma I Novotna K, Hodanova P, Rapantova b, Hanus J, Vavrikova J, et al. 2018. Interactive effects of water deficit and nitrogen nutrition on winter wheat. Remote sensing methods for their detection. *Agric. Water Manag.* 210: 171-184.
45. Liu P, Shi R, Gao W. 2018. Estimating leaf chlorophyll contents by combining multiple spectral indices with an artificial neural network. *Earth Sci. Inf.* 2018. 11: 147-156.
46. Meng J, Xu J, You X. 2015. Optimizing soybean harvest date using HJ-1 satellite imagery. *Precis. Agric.* 16: 164-179.
47. Shang J, Liu J, Ma B, Zaho T, Jiao X, Geng X, Huffman T, Kovacs JM, Walters D. 2015. Mapping Spatial variability of crop growth conditions using rapid eye data in Northern Ontario, Canada. *Remote sens. Environ.* 168: 113-125.
48. Rapaport T, Hochberg U, Cochavi A, Karnieli SC, Roberts P. the potential of the spectral 'water balance index' (WABI) for crop irrigation scheduling. *New Phytol* 2017. 216: 741-757.
49. Gao Y, Walker JP, Allahmoradi M, Moneris A, Ryu D, Jakson TJ. 2015. Optical sensing of vegetation water content: A synthesis study. *IEEE J. STARS.* 8: 1456-1464.
50. Malenovsky Ac AZ, Olejnickova J, Galle A, Rascher U and Mohammen G. 2015. Meta-analysis assessing potential of steady-state chlorophyll fluorescence for remote sensing detection of plant water, temperature and nitrogen stress. *Remote Sens. Environ.*, 168: 420-436.
51. Maxwell K, and Johason GN. 2000. Chlorophyll fluorescence-A practical guide. *J. Exp. Bot.*, 51: 659-668.
52. Lang M, Litchenthaler, Sowinskia M Heisel and Miehe JA. 1996. Fluorescence imaging of water and temperature stress in plant leaves. *J, Plant Physiol.*, 34: 2114-2126.
53. Kumari M, Patel NR, Raj R, Saha SK and Dadhwal VK. 2012. Parametric estimation of net photosynthesis in rice from *in-situ* spectral reflectance measurements. *Current Science.* 103: No. 1.
54. Dutta SK, Fangzauv, Jena S, Nath R, Mazumder and Chakraborty PK. 2011. Absorption of photosynthetically active radiation (PAR) and its effect on yield and dry matter production of rice at different dates of transplanting. *Journal of Crop and weed* 7 (2): 138-142.
55. Vadivambal R and Jayas DS. 2011. Applications of thermal imaging in agriculture and food industry- A review. *Food Bioprocess Technol.*, 4: 186-199.

56. Tilling AK, Oleary GJ, Ferwerda, Jones SD, Jones GJ, D Rodriguez and Belford R. 2007. Remote sensing of nitrogen and water stress in wheat. *Field crops Research.*, 104: 77-85.
57. Kumar V, Naresh RK, Kumar S, Kumar S, Kumar A, Gupta RK, RS Rathore, Singh SP, Dwivedi A, Tyagi S and Mahajan NC. 2017. Efficient nutrient management practices for sustaining soil health and improving Rice-Wheat Productivity: A review. *Journal of Pharmacognosy and Phytochemistry.* 7 (1): 585-597.
58. Shivagnam S, Duraisami VP, and Jagadeeswaran R. Effect of nitrogen on nitrogen content, pigment, growth and spectral reflectance of rice. 2015. *Trends in Biosciences* 8(8): 0974-8, 2126-2132.
59. XuYimming, Smith SE, Grunwaid S, Elrahman AA Wani SP Nair VD. Estimating total nitrogen in small holder farm setting using remote sensing spectral indices and regression kriging. *Catena* 163: 111-122.
60. Evans RG, LaRue J, Stone KC. King BA. 2013. Adoption of site-specific variable rate sprinkler irrigation Management systems. *Irrig. Sci.* 2013, 31: 871-887.
61. McDowell RW. 2017. Does variable rate irrigation decrease nutrient leaching losses from grazed dairy farming? *Soil Use Manag.* 33: 530-537.
62. Shanmugapriya P, Rathika S, Ramesh T and Janki P. 2019. Applications of remote sensing in agriculture – A Review. *Int. J. Curr. Microbiol. App. Sci.* 8 (01): 2270-2283.
63. Das KC, Singh J and Hazra J. 2018. Development of a high resolution soil moisture for precision Agriculture in India. A proceedings of 14<sup>th</sup> International Conference on Precision Agriculture. Le Central Sheraton, Montreal, Canada.
64. Wiess M, Jacob F, Duveillerc G. 2020. Remote sensing for agriculture applications: A meta-review. *Remote Sens. Environ.* 236: 111402.
65. Sishodhia R P, Ray RL, Singh SK. 2020. Applications of remote sensing in precision agriculture: A Review. *Remote Sens.* 12, 3136
66. Kaur R, Jaidka M and Kingra PK. 2013. Study of optimum time span for distinguishing *Rumexspinosus* in wheat crop through spectral reflectance characteristics. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.*
67. Kaur R, Jaidka M. 2014. Spectral reflectance characteristics to distinguish *Malvaneglecta* in wheat (*Triticumaestivum*). *Indian Journal of Agricultural Science.* 84(10): 1243-1249.
68. Franklin S. 2001. *Remote Sensing for sustainable Forest Management* Lewis publisher, Boca Raton Florida, p. 407.
69. Kundu. R, Dutta D, Nanda MK, Chakrabarty. A real time monitoring of potato Late blight disease severity using field hyperspectral Observation. *Smart Agric. Technol.* 2021. 1, 100019.
70. Riedell WE, Osborne SL and Heseler LS.2004. Insect pest and disease detection using remote sensing techniques. *Proceedings of 7<sup>th</sup> International Conference on Precision Agriculture.* Minneapolis MN USA.
71. William D, Stauffer M and Leung K.1979. A forester's look at the application of image manipulation techniques
72. Mahelin AK. 2016. Plant disease detection by imaging sensor-parallel and specific demands for precision agriculture and plant phenotyping. *Plant Dis.* 2016. 100: 241-251.
73. Mahlein AK, Rumpf T, Welke P, Dehne HW, Plumer L, Steiner U, Oerke EC. 2013. Development of spectral indices for detecting and identifying plant diseases. *Remote Sens. Environ.* 128:21-30.

74. AL-Saddik H, Simon J, Cointault F. 2017. Development of spectral disease indices for 'FlavescenceDoree' grapevine disease identification. *Sensors*. 17: 2772.

UNDER PEER REVIEW