

Case report

Abdominal Wall Schwannoma: A Case Report and Review of the Literature

Abstract:

Abdominal wall schwannomas are rare neoplasms originating from peripheral nerve sheaths. We present a case report of a 21-year-old patient with a ten-year history of a subcutaneous mass on the left flank of the abdomen. The mass exhibited a hard consistency and was adherent to the deep plane. Palpation elicited pain, and magnetic resonance imaging (MRI) initially suggested a dermoid cyst in intimate contact with the external oblique muscle. The patient underwent wide excision of the mass with a 2 cm margin and subsequent closure of the resulting muscle and aponeurotic defect by direct suturing. Histopathological examination confirmed the diagnosis of a completely excised subcutaneous schwannoma. This case emphasizes the importance of considering schwannomas as a differential diagnosis for abdominal wall masses and highlights the challenges in diagnosing and managing abdominal wall schwannomas. By presenting a comprehensive review of the literature, we aim to contribute to the existing knowledge of this rare entity, facilitating early recognition and appropriate management of abdominal wall schwannomas in clinical practice.

Keywords: abdominal wall, schwannoma, benign neoplasms, MRI

1. Introduction:

Schwannomas are benign neoplasms originating from Schwann cells, which are responsible for myelination of peripheral nerves. While they commonly occur in the head, neck, and extremities [1–3], schwannomas arising from the abdominal wall are extremely

rare[3–7]. We present a case of an abdominal wall schwannoma and provide a comprehensive review of the existing literature on this topic.

2. Case Presentation:

A 21-year-old patient presented with a subcutaneous mass on the left flank of the abdomen. The mass was discovered by the patient 10 years ago, initially very small and painless, but it had progressively increased in size, becoming painful over the last 6 months, which prompted the patient to seek medical consultation. Clinical examination revealed a 2 cm diameter subcutaneous mass, Well-circumscribed and with a hard consistency on the left flank of the abdomen. Upon palpation, the mass was painful and firmly adhered to the deep plane.

Magnetic resonance imaging initially suggested a dermoid cyst of the soft tissues in close proximity to the external oblique muscle (Figure 1).

We requested a review of the MRI images with another radiologist, as well as an additional ultrasound examination, but the same diagnosis was reiterated.

The mass's firm adherence to the deep plane and location raised our suspicions, leading us to perform an ultrasound-guided fine-needle aspirate biopsy. Unfortunately, the results were inconclusive and did not alleviate our suspicion of a malignant tumor.




Figure 1: Axial magnetic resonance imaging (MRI) of the abdomen reveals an oval-shaped, well-circumscribed, homogeneous mass in the left abdominal wall, in intimate contact with the left oblique muscle. The mass is hypointense on

T1-weighted images and shows intermediate signal intensity on T2-weighted images, with areas of fatty signal. There is no enhancement of the mass with contrast administration.

3. Treatment:

Considering the clinical presentation and radiological findings, the patient underwent surgical intervention. The strong adherence to the deep plane raised suspicion of a neoplastic process, prompting us to choose wide excision of the mass for safety. We maintained a 2 cm margin to ensure complete removal. Subsequently, we closed the resulting defect in the muscle and aponeurotic tissue using direct suturing (Figure 2).



Figure 2:

a: A subcutaneous mass on the left flank of the abdomen with a hard consistency was painful on palpation and firmly adhered to the deep plane.

b,c: Wide excision of the mass was performed with a 2 cm margin to ensure complete removal.

d: The resulting defect in the muscle and aponeurotic tissue was closed using direct suturing.

4. Histopathological Findings:

Histopathological examination of the surgical specimen revealed the characteristic appearance of a subcutaneous schwannoma. Microscopic evaluation demonstrated spindle-shaped cells arranged in interlacing fascicles, with Antoni A and Antoni B areas (Figure 3). Immunohistochemical staining for S100 protein was deemed unnecessary and was not performed.

At the 12-month follow-up, no recurrence was observed.




Figure 3: Microscopic findings: The tumor proliferation comprised spindle-shaped cells arranged in interlacing fascicles, with both Antoni A and Antoni B areas showing a benign appearance. There was no evidence of mitosis or cytonuclear atypia.

5. Literature Review:

We conducted a comprehensive review of published cases of abdominal wall schwannomas to gain further insights into this rare entity [3-18]. The literature search identified a total of 25 cases, including the present report, which were included in our analysis. The cases spanned a wide range of demographics, with patients ranging from 18 to 67 years of age. There was no apparent gender predilection, as schwannomas were reported in both males and females.

The clinical presentation of abdominal wall schwannomas varied among the cases [1,2]. The most common presenting symptom was the presence of a subcutaneous mass, which was often painless or associated with mild discomfort. Some patients reported a progressive increase in the size of the mass over time, while others noted a stable or slow-growing nature. Rarely, patients experienced pain or tenderness at the site of the mass. It is important to note that the clinical presentation of schwannomas is nonspecific and may overlap with other subcutaneous lesions, emphasizing the significance of thorough evaluation and histopathological confirmation.

Radiological imaging played a crucial role in the diagnosis and preoperative evaluation of abdominal wall schwannomas. Magnetic resonance imaging (MRI) was the most commonly employed imaging modality, followed by ultrasound and computed tomography (CT). Schwannomas typically appeared as well-defined masses with homogenous or heterogeneous signal intensity on MRI [10, 11]. They frequently exhibited a close association with peripheral nerves, which could aid in distinguishing them from other soft tissue tumors. However, the imaging features alone were often insufficient for a definitive diagnosis, emphasizing the importance of histopathological examination.

Histopathological examination remains the gold standard for diagnosing abdominal wall schwannomas. Microscopically, schwannomas exhibit characteristic features, including spindle-shaped cells arranged in interlacing fascicles and the presence of Antoni A and Antoni B areas [9]. Immunohistochemical staining for S100 protein, a marker of Schwann cells, is routinely performed to confirm the diagnosis. Positive staining for S100

protein supports the Schwann cell origin of the tumor and differentiates schwannomas from other soft tissue neoplasms [12].

Surgical excision with complete removal of the tumor is the primary treatment modality for abdominal wall schwannomas. The main goal of surgery is to achieve negative margins while preserving the integrity of surrounding structures. In the reviewed cases, various surgical approaches were employed, including wide local excision, enucleation, and excision with muscle or fascial reconstruction. The reported margins ranged from 0.5 cm to 5 cm. Despite the variability in surgical techniques, complete excision was achieved in the majority of cases, resulting in favorable outcomes and low recurrence rates.

Follow-up data were available for a subset of the reviewed cases. The duration of follow-up ranged from 6 months to 10 years. Recurrence was uncommon, with only a few reported cases [13-16]. Long-term outcomes were generally favorable, with no instances of malignant transformation or metastasis observed. The limited available data suggest that abdominal wall schwannomas have a benign clinical course, consistent with schwannomas arising in other anatomical locations.

In conclusion, our comprehensive review of the literature on abdominal wall schwannomas underscores the rarity of this entity and the challenges associated with its diagnosis and management. Radiological imaging, particularly MRI, can provide valuable information for preoperative evaluation. However, histopathological examination remains essential for a definitive diagnosis. Surgical excision with negative margins is the preferred treatment approach, resulting in excellent long-term outcomes. Further studies and case reports are needed to elucidate the pathogenesis, optimal surgical techniques, and long-term follow-up of abdominal wall schwannomas, enhancing our understanding and guiding clinical management decisions.

6. Discussion:

Abdominal wall schwannomas are exceedingly rare, with only a limited number of cases reported in the literature [5–7].

Malignant transformation is extremely rare in isolated lesions [2, 8, 9]. **Their clinical presentation can vary, often leading to diagnostic challenges.** In the present case, the patient presented with a subcutaneous mass on the left flank of the abdomen, which had been progressively growing over a ten-year period. The hard consistency of the mass and its adherence to the deep plane raised suspicion of a neoplastic process. However, the initial MRI findings suggested a dermoid cyst in intimate contact with the external oblique muscle. This highlights the importance of considering schwannomas as a differential diagnosis for abdominal wall masses, even in cases where imaging characteristics may initially resemble other soft tissue tumors.

Radiological imaging, including MRI, plays a crucial role in the evaluation of abdominal wall masses. Although schwannomas may have variable appearances on imaging, certain features can provide clues for their diagnosis. These features include a well-defined margin, homogeneity, and a close relationship to peripheral nerves [10, 11]. In our case, the initial imaging findings, while suggestive of a dermoid cyst, did not definitively exclude the possibility of a schwannoma. Therefore, histopathological examination remained crucial for an accurate diagnosis.

Histopathological examination of the surgical specimen confirmed the presence of a subcutaneous schwannoma. Microscopic evaluation demonstrated characteristic features of schwannomas, such as spindle-shaped cells arranged in interlacing fascicles and the presence of Antoni A and Antoni B areas [9]. Immunohistochemical staining for S100 protein, a marker of Schwann cells, can further support the diagnosis, but it's not mandatory. These findings align with previous reports in the literature, confirming the histological characteristics of schwannomas [12].

Surgical excision with complete removal of the tumor is the treatment of choice for abdominal wall schwannomas. The main goal of surgery is to achieve negative margins while minimizing damage to surrounding structures. In our case, wide excision of the mass with a 2 cm margin was performed, ensuring complete removal due to the suspicion of a neoplastic

process [13-16]. The resulting defect in the muscle and aponeurotic tissue was successfully closed by direct suturing. This approach aimed to restore the structural integrity of the abdominal wall while minimizing the risk of recurrence.

7. Conclusion

Abdominal wall schwannomas are rare neoplasms that should be considered in the differential diagnosis of subcutaneous masses. This case report highlights the challenges in diagnosing and managing abdominal wall schwannomas. By presenting a comprehensive review of the literature, we aim to contribute to the existing knowledge of this rare entity, facilitating early recognition and appropriate management of abdominal wall schwannomas in clinical practice.

Consent

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

References:

[1] Zhou H, Yao C, Dong Y, Alhaskawi A, Wang Z, Lai J, Ezzi SHA, Kota VG, Abdulla MHAH, Lu H. Clinical characteristics and management experience of schwannoma in extremities: Lessons learned from a 10-year retrospective study. *Front Neurol.* 2022 Dec 15;13:1083896.

[2] Schraepen C, Donkersloot P, Duyvendak W, Plazier M, Put E, Roosen G, Vanvolsem S, Wissels M, Bamps S. What to know about schwannomatosis: a literature review. *Br J Neurosurg.* 2022 Apr;36(2):171-174.

[3] R. Lam, B. L. Hunt, and O. Arreola-Owen, "Abdominal wall schwannoma," *Federal Practitioner*, vol. 36, no. 3, pp. 129–133, 2019.

[4] R. K. Bhatia, A. Banerjea, M. Ram, and B. E. Lovett, "Benign ancient schwannoma of the abdominal wall: an unwanted birthday present," *BMC Surgery*, vol. 10, no. 1, 2010.

[5] A. Mishra, M. Hamadto, M. Azzabi, and M. Elfagieh, "Abdominal wall schwannoma: case report and review of the literature," *Case Reports in Radiology*, vol. 2013, Article ID 456863, 4 pages, 2013.

[6] R. Balzarotti, F. Rondelli, J. Barizzi, and R. Cartolari, "Symptomatic schwannoma of the abdominal wall: a case report and review of the literature," *Oncology Letters*, vol. 9, no. 3, pp. 1095–1098, 2015.

[7] G. C. Ginesu, M. Puledda, C. F. Feo et al., "Abdominal wall schwannoma," *Journal of Gastrointestinal Surgery*, vol. 20, no. 10, pp. 1781–1783, 2016.

[8] D. A. Hilton and C. O. Hanemann, "Schwannomas and their pathogenesis," *Brain Pathology*, vol. 24, no. 3, pp. 205–220, 2014.

[9] F. J. Rodriguez, A. L. Folpe, C. Giannini, and A. Perry, "Pathology of peripheral nerve sheath tumors: diagnostic overview and update on selected diagnostic problems," *Acta Neuropathologica*, vol. 123, no. 3, pp. 295–319, 2012.

[10] Behzad B, Dianat S. Magnetic Resonance Imaging of Nerve Tumors. *Semin Musculoskelet Radiol.* 2022 Apr;26(2):172-181.

[11] J. Crist, J. R. Hodge, M. Frick et al., "Magnetic resonance imaging appearance of schwannomas from head to toe: a pictorial review," *Journal of Clinical Imaging Science*, vol. 7, p. 38, 2017.

[12] J. Guedes-Corrêa and R. Cardoso, "Immunohistochemical markers for schwannomas, neurofibromas and malignant peripheral nerve sheath tumors—what can the recent literature tell us?," *Arquivos Brasileiros de Neurocirurgia: Brazilian Neurosurgery*, vol. 37, no. 02, pp. 105–112, 2018.

[13] Estébanez-Ferrero B, Torres-Melero J, Fuentes-Porcel O, Reina-Duarte Á. Schwannoma in the abdominal wall. *Cir Esp (Engl Ed).* 2022 Oct 17:S2173-5077(22)00388-X.

[14] Lam R, Hunt BL, Arreola-Owen O. Abdominal Wall Schwannoma. *Fed Pract.* 2019 Mar;36(3):129-133.

[15]Ginesu GC, Puledda M, Feo CF, Cossu ML, Fancellu A, Addis F, Porcu A. Abdominal Wall Schwannoma.J Gastrointest Surg. 2016 Oct;20(10):1781-3.

[16]Tarchouli M, Essarghini M, Qamouss O, El Kharras A, Bounaim A. Abdominal wall schwannoma: a case report.Gastroenterol Hepatol Bed Bench. 2020 Winter;13(1):95-100.

[17]Liu Y, Chen X, Wang T, Wang Z.Imaging observations of a schwannoma of low malignant potential in the anterior abdominal wall: A case report.Oncol Lett. 2014 Sep;8(3):1159-1162.

[18]Alsahwan AG, Felemban JM, Al-Othman A, Assiri SY, Alzahir AA.Symptomatic Abdominal Wall Schwannoma Mimicking Infected Subcutaneous Soft Tissue Lesion. A Case report.Int J Surg Case Rep. 2021 Apr;81:105751.

UNDER PEER REVIEW