

## **INTEGRATIVE MOLECULAR APPROACHES TO PLANT DISEASE: A REVIEW**

**ABSTRACT:** Modern molecular and bioinformatics technologies have made understanding host-pathogen interactions easier. Plants have a multi-layered defense system against microbial diseases, including physical barriers, PAMP detection, and R genes that recognize pathogen-effector proteins and activate effector-triggered immunity. Plant pathogen genome databases provide genomic and phenotypic data on plant pathogen species and information on plant-pathogen interactions. Map-based or positional gene cloning is improving our understanding of plant-pathogen interactions, with R genes being used to develop resistance to pathogens. Plant genomes typically contain several hundred nucleotide-binding site-leucine-rich repeats (NLRs), with number, arrangement, and domain combinations varying by species. Bacterial blight (BB) severely impacts rice production and about 37 of 44 resistance genes have been mapped and 15 cloned. Many disease-resistant wheat cultivars have been developed using powdery mildew leaf rust (Lr)-resistance genes from wild relatives of *T. aestivum*. Over 140 genes are linked to powdery mildew resistance *T. aestivum* and MutChromSeq has found new target genes. Cloning Arabidopsis resistance genes is essential for developing resistant cultivars and understanding R gene evolution. Some R genes encode proteins with nucleotide-binding site (NBS) motifs, and an LRR protects against *Erysiphe cruciferarum* powdery mildew. CRISPR/Cas9 gene editing is a major tool in plant genome editing, efficiently introducing target site mutations, and improving plant immunity. High-throughput sequencing can identify and clone candidate resistance genes in different plant species, and gene editing technologies like CRISPR/Cas have illuminated site-specific mutagenesis and durable resistance.

**Keywords:** Defense, database, pathogen, resistance genes and clone.

### **1. INTRODUCTION:**

Phytopathogens threaten livelihoods and societal growth by reducing crop yields and quality. Modern molecular and bioinformatics technologies make host-pathogen interactions easier to understand [1]. Scientists can analyze massive amounts of biological data by sequencing many pathogens and plant species using whole-genome sequencing [2]. Plants have a multi-layered defense system against microbial diseases, including physical barriers, PAMP detection, and R genes that recognize pathogen-effector proteins and activate effector-triggered immunity [3]. Quantitative plant disease resistance is controlled by multiple genes or quantitative trait loci, while qualitative resistance is controlled by single R genes [4]. R proteins' disease resistance is linked to nucleotide binding (NB) and leucine-rich repeat (LRR) domains, which are divided into eight classes. CC-NBS-LRR (CNL) proteins, tobacco N, flax L6, and some RLK and RLP genes are resistance genes [5,6]. *Arabidopsis thaliana*, *Oryza sativa*, *Gossypium* sp., *Brassic napus*, and *B. rapa*, *Vitis vinifera*, *Triticum aestivum*, *Zeamays*, *Hordeum vulgare* [7,8–18]. Mega nucleases (MNs), zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and CRISPR/Cas are needed to improve plant health [19]. The CRISPR/Cas system is popular for its simplicity, low cost, efficiency, and reproducibility. CRISPR/Cas has been used to develop plant species' resistance to bacterial, fungal, and viral diseases [19].

### **2. Plant Pathogen Genome Databases**

Genome databases study genetic diversity, disease, and host-pathogen interactions [20]. PhytoPath uses Ensembl genome portals to provide genomic and phenotypic data on plant pathogen species. The NIAS Genebank preserves and documents Japanese agricultural plant, microorganism, and animal genetic resources [21]. PathoPlant, a database of plant-pathogen interactions, explains signal transduction during plant pathogenesis. PHI-base contains fungal and bacterial pathogens' experimentally verified pathogenicity, virulence, and effector genes [22]. Annotate, predict, and display host-pathogen interactions with HPIDB 3.0. Using integration and IMEx consortium curation protocols, Virus Mentha is a new resource for studying virus-virus and virus-host interactions. PCPPI is a large database for Penicillium-crop protein-protein interactions. Microorganism genome databases can amplify data, but plant pathogen genome databases are needed to understand disease resistance mechanisms [23].

### 3. R genes and Plant NLRs

Map-based or positional gene cloning is improving our understanding of plant-pathogen interactions. Genetic map positions are used to isolate genes with known phenotype and genomic locations. Hm1 from *Z. mays* was the first R gene cloned. *Z. mays* developed resistance to *Chlioboluscarbonum*'s HC toxin [8]. Flax R genes include Cf-9, N, RPS2, RPM1, and L6. Advanced plant genomics and genetic engineering techniques make it easier to clone R genes from crops or their wild relatives and transfer them into elite breeding lines or cultivars [24].

#### 3.1.1 Plant NLRs

Diverse genes encode nucleotide-binding site-leucine-rich repeats (NLRs) in two main classes: CNL (coiled-coil) and TNL (Toll/interleukin-1 receptor) domains. Plants, animals, fungi, and protists have many NLR proteins. Plant genomes typically contain several hundred NLRs, with number, arrangement, and domain combinations varying by species [25]. *Carica papaya*, *Cucumis sativus*, and *Citrullus lanatus* have low NLR copy numbers. There is no correlation between genome size and gene count [26, 27, 28, 29,30].

#### 3.1.2 R Genes in *O. sativa*

Bacterial blight (BB) severely impacts rice production. About 37 of 44 BB resistance genes have been mapped and 15 cloned [7,18,31]. These genes are classified as receptor-like kinase (RLK), sugar will eventually be exported transporter (SWEET), executor genes (Xa10), Xa23 and other genes (Xa1 and Xa5). Over 100 R genes and 27 clones have been found in *Magnaportheoryzae*-caused rice blast. *Pia* protects against blast fungus *M. oryzae* with AVR-*Pia* [32–35]. Pi25, Pi36, Pi-64(t), and Pi-jnw1 from cultivar AS20-1 are blast resistance genes [4,12,36].

#### 3.1.3 R Genes in *T. aestivum*

Breeding programs have developed disease-resistant wheat cultivars using powdery mildew leaf rust (Lr)-resistance genes[91]. First resistance genes Lr10, Lr21, and Lr1 were cloned in *T. aestivum*. Most of the 80 Lr genes characterized are from wild relatives of *T. aestivum*. Over 60 stem rust (Sr)-resistant genes have been found in wild relatives of *T. aestivum* [37–39]. *Blumeriagraminis* f. sp. *tritici* is a common disease of *T.aestivum* and causes severe yield loss. Over 140 T genes are linked to powdery mildew resistance *T.aestivum*. Pm2 [14], Pm2a [15], Pm3 [40], Pm3b [41], Pm3c and Pm3b [42], Pm5e [17], Pm8 [13], Pm17 [43], Pm21 [16], Pm24 [44], Pm41 [45], Pm60 [46], PmR1 [47], and Pm2b [91] have been cloned using map-based cloning and sequencing. MutChromSeq has found new target genes. Target-sequence Enrichment Sequencing (TEnSeq) pipelines have cloned Pm genes, most of which contain an NLR, while resistance genes Pm38 and Pm46 encode ATP-Binding Cassette (ABC) and hexose transporters, respectively [48, 49].

#### 3.1.4 R genes in *Z. mays*

Fungal diseases threaten global maize production. Hm-1 was the first gene cloned against northern corn leaf blight and *Cochlioboluscarbonum* [8]. A devastating fungal disease caused by *Setosphaeria turcica*. A map-based cloning method identified and cloned four resistance genes, Ht1, Ht2, Ht3, and Htn1. Htn1 works against the most common NCLB races but depends on maize genotype and environment. Only sixteen maize resistance genes have been cloned, with RppC being the most common. *Puccinia polysora* causes southern corn rust (SCR), the most common disease in the US, Canada, Brazil, and China [108]. *Fusarium graminearum* Gibberella stalk rot in maize is greatly improved by qRfg1, a major resistance quantitative trait locus. Map-based cloning was used to clone CCT domain-containing gene ZmCCT. *Rhizoctonia solani* banded leaf and sheath blight is resistant to ZmFBL41. MDR is a useful tool for developing durable resistance, but only one maize MDR gene (ZmMM1) has been cloned [51, 52]. In maize-growing regions, Rab GDP dissociation inhibitor alpha (RabGDI $\alpha$ ) is a host susceptibility factor for rice black-streaked dwarf virus infections. Sugarcane mosaic virus (SCMV) is a severe maize disease, and Scmv1 and Scmv2 provide complete resistance [53,54].

#### 3.1.5 R Genes in *A. thaliana*

Cloning Arabidopsis resistance genes is essential for developing resistant cultivars and understanding R gene evolution. Most Arabidopsis R genes are TIR-NBS-LRR or LZ-NBS-LRR, with receptor-like kinases (RLKs) involved in plant development and defense [55,56]. The TIR-NBS-LRR subclass has an N-terminal region that

resembles the cytoplasmic domain of Toll and interleukin1 transmembrane receptors (TIRs), while the LZ subclass has a leucine zipper-like motif (LZ) [57,58,59,60]. Some R genes encode proteins with nucleotide-binding site (NBS) motifs, and an LRR protects against *Erysiphe cruciferarum* powdery mildew. Defense-signaling components EDS1, NDR1, and PBS1 are linked to RPP4-mediated resistance [61, 62]. RFO, which is identical to WAKL22 in Arabidopsis, was characterized using map-based cloning. RPS5 is an NBS-LRR subclass, and cloning RPS5 genes has helped characterize two rps5-1 mutations that affect several R genes and confer resistance to both pathogens. In *Arabidopsis*, both subclasses confer resistance to *P. parasitica* and *P. syringae*, while RCY1 belongs to the CC-NB-LRR subclass and confers viral resistance [63].

### 3.1.6 R genes in *S. lycopersicum*

The tomato genome has been extensively studied to understand resistance loci structure and organization. *S. lycopersicum*, has over 100 loci, including disease-resistant genes Pto, Ptil, and Fen. which resist *P. syringae* pv. *tomato* bacterial speck [64]. Cf-2 and Cf-9 from *Solanum pimpinellifolium* and Cf-4 and Cf-5 from *Solanum peruvianum* have been transferred into cultivated species to develop resistance to *Cladosporium fulvum*. Sw-5, introduced from the wild species *S. peruvianum*, developed resistance to TSWV, TCSV, and GRSV [60]. *Solanum* sp. has over 60 *P. infestans* resistance genes (Rpi genes). The first Rpi gene was Ph-1, followed by Ph-2 and Ph-2 in *S. pimpinellifolium* [65]. These genes created disease-resistant cultivars like *S. lycopersicum*, *S. pimpinellifolium*, *S. peruvianum*, *C. Fulvum*, *F. oxysporum*, *G. rostockiensis*, *M. javanica*, *P. syringae*, *V. dahliae*, TMV, TSWV, Ac-Ds, CC, NBS-LRR, TIR.

## 3.2 NLR Annotation Tools

### 3.2.1 NLR-Parser

Java application NLR-Parser quickly annotates plant genome NLR complements. MAST output from translated amino acid sequences filters biologically curated motif compositions. The output shows the start and frame of the first NLR-specific motif, whether it is a TNL or CNL, and whether it is complete or fragmented. The NB-ARC domain sequence is useful for phylogenetics. NLR-parser searches for the complete set of NLR protein motifs to distinguish pseudogenes. Download the tool from Git-Hub [66].

### 3.2.2 NLR-Annotator

NLR-Annotator annotates genomic sequence data NLR loci as an extension of NLR-Parser. It splits genomic sequences into overlapping fragments and selects NLR-loci with NLR-Parser. This method can find NLR loci candidates in stem rust, leaf rust, powdery mildew, and yellow rust resistance genes [67].

### 3.2.3 NL Genome Sweeper

NLGenome Sweeper uses the NB-ARC domain to search genomes for NLR disease-resistance genes. It merges NBS-LRR species' customized sequences into one fasta file. With high specificity for complete genes and structurally complete pseudogenes, the pipeline found 152 NBS-LRR proteins. 140 matches the manually annotated Arabidopsis NLR set with 96% sensitivity [68].

### 3.2.4 NLRtracker

In NLRtracker, InterProScan and predefined NLR motifs annotate proteome or transcriptome sequences and extract NLRs from the RefPlantNLR dataset based on core features. The NB-ARC domain is extracted for comparative phylogenetic analysis [69].

## 4. DRAGO2

DRAGO 2, an online platform for analyzing and predicting plant disease-resistance genes, accepts FASTA DNA or protein sequences. A Perl script predicts pathogen receptor genes (PRGs) and LRR, kinase, NBS, and TIR domains in PRGdb. DRAGO2 tools predicted over 1700 PRGs with the highest sensitivity [70].

## 5. CRISPR Gene Editing

Its simplicity, flexibility, success rate, and cost-effectiveness have made the CRISPR/Cas9 system a major tool in plant genome editing. This system efficiently introduces target site mutations, including INDELS and base

substitutions [71] and other plant species have benefited from it. *A. thaliana*, *O. sativa*, *N. Tabacum*, *S. bicolor*, *T. aestivum*, *Z. G. mays*, *S. lycopersicum*, *S. tuberosum*, *P. alba*, *M. domestica* and *Musa* species to fight viruses, fungi, and bacteria [72, 73, 74]. The CRISPR/Cas9 system can be used to develop plant disease resistance by knocking out disease susceptibility genes (e.g., MLO), deleting or modifying cis-elements in promoters, modifying the amino acid sequence of surface receptor proteins to suppress secreted pathogen effectors, knocking down negative regulators of plant immunity, and modifying central regulators of the defense response. The system has developed powdery mildew fungus O-resistant varieties. *O. neolyopersici*, PMR4, *Erysiphe necator*, and *Plasmopara viticola* in *V. vinifera* [75, 76, 77, 78]. The CRISPR/Cas9 system provides primers for guide activity and target validation, as well as accurate sgRNA design tools. It designs sgRNAs in specific regions, checks for target specificity and genomic context, predicts target site specificity and sgRNA design for different CRISPR/Cas systems, provides on-target and off-target scoring and gRNA sequence analysis, and designs and constructs sgRNAs for CRISPR-Cas9-mediated genome editing [79]. APETELA2/ERF superfamily ethylene-responsive factors (ERFs) are essential for biotic stress adaptation. Fungus-resistant rice blast. *M. oryzae* was enhanced by CRISPR/Cas9-mediated ERF922 gene mutation. AP2/ERF transcription factor knockdown reduced abscisic acid accumulation and increased *M. oryzae* resistance. Overexpressing defense genes is a key biotechnological tool for plant pathogen resistance [80, 81].

MORC proteins are essential nuclear regulators in prokaryotes and eukaryotes, silencing transcriptional genes and stabilizing genomes. MORC1 was found in *Arabidopsis* to protect against turnip crinkle virus (TCV), and AtMORC1, AtMORC2 and AtMORC6 are involved in multiple defenses against *P. syringae* and *Hyaloperonospora arabidopsidis*. The *Streptococcus pyogenes* CRISPR-Cas9 system was used to mutate HvMORC1 and HvMORC6a genes in *H. vulgare*. Also studied MORCs in *S. tuberosum* (StMORC1), *Nicotianabenthhamiana* (NbMORC1) and SIMORC1 [82, 83]. The immune responses of various plants involve WRKY transcription factors. Mutant analyses in *Arabidopsis* have linked specific WRKY proteins to *P. syringae* defense responses. The susceptible variety (IR64) of *O. sativa* was resistant to rice tungro streak spherical virus after a CRISPR/Cas9-mediated mutation in eIF4G. *Arabidopsis* and cucumber gained resistance to turnip mosaic virus and cucumber vein yellowing virus after recessive eIF4E gene mutation. Silencing the eIF4E gene has shown RNA virus resistance in *S. lycopersicum* and *C. melo* [84, 85].

FnCas9 (Cas endonucleases) from *Francisellana novicida* may be a new tool for attacking plant RNA virus genomes. *N. benthamiana* and *Arabidopsis* was resistant to CMV and TMV using FnCas9 [86, 87]. The single effector Cas13a protein of *Leptotrichia shahii* (LshCas13a) was a programmable RNA-guided single-stranded RNA (ssRNA) ribonuclease that protected the bacteria from bacteriophages [192]. *O. sativa* was resistant to SRBSDV and RSMV using the LshCas13a system. The simple operation, good knockout effect, low cytotoxicity, high specificity, and universal applicability of the CRISPR/Cas system have made it more effective for disease resistance development by targeting the pathogen genome or host genes to interfere with susceptibility. Since CRISPR/Cas-induced mutations create pathogen-resistant genotypes when natural populations or wild relatives lack resistance resources, the system has garnered attention [89]. Intracellular NLR receptors recognize pathogen effectors and activate the immune system. The mechanisms of plant NLR activation are unknown, but animal NLRs oligomerize to activate downstream signalling after binding to their effectors. CNL and TNL resist some structural data has greatly improved our understanding of plant NLR activation. Genome editing advances will benefit sustainable agriculture [86, 87, 90].

## 6. Conclusions

Plant immunity relies on NLRs to activate the strong resistance response that resists disease. Central nucleotide-binding (NB) and leucine-rich repeat (LRR) domains [91]. Understanding disease resistance is crucial, and high-throughput sequencing can identify and clone candidate resistance genes in different plant species. By pyramiding genes or altering plant and pathogen genomes, gene editing technologies like CRISPR/Cas have illuminated site-specific mutagenesis and durable resistance. Protein engineering has revolutionized NLR molecular recognition, and engineered intracellular immune receptors may improve disease resistance. NLR protein research is limited by the lack of three-dimensional structures and homology models. Modern computational technology like AlphaFold and cryo-electron microscopy have illuminated NLR biological mechanisms and functional complexity. Designer NLR receptors may give crop plants broad-spectrum resistance using these technologies. Understanding protein structures, ligand binding, and host-pathogen interactions requires more tools.

## References:

1. Ali, Q.; Zheng, H.; Rao, M.J.; Ali, M.; Hussain, A.; Saleem, M.H.; Nehela, Y.; Sohail, M.A.; Ahmed, A.M.; Kubar, K.A.; et al. Advances, limitations, and prospects of biosensing technology for detecting phytopathogenic bacteria. *Chemosphere* 2022, 296, 133773.
2. Mu, H.; Wang, B.; Yuan, F. Bioinformatics in Plant Breeding and Research on Disease Resistance. *Plants* 2022, 11, 3118.
3. Boutrot, F.; Zipfel, C. Function, discovery, and exploitation of plant pattern recognition receptors for broad-spectrum disease resistance. *Annu. Rev. Phytopathol.* 2017, 55, 257–286.
4. Srivastava, D.; Shamim, M.; Kumar, M.; Mishra, A.; Pandey, P.; Kumar, D.; Yadav, P.; Siddiqui, M.H.; Singh, K.N. Current status of conventional and molecular interventions for blast resistance in rice. *Rice Sci.* 2017, 24, 299–321.
5. Wei, C.; Kuang, H.; Li, F.; Chen, J. The I2 resistance gene homologues in *Solanum* have complex evolutionary patterns and are targeted by miRNAs. *BMC Genom.* 2014, 15, 743.
6. Hatsugai, N.; Hillmer, R.; Yamaoka, S.; Hara-Nishimura, I.; Katagiri, F. The  $\mu$  Subunit of Arabidopsis Adaptor Protein-2 Is Involved in Effector-Triggered Immunity Mediated by Membrane-Localized Resistance Proteins. *Mol. Plant Microbe Interact.* 2016, 29, 345–351.
7. Song, W.Y.; Wang, G.L.; Chen, L.L.; Kim, H.S.; Pi, L.Y.; Holsten, T.; Gardner, J.; Wang, B.; Zhai, W.X.; Zhu, L.H.; et al. A receptor kinase-like protein encoded by the rice disease resistance gene, Xa21. *Science* 1995, 270, 1804–1806.
8. Johal, G.S.; Briggs, S.P. Reductase activity encoded by the HM1 disease resistance gene in maize. *Science* 1992, 258, 985–987.
9. Wang, Z.X.; Yano, M.; Yamanouchi, U.; Iwamoto, M.; Monna, L.; Hayasaka, H.; Katayose, Y.; Sasaki, T. The Pib gene for rice blast resistance belongs to the nucleotide binding and leucine-rich repeat class of plant disease resistance genes. *Plant J.* 1999, 19, 55–64.
10. Qu, S.; Liu, G.; Zhou, B.; Bellizzi, M.; Zeng, L.; Dai, L.; Han, B.; Wang, G.L. The broad-spectrum blast resistance gene Pi9 encodes a nucleotide-binding site-leucine-rich repeat protein and is a member of a multigene family in rice. *Genetics* 2006, 172, 1901–1914.
11. Yadav, M.K.; Aravindan, S.; Ngangkham, U.; Raghu, S.; Prabhukarthikeyan, S.R.; Keerthana, U.; Marndi, B.C.; Adak, T.; Munda, S.; Deshmukh, R.; et al. Blast resistance in Indian rice landraces: Genetic dissection by gene-specific markers. *PLoS ONE* 2019, 14, e0211061.
12. Liu, X.; Lin, F.; Wang, L.; Pan, Q. The in-silico map-based cloning of Pi36, a rice coiled-coil nucleotide-binding site leucine-rich repeat gene that confers race-specific resistance to the blast fungus. *Genetics* 2007, 176, 2541–2549.
13. Hurni, S.; Brunner, S.; Buchmann, G.; Herren, G.; Jordan, T.; Krukowski, P.; Wicker, T.; Yahiaoui, N.; Mago, R.; Keller, B. Rye *Pm8* and wheat *Pm3* are orthologous genes and show evolutionary conservation of resistance function against powdery mildew. *Plant J.* 2013, 76, 957–969.
14. Sánchez-Martín, J.; Steuernagel, B.; Ghosh, S.; Herren, G.; Hurni, S.; Adamski, N.; Vrána, J.; Kubaláková, M.; Krattinger, S.G.; Wicker, T.; et al. Rapid gene isolation in barley and wheat by mutant chromosome sequencing. *Genome Biol.* 2016, 17, 221.
15. Steuernagel, B.; Vrána, J.; Karafiátová, M.; Wulff, B.B.H.; Doležel, J. Rapid Gene Isolation Using MutChromSeq. *Methods Mol. Biol.* 2017, 1659, 231–243.
16. Xing, L.; Hu, P.; Liu, J.; Witek, K.; Zhou, S.; Xu, J.; Zhou, W.; Gao, L.; Huang, Z.; Zhang, R.; et al. Pm21 from *Haynaldia villosa* encodes a CC-NBS-LRR protein conferring powdery mildew resistance in wheat. *Mol. Plant.* 2018, 11, 874–878.
17. Xie, J.; Guo, G.; Wang, Y.; Hu, T.; Wang, L.; Li, J.; Qiu, D.; Li, Y.; Wu, Q.; Lu, P.; et al. A rare single nucleotide variant in *Pm5e* confers powdery mildew resistance in common wheat. *New Phytol.* 2020, 228, 1011–1026.
18. Yang, Y.; Zhou, Y.; Sun, J.; Liang, W.; Chen, X.; Wang, X.; Zhou, J.; Yu, C.; Wang, J.; Wu, S.; et al. Research Progress on Cloning and Function of *Xa* Genes Against Rice Bacterial Blight. *Front Plant Sci.* 2022, 13, 847199.
19. Karmakar, S.; Das, P.; Panda, D.; Xie, K.; Baig, M.J.; Molla, K.A. A detailed landscape of CRISPR-Cas-mediated plant disease and pest management. *Plant Sci.* 2022, 323, 111376.
20. Pedro, H.; Maheswari, U.; Urban, M.; Irvine, A.G.; Cuzick, A.; McDowall, M.D.; Staines, D.M.; Kulesha, E.; Hammond-Kosack, K.E.; Kersey, P.J. PhytoPath: An integrative resource for plant pathogen genomics. *Nucleic Acids Res.* 2016, 44, D688–D693.
21. Takeya, M.; Yamasaki, F.; Uzuhashi, S.; Aoki, T.; Sawada, H.; Nagai, T.; Tomioka, K.; Tomooka, N.; Sato, T.; Kawase, M. NIASGBdb: NIAS Genebank databases for genetic resources and plant disease information. *Nucleic Acids Res.* 2011, 39, D1108–D1113.
22. Urban, M.; Cuzick, A.; Seager, J.; Wood, V.; Rutherford, K.; Venkatesh, S.Y.; Sahu, J.; Iyer, S.V.; Khamari, L.; De Silva, N.; et al. PHI-base in 2022: A multi-species phenotype database for Pathogen-Host Interactions. *Nucleic Acids Res.* 2022, 50, D837–D847.
23. Dong, A.Y.; Wang, Z.; Huang, J.J.; Song, B.A.; Hao, G.F. Bioinformatic tool support decision-making in plant disease management. *Trends Plant Sci.* 2021, 26, 953–967.
24. Lawrence, G.J.; Finnegan, E.J.; Ayliffe, M.A.; Ellis, J.G. The L6 Gene for Flax Rust Resistance Is Related to the Arabidopsis Bacterial Resistance Gene RPS2 and the Tobacco Viral Resistance Gene N. *Plant Cell* 1995, 7, 1195–1206.

25. Kourelis, J.; vander Hoorn, R.A.L. Defended to the Nines: 25 Years of Resistance Gene Cloning Identifies Nine Mechanisms for R Protein Function. *Plant Cell* 2018, *30*, 285–299.
26. Meyers, B.C.; Kozik, A.; Griego, A.; Kuang, H.; Michelmore, R.W. Genome-wide analysis of NBS-LRR-encoding genes in Arabidopsis. *Plant Cell* 2005, *15*, 809–834.
27. Wan, H.; Yuan, W.; Bo, K.; Shen, J.; Pang, X.; Chen, J. Genome-wide analysis of NBS-encoding disease resistance genes in Cucumis sativus and phylogenetic study of NBS-encoding genes in Cucurbitaceae crops. *BMC Genom.* 2013, *14*, 109.
28. Lin, X.; Zhang, Y.; Kuang, H.; Chen, J. Frequent loss of lineages and deficient duplications accounted for low copy number of disease resistance genes in Cucurbitaceae. *BMC Genom.* 2013, *14*, 335.
29. Velasco, R.; Zharkikh, A.; Affourtit, J.; Dhingra, A.; Cestaro, A.; Kalyanaraman, A.; Fontana, P.; Bhatnagar, S.K.; Troggio, M.; Pruss, D.; et al. The genome of the domesticated apple (*Malus domestica* Borkh.). *Nat. Genet.* 2010, *42*, 833–839.
30. Baggs, E.; Dagdas, G.; Krasileva, K.V. NLR diversity, helpers and integrated domains: Making sense of the NLR identity. *Curr. Opin. Plant Biol.* 2017, *38*, 59–67.
31. Chen, L.; Yin, F.; Zhang, D.; Xiao, S.; Zhong, Q.; Wang, B.; Ke, X.; Ji, Z.; Wang, L.; Zhang, Y.; et al. Unveiling a Novel Source of Resistance to Bacterial Blight in Medicinal Wild Rice, *Oryza officinalis*. *Life* 2022, *12*, 827.
32. Liu, G.; Lu, G.; Zeng, L.; Wang, G.L. Two broad-spectrum blast resistance genes, Pi9(t) and Pi2(t), are physically linked on rice chromosome 6. *Mol. Genet Genom.* 2002, *267*, 472–480. [CrossRef]
33. Shang, J.; Tao, Y.; Chen, X.; Zou, Y.; Lei, C.; Wang, J.; Li, X.; Zhao, X.; Zhang, M.; Lu, Z.; et al. Identification of a new rice blast resistance gene, Pi-d3, by genome-wide comparison of paired nucleotide-binding site-leucine-rich repeat genes and their pseudogene alleles between the two sequenced rice genomes. *Genetics* 2009, *182*, 1303–1311.
34. Okuyama, Y.; Kanzaki, H.; Abe, A.; Yoshida, K.; Tamiru, M.; Saitoh, H.; Fujibe, T.; Matsumura, H.; Shenton, M.; Galam, D.C.; et al. A multifaceted genomics approach allows the isolation of the rice Pi-a blast resistance gene consisting of two adjacent NBS-LRR protein genes. *Plant J.* 2011, *66*, 467–479.
35. Das, A.; Soubam, D.; Singh, P.K.; Thakur, S.; Singh, N.K.; Sharma, T.R. A novel blast resistance gene, Pi54rh cloned from wild species of rice, *Oryza rhizomatis* confers broad spectrum resistance to Magnaporthe oryzae. *Funct. Integr. Genom.* 2012, *12*, 215–228.
36. Chen, J.; Shi, Y.; Liu, W.; Chai, R.; Fu, Y.; Zhuang, J.; Wu, J. A Pi-d3 allele from rice cultivar Gumei 2 confers resistance to Magnaporthe oryzae. *J. Genet Genom.* 2011, *38*, 209–216.
37. Feuillet, C.; Travella, S.; Stein, N.; Albar, L.; Nublat, A.; Keller, B. Map-based isolation of the leaf rust disease resistance gene Lr10 from the hexaploid wheat (*Triticum aestivum* L.) genome. *Proc. Natl. Acad. Sci. USA* 2003, *100*, 15253–15258.
38. Huang, L.; Brooks, S.A.; Li, W.; Fellers, J.P.; Trick, H.N.; Gill, B.S. Map-based cloning of leaf rust resistance gene Lr21 from the large and polyploid genome of bread wheat. *Genetics* 2003, *164*, 655–664.
39. Cloutier, S.; McCallum, B.D.; Loutre, C.; Banks, T.W.; Wicker, T.; Feuillet, C.; Keller, B.; Jordan, M.C. Leaf rust resistance gene Lr1, isolated from bread wheat (*Triticum aestivum* L.) is a member of the large psr567 gene family. *Plant Mol. Biol.* 2007, *65*, 93–106.
40. Brunner, S.; Hurni, S.; Streckeisen, P.; Mayr, G.; Albrecht, M.; Yahiaoui, N.; Keller, B. Intragenic allelic pyramiding combines different specificities of wheat Pm3 resistance alleles. *Plant J.* 2010, *64*, 433–445.
41. Yahiaoui, N.; Srichumpa, P.; Dudler, R.; Keller, B. Genome analysis at different ploidy levels allows cloning of the powdery mildew resistance gene Pm3b from hexaploid wheat. *Plant J.* 2004, *37*, 528–538.
42. Srichumpa, P.; Brunner, S.; Keller, B.; Yahiaoui, N. Allelic series of four powdery mildew resistance genes at the Pm31 locus in hexaploid bread wheat. *Plant Physiol.* 2004, *139*, 885–895.
43. Singh, S.P.; Hurni, S.; Ruinelli, M.; Brunner, S.; Sanchez-Martin, J.; Krukowski, P.; Peditto, D.; Buchmann, G.; Zbinden, H.; Keller, B. Evolutionary divergence of the *Pm17* and *Pm8* resistance genes reveals ancient diversity. *Plant Mol. Biol.* 2018, *98*, 249–260.
44. Lu, P.; Guo, L.; Wang, Z.; Li, B.; Li, J.; Li, Y.; Qiu, D.; Shi, W.; Yang, L.; Wang, N.; et al. A rare gain of function mutation in wheat tandem kinase confers resistance to powdery mildew. *Nat. Commun.* 2020, *11*, 680.
45. Li, M.; Dong, L.; Li, B.; Wang, Z.; Xie, J.; Qiu, D.; Li, Y.; Shi, W.; Yang, L.; Wu, Q.; et al. A CNL protein in wild emmer wheat confers powdery mildew resistance. *New Phytol.* 2020, *228*, 1027–1037.
46. Zou, S.; Shi, W.; Ji, J.; Wang, H.; Tang, Y.; Yu, D.; Tang, D. Diversity and similarity of wheat powdery mildew resistance among three allelic functional genes at the Pm60 locus. *Plant J.* 2022, *110*, 1781–1790.
47. Zou, S.; Wang, H.; Li, Y.; Kong, Z.; Tang, D. The NBS-LRR gene Pm60 confers powdery mildew resistance in wheat. *New Phytol.* 2018, *218*, 298–309.
48. Moore, J.W.; Herrera-Foessel, S.; Lan, C.; Schnippenkoetter, W.; Ayliffe, M.; Huerta-Espino, J.; Lillemo, M.; Viccars, L.; Milne, R.; Periyannan, S.; et al. A recently evolved hexose transporter variant confers resistance to multiple pathogens in wheat. *Nat. Genet.* 2015, *47*, 1494–1498.
49. Krattinger, S.G.; Kang, J.; Bräunlich, S.; Boni, R.; Chauhan, H.; Selter, L.L.; Robinson, M.D.; Schmid, M.W.; Wiederhold, E.; Hensel, G.; et al. Abscisic acid is a substrate of the ABC transporter encoded by the durable wheat disease resistance gene Lr34. *New Phytol.* 2019, *223*, 853–866.

50. Deng, C.; Leonard, A.; Cahill, J.; Lv, M.; Li, Y.; Thatcher, S.; Li, X.; Zhao, X.; Du, W.; Li, Z.; et al. The RppC-AvrRppC NLR-effector interaction mediates the resistance to southern corn rust in maize. *Mol. Plant.* 2022, *15*, 904–912.
51. Li, N.; Lin, B.; Wang, H.; Li, X.; Yang, F.; Ding, X.; Yan, J.; Chu, Z. Natural variation in ZmFBL41 confers banded leaf and sheath blight resistance in maize. *Nat. Genet.* 2019, *51*, 1540–1548.
52. Wang, H.; Hou, J.; Ye, P.; Hu, L.; Huang, J.; Dai, Z.; Zhang, B.; Dai, S.; Que, J.; Min, H.; et al. A teosinte-derived allele of a MYB transcription repressor confers multiple disease resistance in maize. *Mol. Plant.* 2021, *14*, 1846–1863.
53. Liu, Q.; Liu, H.; Gong, Y.; Tao, Y.; Jiang, L.; Zuo, W.; Yang, Q.; Ye, J.; Lai, J.; Wu, J.; et al. An Atypical Thioredoxin Imparts Early Resistance to Sugarcane Mosaic Virus in Maize. *Mol. Plant.* 2017, *10*, 483–497.
54. Leng, P.; Ji, Q.; Asp, T.; Frei, U.K.; Ingvarsdson, C.R.; Xing, Y.; Studer, B.; Redinbaugh, M.; Jones, M.; Gajjar, P.; et al. Auxin Binding Protein 1 Reinforces Resistance to Sugarcane Mosaic Virus in Maize. *Mol. Plant.* 2017, *10*, 1357–1360.
55. Gómez-Gómez, L.; Boller, T. FLS2: An LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. *Mol. Cell* 2000, *5*, 1003–1011.
56. Kim, M.H.; Kim, Y.; Kim, J.W.; Lee, H.S.; Lee, W.S.; Kim, S.K.; Wang, Z.Y.; Kim, S.H. Identification of Arabidopsis BAK1-associating receptor-like kinase 1 (BARK1) and characterization of its gene expression and brassinosteroid-regulated root phenotypes. *Plant Cell Physiol.* 2013, *54*, 1620–1634.
57. Bent, A.F.; Kunkel, B.N.; Dahlbeck, D.; Brown, K.L.; Schmidt, R.; Giraudat, J.; Leung, J.; Staskawicz, B.J. RPS2 of Arabidopsis thaliana: A leucine-rich repeat class of plant disease resistance genes. *Science* 1994, *265*, 1856–1860.
58. Grant, M.R.; Godiard, L.; Straube, E.; Ashfield, T.; Lewald, J.; Sattler, A.; Innes, R.W.; Dangl, J.L. Structure of the Arabidopsis RPM1 gene enabling dual specificity disease resistance. *Science* 1995, *269*, 843–846.
59. McDowell, J.M.; Dhandaydham, M.; Long, T.A.; Aarts, M.G.; Goff, S.; Holub, E.B.; Dangl, J.L. Intragenic recombination and diversifying selection contribute to the evolution of downy mildew resistance at the RPP8 locus of Arabidopsis. *Plant Cell* 1998, *10*, 1861–1874.
60. Bittner-Eddy, P.D.; Crute, I.R.; Holub, E.B.; Beynon, J.L. RPP13 is a simple locus in Arabidopsis thaliana for alleles that specify downy mildew resistance to different avirulence determinants in *Peronospora parasitica*. *Plant J.* 2000, *21*, 177–188.
61. Aarts, N.; Metz, M.; Holub, E.; Staskawicz, B.J.; Daniels, M.J.; Parker, J.E. Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signaling pathways in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 1998, *95*, 10306–10311.
62. Century, K.S.; Shapiro, A.D.; Repetti, P.P.; Dahlbeck, D.; Holub, E.; Staskawicz, B.J. NDR1, a pathogen-induced component required for Arabidopsis disease resistance. *Science* 1997, *278*, 1963–1965.
63. Takahashi, H.; Miller, J.; Nozaki, Y.; Takeda, M.; Shah, J.; Hase, S.; Ikegami, M.; Ehara, Y.; Dinesh-Kumar, S.P.; Sukamto, R.C.Y.1, an Arabidopsis thaliana RPP8/HRT family resistance gene, conferring resistance to cucumber mosaic virus requires salicylic acid, ethylene and a novel signal transduction mechanism. *Plant J.* 2002, *32*, 655–667.
64. Bashir, S.; Rehman, N.; Fakhar Zaman, F.; Naeem, M.K.; Jamal, A.; Tellier, A.; Ilyas, M.; Silva Arias, G.A.; Khan, M.R. Genome-wide characterization of the NLR gene family in tomato (*Solanum lycopersicum*) and their relatedness to disease resistance. *Front. Genet.* 2022, *13*, 931580.
65. Deslandes, L.; Olivier, J.; Theuclier, F.; Hirsch, J.; Feng, D.X.; Bittner-Eddy, P.; Beynon, J.; Marco, Y. Resistance to *Ralstonia solanacearum* in Arabidopsis thaliana is conferred by the recessive RRS1-R gene, a member of a novel family of resistance genes. *Proc. Natl. Acad. Sci. USA* 2002, *99*, 2404–2409.
66. Steuernagel, B.; Jupe, F.; Witek, K.; Jones, J.D.; Wulff, B.B. NLR-parser: Rapid annotation of plant NLR complements. *Bioinformatics* 2015, *31*, 1665–1667.
67. Huang, Z.; Qiao, F.; Yang, B.; Liu, J.; Liu, Y.; Wulff, B.B.H.; Hu, P.; Lv, Z.; Zhang, R.; Chen, P.; et al. Genome-wide identification of the NLR gene family in *Haynaldia villosa* by SMRT-RenSeq. *BMC Genom.* 2022, *23*, 118.
68. Toda, N.; Rustenholz, C.; Baud, A.; LePaslier, M.C.; Amselem, J.; Merdinoglu, D.; Faivre-Rampant, P. NLGenomeSweeper: A Tool for Genome-Wide NBS-LRR Resistance Gene Identification. *Genes* 2020, *11*, 333.
69. Jupe, F.; Witek, K.; Verweij, W.; Sliwka, J.; Pritchard, L.; Etherington, G.J.; Maclean, D.; Cock, P.J.; Leggett, R.M.; Bryan, G.J.; et al. Resistance gene enrichment sequencing (RenSeq) enables reannotation of the NB-LRR gene family from sequenced plant genomes and rapid mapping of resistance loci in segregating populations. *Plant J.* 2013, *76*, 530–544.
70. Kourelis, J.; Sakai, T.; Adachi, H.; Kamoun, S. RefPlantNLR: A comprehensive collection of experimentally validated plant NLRs. *PLoS Biol.* 2021, *19*, e3001124.
71. Armario Najera, V.; Twyman, R.M.; Christou, P.; Zhu, C. Applications of multiplex genome editing in higher plants. *Curr Opin Biotechnol.* 2019, *59*, 93–102.
72. Li, C.; Chu, W.; Gill, R.A.; Sang, S.; Shi, Y.; Hu, X.; Yang, Y.; Zaman, Q.U.; Zhang, B. Computational tools and resources for CRISPR/Cas genome editing. *Genom. Proteom. Bioinform.* 2022, *S1672-0229(22)00027-4*.

73. Fister, A.S.; Landherr, L.; Maximova, S.N.; Gultinan, M.J. Transient expression of CRISPR/Cas9 machinery targeting *TcNPR3* enhances defense response in *Theobromacacao*. *Front. Plant Sci.* 2018, *9*, 268.
74. Sun, Q.; Lin, L.; Liu, D.; Wu, D.; Fang, Y.; Wu, J.; Wang, Y. CRISPR/Cas9-Mediated multiplex genome editing of the BnWRKY11 and BnWRKY70 genes in *Brassica napus* L. *Int. J. Mol. Sci.* 2018, *19*, 2716.
75. Wang, Y.; Cheng, X.; Shan, Q.; Zhang, Y.; Liu, J.; Gao, C.; Qiu, J.L. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* 2014, *32*, 947–951.
76. Santillán Martínez, M.I.; Bracuto, V.; Koseoglou, E.; Appiano, M.; Jacobsen, E.; Visser, R.G.F.; Wolters, A.A.; Bai, Y. CRISPR/Cas9-targeted mutagenesis of the tomato susceptibility gene *PMR4* for resistance against powdery mildew. *BMC Plant Biol.* 2020, *20*, 284.
77. Jarosch, B.; Kogel, K.H.; Schaffrath, U. The ambivalence of the barley Mlo locus: Mutations conferring resistance against powdery mildew (*Blumeria graminis* f. sp. *hordei*) enhance susceptibility to the rice blast fungus *Magnaporthe grisea*. *Mol. Plant Microbe Interact.* 1999, *12*, 508–514.
78. Wan, D.Y.; Guo, Y.; Cheng, Y.; Hu, Y.; Xiao, S.; Wang, Y.; Wen, Y.Q. CRISPR/Cas9-mediated mutagenesis of VvMLO3 results in enhanced resistance to powdery mildew in grapevine (*Vitis vinifera*). *Hortic. Res.* 2020, *7*, 116.
79. Zhang, Y.; Bai, Y.; Wu, G.; Zou, S.; Chen, Y.; Gao, C.; Tang, D. Simultaneous modification of three homoeologs of TaEDR1 by genome editing enhances powdery mildew resistance in wheat. *Plant J.* 2017, *91*, 714–724.
80. Liu, D.; Chen, X.; Liu, J.; Ye, J.; Guo, Z. The rice ERF transcription factor *OsERF922* negatively regulates resistance to *Magnaporthe oryzae* and salt tolerance. *J. Exp. Bot.* 2012, *63*, 3899–3912.
81. Lu, W.; Deng, F.; Jia, J.; Chen, X.; Li, J.; Wen, Q.; Li, T.; Meng, Y.; Shan, W. The Arabidopsis thaliana gene AtERF019 negatively regulates plant resistance to *Phytophthora parasitica* by suppressing PAMP-triggered immunity. *Mol. Plant Pathol.* 2020, *21*, 1179–1193.
82. Langen, G.; von Einem, S.; Koch, A.; Imani, J.; Pai, S.B.; Manohar, M.; Ehlers, K.; Choi, H.W.; Claar, M.; Schmidt, R.; et al. The compromised recognition of turnip crinkler virus I subfamily of microorchidia ATPases regulates disease resistance in barley to biotrophic and necrotrophic pathogens. *Plant Physiol.* 2014, *164*, 866–878.
83. Kumar, N.; Galli, M.; Ordon, J.; Stuttmann, J.; Kogel, K.H.; Imani, J. Further analysis of barley MORC1 using a highly efficient RNA-guided Cas9 gene-editing system. *Plant Biotechnol. J.* 2018, *16*, 1892–1903.
84. Mazier, M.; Flamain, F.; Nicolai, M.; Sarnette, V.; Caranta, C. Knock-down of both the IF4E1 and IF4E2 genes confers broad-spectrum resistance against potyviruses in tomato. *PLoS ONE* 2011, *6*, e29595.
85. Rodríguez-Hernández, A.M.; Gosalvez, B.; Sempere, R.N.; Burgos, L.; Aranda, M.A.; Truniger, V. Melon RNA interference (RNAi) lines silenced for Cm-eIF4E show broad virus resistance. *Mol. Plant Pathol.* 2012, *13*, 755–763.
86. Ma, S.; Lapin, D.; Liu, L.; Sun, Y.; Song, W.; Zhang, X.; Logemann, E.; Yu, D.; Wang, J.; Jirschitzka, J.; et al. Direct pathogen-induced assembly of an NLR immunoreceptor complex to form a holoenzyme. *Science* 2020, *370*, eabe3069.
87. Martin, R.; Qi, T.; Zhang, H.; Liu, F.; King, M.; Toth, C.; Nogales, E.; Staskawicz, B.J. Structure of the activated ROQ1 resistosome directly recognizing the pathogen effector XopQ. *Science* 2020, *370*, eabd9993.
88. Li, S.; Shen, L.; Hu, P.; Liu, Q.; Zhu, X.; Qian, Q.; Wang, K.; Wang, Y. Developing disease-resistant thermosensitive male sterility rice by multiplex gene editing. *J. Integr. Plant Biol.* 2019, *61*, 1201–1205.
89. Peng, A.; Chen, S.; Lei, T.; Xu, L.; He, Y.; Wu, L.; Yao, L.; Zou, X. Engineering canker-resistant plants through CRISPR/Cas9-targeted editing of the susceptibility gene *CsLOB1* promoter in citrus. *Plant Biotechnol. J.* 2017, *15*, 1509–1519.
90. Ahn, H.K.; Lin, X.; Olave-Achury, A.C.; Derevnina, L.; Contreras, M.P.; Kourelis, J.; Wu, C.H.; Kamoun, S.; Jones, J.D.G. Effector-dependent activation and oligomerization of plant NRC class helper NLRs by sensor NLR immune receptors Rpi-amr3 and Rpi-amr1. *EMBO J.* 2023, *42*, e111484.
91. Joshi, A.; Yang, S.Y.; Lee, J.H. Integrated Molecular and Bioinformatics Approaches for Disease-Related Genes in Plants. *Plants*. 2023, *12*, 245.