

Micro Nutrient Management for Enhanced Growth, Yield, and Quality of Mango: A Comprehensive Review

ABSTRACT

This review paper delves into the intricate interrelationships among micronutrient interventions, carbohydrates, and osmostimulants, investigating their collective influence on diverse aspects of mango tree growth and fruit characteristics. Micronutrients, essential elements required in small quantities for plant health, play a pivotal role in shaping optimal plant development. The paper underscores the significance of these micronutrients by highlighting their roles in stimulating vegetative growth, influencing flowering dynamics, regulating fruit set processes, impacting yield metrics, and contributing to the nuanced attributes of fruit quality. This synthesis illuminates the complex network of processes governing mango tree growth and fruit production. Moreover, the study underscores the potential of well-structured nutrient management strategies as key catalysts for optimizing mango production outcomes. By carefully balancing the imperative of yield quantity and fruit quality, such strategies hold the promise of enhancing overall agricultural efficiency. Drawing insights from a comprehensive review of pertinent studies, this research paper contributes to a holistic comprehension of the intricate interactions involving micronutrients, carbohydrates, and osmostimulants, intricately connected with the diverse growth and fruit parameters within mango trees. Through this endeavor, the paper provides valuable insights into the intricate interplay of micronutrients within mango cultivation, offering a knowledge foundation for informed agricultural practices geared towards sustainable and high-quality fruit production. The paramount importance of micronutrients in fostering robust crop health and productivity underscores the broader implications of these findings for contemporary agricultural methodologies.

Introduction

Mango (*Mangifera indica L.*) is a premier fruit crop of India considering its area, production and popularity among people is designated as “national fruit of India. Mango is considered the king of fruits in many countries (Purseglove, 1972)[1]. In mango many problems

are associated with fruit set, yield and quality due to an imbalance supply of nutrients and its results in poor health of plants that are also prone to attacks of insect-pests and diseases.

In spite of adequate flowering, low fruit yield in mango orchards has been experienced because of low initial fruit set and subsequently higher fruitlet abscission Singh and Singh, (1995)[2]. Fruitlet abscission is a very complex physiological process, occurs in many cultivars of mango and at all stages of development, but it is particularly high during the first 3-4 weeks after pollination and accounts for over 90 % loss of set fruitlets Bains, et al., (1997)[3] and Wahdan and Melouk (2004)[4].

The use of growth substances and some chemical compounds may regulate fruit sets in many fruit crops. Many investigators found that spraying mango trees with Naphthalene acetic acid (NAA) at different concentrations (20, 25 and 40 ppm) increased fruit set percentages and fruit retention (Oksher et al., 1980[5] and Singh and Ram, 1983[6]). Auxin is well known as inhibitors of ethylene action in a number of plants (Beyer, 1976[7]). Moti Singh et al., (1987) [8] with Langra and Dashehari cvs stated that NAA or GA₃ each 5-25 ppm once sprayed at full bloom or twice at full bloom and at pea stage or thrice plus at marble stage increased fruit retention. Singh et al. (1991)[9] found that the highest fruit retention and yield/tree were recorded on mango cv. (Amrapali) by spraying urea with 3 % at pea stage.

In plants, micronutrients are required for different physiological and metabolic processes, and their deficiency affects a number of processes including hindered plant growth, productivity, and quality (Berdanier and Berdanier 2015[10]; Gurjar et al. 2015[11]; Souri and Aslani 2018[12]; Souri and Bakhtiarizade 2019[13]). The key role of micronutrients is in respiration and photosynthesis (Ahmed et al. 2009)[14]. Iron deficiency affected the yield, chlorophyll contents, fruit quality, and mineral nutrients in a number of fruit trees (Tagliavini et al. 2000[15]; Souri et al. 2018[16]). Iron is an essential micronutrient for plant growth and development. It plays a critical role in various physiological processes, including photosynthesis, respiration, and DNA synthesis. Iron is a component of many enzymes involved in redox reactions and electron transfer within the plant cells. (Sharma and Dietz 2006[17]). Zinc (Zn) stands as a pivotal micronutrient for plants, with its significance underscored by its indispensable roles in crucial aspects such as fruit set, retention, yield, and quality [Marschner 2011[18]]. The absence of adequate zinc can profoundly hinder plant growth, leading to a substantial decrease in overall

yield (Ojeda-Barrios et al. 2014)[19]. This emphasizes the critical nature of maintaining proper zinc levels for optimizing agricultural productivity and ensuring robust plant development. Zinc is a crucial micronutrient required for the synthesis of enzymes and proteins in plants. It is involved in various enzymatic reactions, including those related to carbohydrate metabolism, auxin synthesis, and protein synthesis. Zinc also plays a role in DNA and RNA synthesis and is essential for proper root development. (Broadley et al. 2007[20], Yruela 2005[21]). Boron is essential for cell wall formation and stability in plants. It plays a critical role in the synthesis of nucleic acids and proteins, as well as in the transport of sugars and nutrients within the plant. Boron is also involved in pollen tube growth and is essential for proper seed and fruit development (Goldbach et al. 2001[22], Gupta and Gupta 1997[23]).

Effect of Micronutrients on Various Parameters in Mango Trees

Microscopic nutrients, also known as micronutrients, play a crucial role in influencing various parameters of mango tree growth and development. Here, we explore the diverse effects of these nutrients on different aspects of mango trees' lifecycle:

1. Vegetative Growth Enhancement:

Micronutrients play a vital role in the vegetative growth of mango (*Mangifera indica*), contributing to the overall health, vigor, and development of this economically important fruit tree (Vittal 2023[24] and Das 2019[25]). Mango trees require a diverse range of nutrients in trace amounts to support their various physiological processes. While macronutrients like nitrogen, phosphorus, and potassium are well-recognized for their influence on plant growth, micronutrients, including elements like iron, zinc, copper, manganese, boron, and molybdenum, are equally essential for optimal growth and development. (Ryan 2013[26] and Milošević 2023[27]). The enhancement of vegetative growth in plants has been a subject of extensive research. Notably, Singh and Rajput (1976)[28] conducted a study on mango trees, revealing the positive effects of varying levels of $ZnSO_4$ on vegetative growth parameters. The application of zinc and boron led to increased terminal shoot length, leaf count, and leaf area per shoot, thereby influencing plant height, trunk girth, and young plant spread. In a similar vein, Masroor et al. (2016)[29] highlighted the significance of zinc application, particularly a foliar spray

of 1% ZnSO₄, in elevating zinc levels within mango leaves. This innovative approach demonstrated the potential of targeted micronutrient application for stimulating vegetative growth.

Moreover, the role of nano zinc oxide (nZnO) in influencing vegetative growth was investigated by Elsheery et al. (2020)[30], who observed a significant increase in leaf area with the application of 100 units of nZnO. This study emphasized the efficacy of nano-based formulations in enhancing plant growth. Additionally, Kacha HL (2021)[31] shed light on zinc's crucial involvement in starch formation, a process that subsequently contributes to increased plant weight and overall vegetative growth. These research findings collectively underscore the multifaceted contributions of zinc and its compounds to fostering robust vegetative growth in various plant species.

The empirical data from these studies collectively reinforces the vital significance of micronutrients in promoting robust vegetative growth in mango trees. These elements intricately influence key physiological processes that drive growth, ultimately impacting shoot length, leaf area, stem strength, and canopy density. The application of appropriate micronutrient management strategies, as demonstrated by these findings, holds immense potential for optimizing mango orchard health and productivity. Continued research in this field is imperative to refine nutrient management practices and further enhance the yield and quality of mango crops.

2. Leaf Mineral Contents and Zn Foliar Spray:

Numerous studies have elucidated the specific effects of micronutrient supplementation on the mineral content of mango leaves (Ahamd 2018)[32]. This not only improved the overall micronutrient balance but also positively influenced the content of other minerals, leading to a more comprehensive nutrient profile within the leaves (Razzaq 2013)[33]. Leaf mineral content and its association with Zn foliar spray have garnered significant attention in recent research endeavors. Elsheery et al. (2020)[30] extended their investigations to delve into the effects of Zn foliar spray on leaf mineral content within mango trees. Their findings illuminated the positive impact of this application method on enriching the mineral content of leaves, thereby contributing to enhanced overall plant health. Similarly, Ahmad I et al. (2018)[32] underlined the

considerable influence of Zn foliar spray on leaf mineral content. This relationship between Zn application and leaf mineral composition holds promising implications for influencing not only mineral concentrations but also potentially exerting downstream effects on various vegetative parameters. The convergence of these research insights underscores the significance of Zn foliar spray as a viable strategy to manipulate leaf mineral content and subsequently impact plant vitality.

In essence, the pivotal role of zinc (Zn) as an essential trace element in plants extends beyond enzymatic reactions, encompassing the vital regulation of protein and carbohydrate metabolism. This intricate interplay underscores its indispensability for optimal growth and development. Striking a delicate balance in zinc levels is crucial, as deficiency or excess can both disrupt plant health. Recognizing the multifaceted influence of zinc not only enhances our comprehension of plant physiology but also holds the potential to revolutionize agricultural practices for improved crop yields and sustainability. Swietlik (1999)[34]

3. **Flowering and Fruit Set:**

Micronutrients exert a profound impact on the flowering and fruit set processes of mango (*Mangifera indica*) trees, playing a crucial role in determining the yield and quality of this economically significant fruit crop (Suman 2017)[35]. While macronutrients like nitrogen, phosphorus, and potassium are widely recognized for their roles in plant growth, the contribution of micronutrients—such as iron, zinc, copper, manganese, boron, and molybdenum—cannot be understated when it comes to fostering optimal flowering and fruit set.

The intricate relationship between micronutrients and flowering processes in plants has been a subject of focused investigation. Elsheery et al. (2020)[30] conducted a study showcasing the potential benefits of applying ZnO and Si NPs via spraying on mango trees. This approach notably led to a reduction in floral malformation, consequently promoting an improved flowering process. This finding emphasizes the role of micronutrient supplementation in optimizing flower development.

Furthermore, the pivotal role of zinc in influencing flowering was illuminated by Masroor et al. (2016)[29] and Kacha HL et al. (2021)[21]. Through their independent

studies, both researchers emphasized the substantial impact of ZnSO₄ application on enhancing the number of flowers per panicle. These observations collectively underscore the vital contribution of micronutrients, particularly zinc-based applications, in modulating flowering outcomes in plants. The convergence of research findings highlights the potential for targeted nutrient management strategies to positively influence flowering and subsequent fruit sets, thereby enhancing overall crop productivity.

4. **Fruit Retention and Weight:**

Empirical investigations provide a rich understanding of the specific effects of micronutrient applications on mango fruit retention and weight. Notably, research by Merwad et al. (2016)[36] spotlighted NAA, zinc, boron and calcium role in fortifying fruit retention. Mango trees treated with NAA, zinc, boron and calcium demonstrated an elevated proportion of retained fruits, underscoring the critical contribution of adequate NAA, zinc, boron and calcium levels in bolstering fruit-holding resilience.

The intricate interplay between micronutrients and fruit development has attracted considerable research attention. Ahmad et al. (2018)[32] conducted a study that yielded noteworthy outcomes in terms of high fruit retention percentages, employing specific micronutrient treatments. This finding suggests a potential role for micronutrients in enhancing the retention of fruits on trees.

Furthermore, the influence of zinc on fruit weight was demonstrated by Masroor et al. (2016)[29] and Kacha HL et al. (2021)[31]. Their independent research endeavors elucidated how zinc application strategies, whether foliar or soil-based, could positively impact fruit weight. These studies collectively underscore the significant potential of targeted zinc treatments to influence fruit development and quality.

Expanding on this theme, Dheware (2020)[37] delved into a comprehensive nutrient application strategy involving both soil and foliar applications. This holistic approach yielded substantial improvements in fruit weight and quality. The convergence of these research findings underscores the multifaceted influence of micronutrients, particularly zinc, in shaping the retention and weight of fruits. This insight holds considerable implications for the agricultural sector, suggesting avenues for optimizing

crop yield and enhancing fruit characteristics through strategic nutrient management practices

5. Total Soluble Solids, Sugar Content, and Vitamin C:

The nexus between micronutrient applications and the biochemical composition of fruits has been a subject of keen exploration. RM Dheware (2020)[37] provided insights into the influence of specific nutrient applications on fruit quality, unveiling an increase in Total Soluble Solids (TSS) content. This finding underscores the potential for targeted nutrient management to impact the composition of fruits.

In line with this, Patel (2018)[38] undertook research into novel organic liquid fertilizer treatments, which yielded multifaceted improvements in fruit quality. Notably, these treatments led to elevated TSS levels, along with increased sugar and vitamin C content. This discovery highlights the efficacy of innovative fertilization approaches in enhancing the nutritional attributes of fruits.

Collectively, the research of Dheware 2020[37] and Patel 2018[38] reinforces the interconnectedness between micronutrient applications and key components of fruit quality, including TSS, sugar content, and vitamin C levels. These findings not only contribute to our understanding of nutrient-induced variations in fruit composition but also hold implications for promoting healthier and more nutrient-rich produce for consumers.

6. Effects on Carbohydrates and Osmostimulants:

According to Siddiki et al. (2022)[39] Micronutrients are essential for various enzymatic reactions involved in photosynthesis, which is the process by which plants produce carbohydrates (such as sugars) from carbon dioxide and water in the presence of sunlight. While micronutrients themselves do not directly form carbohydrates, their presence or absence can affect the efficiency of photosynthesis and subsequently impact carbohydrate production.

Osmostimulants are compounds that help regulate osmotic balance in plants, which is critical for water and nutrient uptake Elsheery et al. (2020)[30]. While micronutrients do not directly influence osmostimulants, their impact on overall plant health can indirectly affect osmotic regulation. The impact of micronutrient applications

on carbohydrates and osmostimulants within plants has been a topic of interest and investigation. Elsheery et al. (2020)[30] conducted research that yielded insightful results, indicating an improved total yield resulting from the application of Zn and Si NPs. This finding highlights the positive effect of osmostimulants in promoting overall plant productivity.

Kacha et al. (2021)[31] contributed to this understanding by demonstrating how fruit yield per tree could be enhanced through soil application of Grade-V 400g, suggesting the influence of specific nutrient strategies on carbohydrate accumulation and subsequent yield.

Dheware (2020)[37] further enriched this discourse by presenting evidence of increased fruit yield attributed to comprehensive micronutrient treatments, including the application of copper and Borax. These findings underscore the role of specific nutrients in influencing carbohydrate metabolism and, consequently, fruit production.

The research of Puranik (2019)[41] supplements this narrative by reporting increased fruit yield resulting from specific nutrient treatments involving copper, boron, and molybdenum application. This observation lends credence to the notion that the manipulation of osmostimulants and carbohydrates through targeted micronutrient interventions can significantly impact plant productivity and yield. The collective findings of these studies contribute to a more comprehensive understanding of the intricate relationship between micronutrients, carbohydrates, and osmostimulants in influencing crop yield and overall plant performance.

7. **Yield Enhancement:**

Recent studies have advanced our understanding of how micronutrient applications can positively influence crop yield, shedding light on diverse strategies that can significantly enhance overall plant productivity. Elsheery et al. (2020)[30] demonstrated the potential of nanomaterials by applying zinc (Zn) and silicon (Si) nanoparticles (NPs) to mango trees, resulting in a noteworthy improvement in total yield. This finding underscores the capacity of nanomaterials to exert a nanoscale impact on crop productivity. Similarly, Kacha HL et al. (2021)[31] contributed to this discourse by highlighting the efficacy of soil interventions, showcasing that the application of Grade-V

400g significantly elevated fruit yield per tree. This approach accentuates the importance of nurturing the soil environment for substantial enhancements in crop productivity.

Furthermore, Dheware (2020)[37] explored a holistic treatment strategy involving foliar spray of micronutrients such as zinc sulphate, copper sulphate, and Borax during critical growth stages. This meticulously timed application led to a higher number of fruits per tree, addressing issues related to fruit drop and concurrently elevating fruit weight. The cumulative effect resulted in a remarkable surge in overall fruit yield. Expanding on this theme, Puranik (2019)[41] reported on the successful augmentation of fruit yield through a specific nutrient regimen encompassing copper, boron, and molybdenum application. The combined impact of these essential elements demonstrated their potential to collectively contribute to improved yield outcomes.

In conclusion, these studies collectively underscore the multifaceted avenues through which micronutrient interventions can substantially enhance crop yield. Whether through nanomaterials, soil amendments, or comprehensive treatments, these findings provide valuable insights for optimizing agricultural practices and ensuring sustainable food production in the face of evolving agricultural challenges.

8. **Fruit Quality Improvement:**

Micronutrients play a significant role in enhancing fruit quality in mango trees. While they are required in smaller quantities compared to macronutrients, their presence is essential for various biochemical and physiological processes that contribute to fruit development, flavor, color, texture, and overall quality Barrett et al. (2010)[42]. Here's how different micronutrients can impact fruit quality improvement in mangoes.

- **Iron (Fe):**

Iron is essential for the synthesis of chlorophyll, which is responsible for the green color of leaves and indirectly influences fruit color development Sun (2022)[43]. An adequate supply of iron ensures that photosynthesis is efficient, leading to the production of carbohydrates that contribute to fruit sweetness and flavor Ribeiro and Schieber (2010)[44].

- **Zinc (Zn):**

Zinc is involved in the synthesis of auxins, which are plant hormones that play a role in fruit development and ripening Kazemi (2014)[45]. Adequate zinc levels can

promote proper fruit sizing, uniform ripening, and enhanced flavor development Masroor et al. (2014)[46].

- **Boron (B):**

Boron is crucial for cell wall formation and stability Liu et al. (2017)[47]. It contributes to the texture of the fruit by influencing cell wall strength Huang (2023)[48]. Proper boron levels can improve fruit firmness and minimize disorders like fruit cracking Shivshankar (2014)[49].

- **Manganese (Mn):**

Manganese is a cofactor for various enzymes involved in the synthesis of secondary metabolites, including those responsible for flavor and aroma compounds in fruits Schwab (2003)[50]. Adequate manganese levels can contribute to improved taste and aroma of mangoes Appiah et al. (2011)[51].

- **Copper (Cu):**

Copper is involved in several enzymatic reactions, including those related to lignin synthesis and color development Lin et al. (2005) [52]. It can influence the coloration of fruit skin and contribute to the visual appeal of mangoes Lebaka et al. (2021) [53].

- **Molybdenum (Mo):**

Molybdenum is necessary for nitrogen fixation and the conversion of nitrates to amino acids, which are building blocks of proteins and flavor compounds. Optimal molybdenum levels can lead to improved protein content and enhanced flavor in mangoes Gericó et al. (2020) [54].

- **Magnesium (Mg):**

Magnesium is involved in the synthesis of chlorophyll and enzymes required for carbohydrate metabolism Moretti et al. (2021) [55]. It indirectly affects fruit color development and sweetness by ensuring efficient photosynthesis and sugar production Nagereza and Pawelzik (2016) [56].

- **Calcium (Ca):**

Calcium plays a critical role in cell wall structure and integrity Khaliq et al. (2016)[57]. Adequate calcium levels can prevent disorders like bitter pit, which affects

fruit texture and quality Sharma et al. (2013)[58]. Calcium also helps maintain fruit firmness during storage and transportation Wang et al. (2014)[59].

Furthermore, Ahmad I et al. (2018)[32] illuminated the manifold benefits of zinc applications by emphasizing how both foliar and soil treatments led to significant enhancements in fruit yield and quality. Parameters such as fruit weight, volume, pulp recovery, Total Soluble Solids (TSS), and acidity witnessed notable improvements. This holistic improvement underscores the pivotal role of zinc in shaping fruit attributes.

Elsheery et al. (2020)[30] delved into the intricacies of micronutrient combinations, revealing that the synergy between 50 mg/L of nano zinc oxide (nZnO) and 300 mg/L of nano silicon (nSi) resulted in the highest acidity in mangoes. Interestingly, this outcome could be attributed to the inhibitory effects of a high concentration of nSi on acidity levels. This insight provides a nuanced perspective on how different micronutrients' interactions can influence specific fruit attributes.

Furthermore, Patel et al. (2018)[38] conducted research that showcased the diverse impact of foliar applications of 1% Grade-IV micronutrients. This approach led to an overall acceptability of the produce. This enhancement in fruit quality can be attributed to iron's contribution to flavor-protein development, the presence of adequate zinc improving auxin content, and zinc's catalytic role in oxidation-reduction processes within plants. These findings collectively underscore the intricate interplay between micronutrients and fruit quality, shedding light on the diverse pathways through which these elements influence the sensory and nutritional attributes of produce.

Conclusion

In summary, the influence of micronutrients on various parameters in mango trees emerges as a multifaceted process. From promoting vegetative growth to affecting flowering, fruit set, and fruit quality, these findings underscore the indispensable role that micronutrients play in fine-tuning mango tree development. The combined influence of micronutrients, carbohydrates, and osmostimulants on mango tree growth and fruit parameters is evident across various studies. From enhancing vegetative growth to affecting flowering, fruit set, yield, and quality, these factors collectively shape mango cultivation practices. Proper nutrient management strategies hold promise for optimizing mango production, ensuring both quantity and quality in harvests.

The review underscores the importance of micronutrients in influencing mango tree yield and fruit quality. The findings highlight the multifaceted impact of micronutrient applications, demonstrating their role in enhancing yield through fruit retention, increasing fruit numbers, and influencing fruit weight. Furthermore, the improvement in fruit quality, as evidenced by parameters like acidity and overall acceptability, emphasizes the potential of micronutrients to contribute to the overall success of mango cultivation. Collectively, these insights underline the intricate web of interactions between micronutrients, carbohydrates, osmostimulants, and various growth and fruit parameters in mango trees. These findings offer valuable insights for optimizing nutrient management strategies and practices, ensuring sustainable and high-quality mango production.

UNDER PEER REVIEW

References

1. Purseglove, J. W. (1972). Mangoes west of India. *Acta. Hort.* 24: 170-174.
2. Singh, Z. and, Singh, L. (1995). Increased fruit set and retention in mango with exogenous applications of polyamines. *J. Hort. Sci.*, 70 (2): 271-277.
3. Bains, K. S., Bajwa, G. S. and Singh, Z. (1997). Abscission of mango fruitlets. I. In relation to endogenous concentrations of IAA, GA and ABA in pedicels and fruitlets. *Fruits*, 52: 159- 165.
4. Wahdan, M. T. and Melouk, A. E. (2004). Effect of Amcotone on vegetative growth, fruiting, fruit yield and quality of Succary Abiad mango trees. *Agri. Res. J. Suez Canal University.* 4(2): 69-76.
5. Oksher, A. K., Ramachandran, C. and Pyhodath, J. S. (1980). Effect of planofix on fruit set in mango. *Agric. Res. J. Kereale.* 17(1): 105 (Hort. Abst. 50, 5712).
6. Singh, R. S. and Ram, S. (1983). Studies on the use of plant growth substances for fruit retention in mango cv. Dashehari. *Ind. J. Hort.* 40(3/4): 188.
7. Beyer, E. M. J. (1976). A potent inhibitor of ethylene action in plants. *Plant physiol.* 58, 268- 271
8. Moti-Singh, A. S., Chaudhary, A. S. and Prasad, M. (1987). A note on the effect of some plant regulators on fruit retention in mango (*Mangifera indica* L). *Haryana J. Hort. Sci.* 15 (3/4): 221 (Hort. Abst. 51 (3):2200).
9. Singh, J. N., Rajput, C. B. S. and Prakash, S. (1991). Effect of urea spray on fruit retention and physiochemical composition of mango (*Mangifera indica* L) cv. Amrapali. *Haryana J. Hort. Sci.* 20(1/2): 35. (Hort. Abs. 62(4): 3500).
10. Berdanier CD, Berdanier LA. *Advanced nutrition: macronutrients, micronutrients, and metabolism.* CRC Press; 2015.
11. Gurjar TD, Patel NL, Panchal BH, Chaudhari D. Effect of foliar spray of micronutrients on flowering and fruiting of Alphonso mango (*Mangifera indica* L.). *Bioscan.* 2015;10(3):1053–1056.
12. Souri MK, Aslani M. Beneficial effects of foliar application of organic chelate fertilizers on French bean production under field conditions in a calcareous soil. *Adv Horticulture Sci.* 2018;32(2):265–272.

13. Souri MK, Bakhtiarizade M. Biostimulation effects of Rosemary essential oil on growth and nutrient uptake of tomato seedlings. *Sci Hortic.* 2019;243:472–476.
14. Ahmed N, Ahmad F, Abid M, Ullah MA. Impact of zinc fertilization on gas exchange characteristics and water use efficiency of cotton crop under arid environment. *Pak J Bot.* 2009;41(5):2189–97.
15. Tagliavini M, Abadía J, Rombolà AD, Abadía A, Tsipouridis C, Marangoni B. Agronomic means for the control of iron deficiency chlorosis in deciduous fruit trees. *J Plant Nutri.* 2000;23:2007–2022.
16. Souri MK, Naiji M, Aslani M. Effect of Fe-glycine aminochelate on pod quality and iron concentrations of bean (*Phaseolus vulgaris* L.) under lime soil conditions. *Commun Soil Sci Plant Anal.* 2018;49(2):215–224.
17. Sharma, S. S., & Dietz, K. J. (2006). The significance of amino acids and amino acid-derived molecules in plant responses and adaptation to heavy metal stress. *Journal of Experimental Botany*, 57(4), 711–726.
18. Marschner, H. *Mineral Nutrition of Higher Plants*, 3rd ed.; Academic Press: Cambridge, MA, USA, 2011; pp. 347–364.
19. Ojeda-Barrios, D.L.; Perea-Portillo, E.; Hernández-Rodríguez, O.A.; Martínez-Téllez, J.; Abadía, J.; Lombardini, L. Foliar fertilization with zinc in pecan trees. *HortScience* 2014, 49, 562–566.
20. Broadley, M. R., White, P. J., Hammond, J. P., Zelko, I., & Lux, A. (2007). Zinc in plants. *New Phytologist*, 173(4), 677–702.
21. Yruela, I. (2005). Copper in plants. *Brazilian Journal of Plant Physiology*, 17(1), 145–156.
22. Goldbach, H. E., Wimmer, M. A., & Findelee, P. (2001). Boron in plants and animals: Is there a role beyond cell-wall structure? *Journal of Plant Nutrition and Soil Science*, 164(4), 415–419.
23. Gupta, U. C., & Gupta, S. C. (1997). *Boron: Its role in crop production*. CRC Press.
24. Vittal, H., Sharma, N., Dubey, A. K., Shivran, M., Singh, S. K., Meena, M. C., ... & Sharma, R. M. (2023). Rootstock-mediated carbohydrate metabolism, nutrient

contents, and physiological modifications in regular and alternate mango (*Mangifera indica* L.) scion varieties. *Plos one*, 18(5), e0284910.

25. Das, S. C., Datta, M., Ray, P., Singh, S. K., Jena, R. K., Das, B., & Ray, S. K. (2019). Mango (*Mangifera indica*) Cultivation in North-Eastern Region of India. *Adv. Agric. Res. Technol. J*, 3, 54-66.
26. Ryan, J., Rashid, A., Torrent, J., Yau, S. K., Ibrikci, H., Sommer, R., & Erenoglu, E. B. (2013). Micronutrient constraints to crop production in the Middle East–West Asia region: significance, research, and management. *Advances in Agronomy*, 122, 1-84.
27. Milošević, T., & Milošević, N. (2023). Role of Nutrients in Modifications of Fruit Quality and Antioxidant Activity. *Plant Ionomics: Sensing, Signaling, and Regulation*, 148-170.
28. Singh RR, Rajput CBS (1976). Effect of various concentrations of zinc on vegetative growth characters, flowering, fruiting and physico-chemical composition of fruits in mango cv. Chausa, Haryana J. Hortic. Sci. 5(1-2):10-14.
29. Masroor, H. M., Anjum, M. A., Hussain, S., Ejaz, S., Ahmad, S., Ercisli, S., & Zia-Ul-Haq, M. (2016). Zinc Ameliorates Fruit Yield and Quality of Mangoes Cultivated in Calcareous Soils. *Erwerbs-Obstbau*, 58(1).
30. Elsheery, N. I., Helaly, M. N., El-Hoseiny, H. M., & Alam-Eldein, S. M. (2020). Zinc oxide and silicone nanoparticles to improve the resistance mechanism and annual productivity of salt-stressed mango trees. *Agronomy*, 10(4), 558.
31. KACHA, H. L., Patel, H. C., & Paradava, D. R. (2021). Effect of soil and foliar applications of micronutrients on flowering and yield of mango: Performance of mango influenced by soil and foliar applications of micronutrients. *Journal of AgriSearch*, 8(1), 40-44.
32. Ahmad, I., Bibi, F., Ullah, H., & Munir, T. M. (2018). Mango fruit yield and critical quality parameters respond to foliar and soil applications of zinc and boron. *Plants*, 7(4), 97.
33. Razzaq, K., Khan, A. S., Malik, A. U., Shahid, M., & Ullah, S. (2013). Foliar application of zinc influences the leaf mineral status, vegetative and reproductive

growth, yield and fruit quality of 'Kinnow' mandarin. *Journal of plant nutrition*, 36(10), 1479-1495.

34. Swietlik, D. Zinc nutrition in horticultural crops. *Hort. Rev.* 1999, 23, 109–180.
35. Suman, M., Sangma, P. D., & Singh, D. (2017). Role of micronutrients (Fe, Zn, B, Cu, Mg, Mn and Mo) in fruit crops. *Int. J. Curr. Microbiol. App. Sci*, 6(6), 3240-3250.
36. Merwad, M. A., Eisa, R. A., & Saleh, M. M. S. (2016). The beneficial effect of NAA, Zn, Ca and B on fruiting, yield and fruit quality of Alphonso mango trees. *Intl. J. Chemtech Res*, 9(3), 147-157.
37. Dheware, R. M., Nalage, N. A., Sawant, B. N., Haldavanekar, P. C., Raut, R. A., Munj, A. Y., & Sawant, S. N. (2020). Response of micronutrient on the quality yield of mango cv. Alphonso under Konkan agro-climatic conditions. *Journal of Pharmacognosy and Phytochemistry*, 9(3), 2004-2006.
38. Patel RJ, Patil SJ, Tandel BM, Ahlawat TR and Amarcholi JJ 2018 Effect of micronutrients and banana pseudostem sap at different pH levels of foliar spray solution on fruit quality of mango (*Mangifera indica* L.) cv. Kesar *International Journal of Chemical Studies* 2018; 6(3): 852-854
39. Siddiki, S. Y. A., Mofijur, M., Kumar, P. S., Ahmed, S. F., Inayat, A., Kusumo, F., and Mahlia, T. M. I. (2022). Microalgae biomass as a sustainable source for biofuel, biochemical and biobased value-added products: An integrated biorefinery concept. *Fuel*, 307, 121782.
40. Elsheery, N.I., B. Wilske, J.-L. Zhang, and K.-F. Cao. 2007. Seasonal variations in gas exchange and chlorophyll fluorescence in the leaves of five mango cultivars in southern Yunnan, China. *J. Hort. Sci. Biotechnol.* 82:855–862.
41. Puranik, U. Y., Salvi, V. G., Prabhudesai, S. S., Salvi, B. R., & Dhekale, J. S. (2019). Effect of micronutrient fortified Konkan Annapurna Briquettes on yield of Alphonso mango (*Mangifera indica* L.) in lateritic soils of Konkan. *Journal of Pharmacognosy and Phytochemistry*, 8(2S), 273-276.
42. Barrett, D. M., Beaulieu, J. C., & Shewfelt, R. (2010). Color, flavor, texture, and nutritional quality of fresh-cut fruits and vegetables: desirable levels, instrumental

and sensory measurement, and the effects of processing. *Critical reviews in food science and nutrition*, 50(5), 369-389.

43. Sun, T., Rao, S., Zhou, X., & Li, L. (2022). Plant carotenoids: Recent advances and future perspectives. *Molecular Horticulture*, 2(1), 3.
44. Ribeiro, S. M. R., & Schieber, A. (2010). Bioactive compounds in mango (*Mangifera indica* L.). In *Bioactive foods in promoting health* (pp. 507-523). Academic Press.
45. Kazemi, M. (2014). Influence of foliar application of iron, calcium and zinc sulfate on vegetative growth and reproductive characteristics of strawberry cv. 'Pajaro'. *Trakia Journal of Sciences*, 12(1), 21-26.
46. Masroor, H. M., Anjum, M. A., Hussain, S., Ejaz, S., Ahmad, S., Ercisli, S., & Zia-Ul-Haq, M. (2016). Zinc Ameliorates Fruit Yield and Quality of Mangoes Cultivated in Calcareous Soils. *Erwerbs-Obstbau*, 58(1).
47. Liu, Y., Riaz, M., Yan, L., Zeng, Y., & Cuncang, J. (2019). Boron and calcium deficiency disturbing the growth of trifoliolate rootstock seedlings (*Poncirus trifoliolate* L.) by changing root architecture and cell wall. *Plant Physiology and Biochemistry*, 144, 345-354.
48. Huang, W., Shi, Y., Yan, H., Wang, H., Wu, D., Grierson, D., & Chen, K. (2023). The calcium-mediated homogalacturonan pectin complexation in cell walls contributes the firmness increase in loquat fruit during postharvest storage. *Journal of Advanced Research*, 49, 47-62.
49. Shivashankar, S. (2014). Physiological disorders of mango fruit. *Horticultural Reviews: Volume 42*, 313-348.
50. Schwab, W. (2003). Metabolome diversity: too few genes, too many metabolites?. *Phytochemistry*, 62(6), 837-849.
51. Appiah, F., Kumah, P., & Idun, I. (2011). Effect of ripening stage on composition, sensory qualities and acceptability of keitt mango (*Mangifera indica* L.) chips. *African Journal of Food, Agriculture, Nutrition and Development*, 11(5), 5096-5109.
52. Lin, C. C., Chen, L. M., & Liu, Z. H. (2005). Rapid effect of copper on lignin biosynthesis in soybean roots. *Plant Science*, 168(3), 855-861.

53. Lebaka, V. R., Wee, Y. J., Ye, W., & Korivi, M. (2021). Nutritional composition and bioactive compounds in three different parts of mango fruit. *International Journal of Environmental Research and Public Health*, 18(2), 741.
54. Gericó, T. G., Tavanti, R. F. R., de Lima, J. P., Ribeiro, R. P., dos Santos, L. C. C., da Silva, M. S., & dos Reis, A. R. (2020). Cobalt and molybdenum stimulate compounds of primary metabolism, nitrogen forms, and photosynthetic pigments in peanut plants (*Arachis hypogaea* L.). *Journal of Plant Nutrition*, 43(12), 1907-1922.
55. Moretti, L. G., Crusciol, C. A. C., Bossolani, J. W., Garcia, A., Rossi, R., & Moreira, A. (2021). Thermomagnesium as a fertilizer for soybean: Carbohydrate metabolism, silicon–magnesium fertilizer, and grain yield. *Journal of Plant Nutrition*, 44(14), 2108-2122.
56. Ngereza, J. A., & Pawelzik, E. (2016). Nutritional characterization of organically and conventionally grown mango (*Mangifera indica* L.) and pineapple (*Ananas cosmus*) of different origins. *J. Crop Sci. Agron*, 1(1), 1-17.
57. Khaliq, G., Mohamed, M. T. M., Ghazali, H. M., Ding, P., & Ali, A. (2016). Influence of gum arabic coating enriched with calcium chloride on physiological, biochemical and quality responses of mango (*Mangifera indica* L.) fruit stored under low temperature stress. *Postharvest Biology and Technology*, 111, 362-369.
58. Sharma, R. R., Singh, D., & Pal, R. K. (2013). Synergistic influence of pre-harvest calcium sprays and postharvest hot water treatment on fruit firmness, decay, bitter pit incidence and postharvest quality of royal delicious apples (*Malus x domestica* Borkh).
59. Wang, Y., Xie, X., & Long, L. E. (2014). The effect of postharvest calcium application in hydro-cooling water on tissue calcium content, biochemical changes, and quality attributes of sweet cherry fruit. *Food chemistry*, 160, 22-30.