

### **RELATIONSHIP BETWEEN ATTENTION DEFICIT HYPERACTIVITY DISORDER AND SEVERE SUBSTANCE USE DISORDER IN ADULT INPATIENTS: A CASE-CONTROL STUDY**

#### **ABSTRACT**

**Objectives:** To evaluate the frequency of Attention Deficit Hyperactivity Disorder (ADHD) in adults interned for treatment of severe chemical dependence in the Therapeutic Community Rios de Água Viva in Coronel Fabriciano, Minas Gerais.

**Method:** A case-control study with male subjects aged 18 years or older, compared to age-matched men without chemical dependence on a 1:1 ratio. Data collection was performed by applying a questionnaire with the Fifth Edition of Diagnostic and Statistical Manual of Mental Disorders (DSM-5) of American Psychiatric Association criteria for diagnosis of ADHD, and information on demographic profile, smoking and age of onset of psychoactive chemical substance use.

**Results:** The vast majority of patients have high school education (72.7%), are not married (93.2%) and are smokers (86.4%). From the 44 patients with Substance Use Disorder who were screened for ADHD symptoms, 22.7% screened positive for the disorder, while only 2.3% of the control group received this diagnosis ( $p < 0.001$ ). The most used drug was alcohol (100%), followed by cocaine-crack (47%) and in third place cannabis (4.5%).

**Conclusion:** Higher frequencies of ADHD symptoms were found among severe chemical dependents interned in Therapeutic Community.

*Keywords:* Attention Deficit Hyperactivity Disorder, Substance Use Disorder, prevalence, diagnosis, Therapeutic Community, adult inpatients.

#### **1 - INTRODUCTION**

Attention Deficit Hyperactivity Disorder (ADHD) is a neurodevelopmental disorder defined in Diagnostic and Statistical Manual of Mental Disorders 5th edition (DSM-5 – American Psychiatric Association 2013) [1] as a persisting pattern of inattention and hyperactivity/ impulsivity that interferes with functioning or development.

ADHD has developmental onset, typically in childhood, and has a high heritability [2]. The etiology of ADHD is also strongly influenced by environmental interactions [3]. Worldwide, this disorder has a childhood prevalence of 4-7% [4,5,6] and 2.5% in adults [7,8]. In approximately half to two-thirds of cases, childhood ADHD persists into adulthood [9,10,11,12], but ADHD is often underdiagnosed in adults [13,14] and brings harmful consequences for the individual when not treated. Among individuals with ADHD, the co-occurrence of other psychiatric disorders is frequent, in particular Substance Use Disorder (SUD) [9].

International research revealed a wide variance in the prevalence of ADHD in SUD patients across countries and cultures [15,16,17], according to the main substance used, methodological factors, the setting (e.g. outpatient vs. inpatient) and different sample sizes. The IASP (The International ADHD in Substance Use Disorders Prevalence)

study estimated an international prevalence of 13.8% of ADHD in SUD patients [18,19].

The comorbidity between drug addiction and ADHD is associated with an earlier onset of substance use and a higher likelihood of use of a variety of substances [22, 23,24,25]. People with ADHD are at increased risk for poorer long-term outcomes, including social, academic and occupational functioning, poorer intellectual performance, worse health outcomes, teenage pregnancy, increased criminal convictions and mortality [25, 26, 27].

Impulsivity is a core symptom in ADHD and is associated with an increased risk for SUD, including alcoholism [29,30]. Cognitive impulsiveness (i.e. impulsive decision making) remains prominent in adult ADHD and is associated with a range of clinical consequences, including poor relational or professional stability [25,34]. Impulsive risk-taking activities among early adolescents may include drug and alcohol use, which may put them at higher risk of long term use. Many studies and reviews show the complex correlation between ADHD and SUD [20,21]. Longitudinal follow up of children in the Multimodal Treatment Study of ADHD found that 64% of adolescents with ADHD were likely to engage in both alcohol and illicit drug use compared with 24% non-ADHD controls [25], making exposure to alcohol/substances one of the most important health-related issues for youth and young adults with ADHD [34]. The relationship is bidirectional: about 40% of adults with ADHD have also SUD [6,32]. It is still a knowledge gap to understand the role of ADHD in the context of patients institutionalized due to chemical dependence.

The treatment of patients with severe chemical dependence is a great challenge. In 1971, the German-French psychiatrist Doctor Claude Olievenstein, specialist in the field of drug addiction, founded the first therapeutic community for treatment of severe chemical dependents in Paris, the Marmottan Center, due to low success rates achieved with other treatment attempts [33].

The treatment in Therapeutic Communities (TC) is based on isolation, work, discipline and spirituality, and lasts nine months. The patients also receive medication and psychotherapy.

The main indications for institutionalization in TC are: serious health problems caused by substance use, loss of work capacity, frequent physical aggression against family members, severe property damage (spending on drugs the necessary resources to support the family) and conflicts with the law (mainly drug trafficking and thefts).

Given the above, the aim of this study was to evaluate the relationship between ADHD and severe substance use disorder in adult inpatient men.

## **2 - MATERIALS AND METHOD**

A case-control study whose cases are inpatient men aged 18 years or older, institutionalized for treatment of severe substance use disorder at the Therapeutic Community Rios de Água Viva (TCRAV) in the city of Coronel Fabriciano, Minas Gerais, Brazil from January 01, 2022 to December 01, 2022. The controls are men with no history of addiction, matched for age on a 1:1 ratio. Individuals who showed signs of dementia and those who had decompensated psychiatric comorbidities during the period of application of the questionnaire were excluded: psychoses, severe depression, manic episode.

The control group are men without chemical dependency, living in the same region as the case group, randomly selected, age-matched and unrelated to drug addicts.

Data collection was performed by applying a questionnaire with the DSM-5 of American Psychiatric Association criteria in addition to sociodemographic data, smoking and age of onset of psychoactive substance use. All participants signed the Informed Consent Form.

This study was approved by the Ethics Committee of Faculdade de Ciências Médicas de Minas Gerais. According to DSM-5 criteria, the diagnosis of ADHD is done in adults who meet five criteria for inattention plus five criteria for hyperactivity/impulsivity. In addition, the symptoms must have started before age 12 years, must occur in more than one location (e.g., home and work), and must interfere with or reduce the quality of social, academic, or professional functioning.

#### *Sample size*

The sample size was calculated to test the difference in prevalence of ADHD among men who are or are not in treatment of severe chemical dependence. Considering a significance level of 5%, minimum power of 80%, using prevalence data from previous studies [18,34,38], it was calculated that 44 cases and 44 controls were needed.

#### *Data analysis*

For data analyses, exploratory statistic technics were used that allowed a better visualization of the general characteristics of the data.

Data were presented in frequency tables with absolute frequencies and their respective percentages, as well as descriptive measures (mean, standard deviation, median, IQR – interquartile range - percentiles 25% and 75%) and the minimum and maximum values for quantitative data.

For comparing categorical data, the Chi-square test was used and, when necessary, the Fisher test.

For the comparison of quantitative variables, the T test was used, due to its normal distribution.

In all tests, the significance level adopted was 5%, therefore comparisons whose p-value was less than or equal 5% were considered significant.

The software used for the analyses was SPSS version 25.0.

### **3 – RESULTS**

#### *3.1 – Proportion of ADHD*

From the 44 SUD patients who were screened for ADHD symptoms, 10 (22.7%) screened positive for the disorder ( $p < 0.001$ ), while only one men of the control group (2.3%) met the DSM-5 diagnostic criteria for ADHD.

Patients with chemical dependence have a higher percentage of ADHD (22.7%) compared to those who do not have chemical dependence (2.3%),  $p = 0.004$  (Figure 1).



**Figure 1: Proportion of ADHD in the evaluated groups**

### 3.2 – Characterization of demographic profile of the patients

The average age was 42.3 years, ranging from 21 to 66 years. Most with high school education (72.7%) and single (72.7%) - Table 1

**Table 1** – Characterization of the demographic profile of the interned patients (n=44) and the controls (n=44)

Variables	Mean (SD)	Median (P25 - P75)	Min - Max
<b>Age</b>	42.32 (11.98)	39.5 (34.75 - 50)	21 - 66

  

Variables	Group		P Value	n (%)
	Patients	Controls		
	n (%)	n (%)		
<b>Marital status</b>				
Married	3 (6.8%)	24 (54.5%)	<b>&lt;0,001<sup>a</sup></b>	
Not married	41 (93.2%)	20 (45.5%)		
<b>Schooling</b>				
Illiterate	1 (2.3%)			
	10 (22.7%)			
Elementary school				
	32 (72.7%)			
High school				
University Education	1 (2.3%)			

SD: standard deviation; P25: percentile 25; P75: percentile 75

### 3.3 – Type of substance used by the patients

Table 2 shows the classification of the group of patients with chemical dependency according to the type of the substance used, and the age of onset of substance use. The use of alcohol and crack/cocaine were the most frequent: 100% e 47.7% respectively.

**Table 2** – Evaluation of the patients with chemical dependence according of the type of substance used and the age of onset (years)

Variables	Mean (DP)	Median (P25 - P75)	Min - Max
<b>Onset of use (in years)</b>	14.2 (3.7)	14 (12 – 15.25)	7 - 28
n (%)			
Type of substance used			
<b>Alcohol</b>		44 (100%)	
<b>Cocaine/Crack</b>		21 (47.7%)	
<b>Cannabis</b>		2 (4.5%)	

### 3.4 – ADHD prevalence and frequency of smoking

Table 3 shows the frequency of smoking in men with and without chemical dependence. Both assessments showed statistically significant difference ( $p \leq 0.05$ ), which shows an association with chemical dependence. A high degree of smokers was observed in patients with chemical dependence (86.4%), with a significant difference in relation to those who do not have this dependence (47.7%),  $p < 0.001$ .

**Table 3** – Evaluation of ADHD and smoking in relation to patients with and without chemical dependence

Variables			P Value
	Patients	Controls	
	n (%)	n (%)	
<b>Smoking</b>			
No	6 (13.6%)	23 (52.3%)	<b>&lt;0,001<sup>q</sup></b>
Yes	38 (86.4%)	21 (47.7%)	

q: Pearson chi square test; fi: Fisher's test; mc: Monte Carlo

### 3.5 – Age of onset of substance used in patients with and without ADHD

In Table 4 we have the evaluation of the age of beginning of substance use and the diagnosis of ADHD. There were no significant statistical differences among patients with severe Chemical dependence, with or without ADHD: the age of onset was similar in both ( $p = 0.592$ ).

**Table 4** – Assessment of the age at onset of substance use in patients with and without ADHD

Variable	ADHD						P (v)
	No (n=34)			Yes (n=10)			
	Mean (SD)	Median (IQR)	Min - Max	Mean (SD)	Median (IQR)	Min - Max	
Age of onset (years)	14.38 (3.37)	14.00 (12.00 – 15.75)	10.00 – 28.0	13.50 (4.72)	13.50 (12.00 – 14.00)	7.00 – 24.00	0.592

t Test

### 3.6 – Number of substances used

The vast majority of interned patients for treatment of Chemical dependence used more than one substance, as shown on the Table 5.

**Table 5** - Characterization of profile of the number of substance used by the interned patients for treatment of severe chemical dependence

Variables	n (%)	P(v)
<b>Substance use</b>		
One substance	26 (59.10%)	-
Two or more substances	18 (40.90%)	
<b>Age of onset (years)</b>		
One substance	Mean (SD) 13.31 (2.619)	0.057*
Two or more substances	15.44 (4.605)	

\*t Test

## 4 – DISCUSSION

As the main result of this study, we found that 22.7% of the individuals in the case group have ADHD symptoms, while only 2.3% of the control group received this diagnosis.

The definition of ADHD diagnosis were based on self-reported DSM-5 questionnaire, with the appropriate explanations made by the doctor and psychologists. In adults, instruments for self-reported symptoms of ADHD have shown adequate psychometric properties [35, 36,37].

It has been estimated that one quarter of those suffering from SUD have comorbid ADHD [38]. The present study corroborates these findings, since 22.7% of the SUD patients meet the DSM-5 diagnostic criteria for ADHD. The prevalence of ADHD among the control group encountered was according to the international literature findings: 2,3% of the non SUD versus 2.5% in the literature [7,8].

Although previous studies showed that substance using individuals with ADHD have earlier onset of substance abuse, higher impairment across several domains of daily life, and worse prognosis in treatment compared to non-ADHD abusers [39, 40], the difference of age onset of substance use did not appear in our sample. This may be due to the severity of SUD in patients who are referred for treatment in Therapeutic Communities and the small sample of cases.

This study also found a high prevalence of smoking and alcohol use among severe chemical dependents. These findings point to the importance of doing efforts to prevent alcohol and tobacco use among young people.

The prevalence of smoking in the control group was much higher than the median of Brazilian population. According to the National Cancer Institute (INCA, 2022), 9.1% of Brazilian population is a smoker: 11.8% of men and 6,7% of women. We assign this high prevalence of smoking in the control group (47.7%) possibly to the low economic level of most individuals in this group and the small sample size.

According to the Brazilian Institute of Geography and Statistics (IBGE), 21% of Brazilians have an University degree. Among the inpatients, only 2.7% had completed higher education. The Therapeutic Community Rios de Água Viva (TCRAV) serves needy patients at low cost and also many patients free of payment. In addition, the average age of the patients is higher than the general population, which helps to explain the finding of only one patient with a university degree. Despite these considerations, further studies are needed to verify whether the prevalence of severe chemical dependence is lower among men who have a university degree.

Among patients with severe chemical dependence, only 6% are married. This data also contrasts with the data of Brazilian population: 47% of adult men are married and 57% declare to have some type of stable union, according to the IBGE. Severe chemical dependence is known to cause separation between couples [40]. Furthermore, being married may also act as a protective factor against substance abuse.

As a strength of this study, we point out the innovation of the investigation the relationship between ADHD and chemical dependence in the context of the patient interned in TC for treatment of severe chemical dependence.

As limitations, we point out the self-reported collection of age at onset of drug use: the memory bias may have altered the final data.

## **5 – CONCLUSIONS**

The aim of this study was to determine association between severe SUD and ADHD. Aside from this, we wanted to explore the relationships between severe SUD and marital status, schooling, smoking and onset of substance use.

We found high prevalence of ADHD symptoms among the patients with severe SUD, low educational level and predominance of not married.

The main substance used by the patients interned for treatment of severe SUD is alcohol, followed by crack/cocaine. Cannabis appears in the third position. This can also be explained by the severity of the substance use disorder presented by patients who need institutionalization.

ADHD may be one of the risk factors for late chemical dependence. From a public health perspective, identifying ADHD in childhood or adolescence and continuous treatment and support would be best to lower risks of SUD, but identification of ADHD in early adulthood may be still relevant for the start of specific ADHD and SUD interventions.

The association between SUD and smoking, low educational status and not being married should be considered for future public policies to combat drug use, aiming for a less violent society, with better mental health.

## **8 – CONSENT**

As per international standard, patients and control group written has been collected and preserved by the authors.

## **9 – ETHICAL APPROVAL**

This study was previously approved by the Ethics Research Committee of Faculdade de Ciências Médicas de Minas Gerais.

Registration: CAAE 59854722.9.0000.5134

## **10 - REFERENCES**

- 1 - American Psychiatry Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-5. 5<sup>a</sup> Ed. Arlington, VA: American Psychiatric Publishing; 2013
- 2 - Faraone SV, Larsson H. Genetics of attention deficit hyperactivity disorder. *Mol. Psychiatry*. 2019;24 (4):562-75. doi: 10.1038/s41380-018-0070-0.
- 3 – Kostenius M et al. Childhood trauma exposure in substance use disorder patients with and without ADHD. *Addict Behav*.2017 Feb;65-118-125. doi: 10.1016/j.addbeh.2016.10.016.
- 4 – Polanczyk G, de Lima MS, Horta ML, Biederman J, Rhode LA. The worldwide prevalence of ADHD: a systematic review and metaregression analyses. *Am J Psychiatry*.2007 Jun;164 (6):942-8. doi: 10.1176/ajp.2007.164.6.942.

- 5 – Willcut EG. The prevalence of DSM-IV attention deficit hyperactivity disorder: a meta-analytic review. *Neurotherapeutics*. 2012 Jul;9(3):490-9. doi: 10.1007/s13311-012-0135-8.
- 6 – Kessler RC et al. The prevalence and correlates of adult ADHD in the United States: results from the National Comorbidity Survey Replication. *Am J Psychiatry*. 2006 Apr;163(4):716-23. doi: 10.1176/ajp.2006.163.4.716.
- 7 – Biederman J, Faraone SV. Attention-deficit hyperactivity disorder. *Lancet*. 2005 Jul 16-22; 366(9481):237-48. doi: 10.1016/S0140-6736(05)66915-2.
- 8 – Simon V, Czobor P, Bálint S, Mészáros A, Bitter I. *Br J Psychiatry*. 2009 Mar;194(3):204-11. doi: 10.1192/bjp.bp.107.048827.
- 9 – Kaye S et al. Persistence and Subtype Stability of ADHD Among Substance Use Disorder Treatment Seekers. *J Atten Disord*. 2019 Oct;23(12):1438-1453. Doi: 10.1177/1087054716629217.
- 10 – Wilens TE, Spencer TJ. Understanding attention deficit/hyperactivity disorder from childhood to adulthood. *Postgrad Med*. 2010 Sep;122(5):97-109. Doi:10.3810/pgm.2010.09.2206.
- 11 – Faraone SV, Biederman J, Mick E. The age-dependent decline of attention deficit hyperactivity disorder: a meta-analysis of follow-up studies. *Psychol Med*. 2006 Feb;36(2):159-65. Doi: 10.1017/S003329170500471X.
- 12 - Lara C et al., Childhood predictors of adult attention-deficit hyperactivity disorder: results from the World Health Organization World Mental Survey Initiative. *Biol Psychiatry*. 2009 Jan 1;65(1):46-54. doi: 10.1016/j.biopsych.2008.10.005.
- 13 – Mc Aweeney M, Rogers NL, Huddleston C, Moore D, Gentile JP. Symptom prevalence of ADHD in a community residential substance abuse treatment program. *J Atten Disord*. 2010 May;13(6):601-8. doi: 10.1177/1087054708329973.
- 14 – Huntley Z et al. Rates of undiagnosed attention deficit disorder in London drug and alcohol detoxification units. *BMC Psychiatry*. 2012 Dec 6;12-23. doi: 10.1186/1471-244X-12-223.
- 15 – Faraone SV, Banaschewski T, Coghill D, Zheng Y, Biederman J, Bellgrove MA, et al. The world federation of ADHD international consensus statement: 208 evidence-based conclusions about the disorder. *Neurosci Biobehav Rev*. 2021; 128:789-818. doi: 10.1016/j.neurobiorev.2021.01.022.

- 16 – Libutzki B, Ludwig S, May M, Jacobsen RH, Reif A, Hartman CA. Direct medical costs of ADHD and its comorbid conditions on basis of a claims data analysis. *Eur Psychiatry*. 2019;58:38-44. doi: 10.1016/eurpsy.2019.01.019. Cdep.2013.09.026.
- 17 – van de Glind G, et al. Variability in the prevalence of adult ADHD in treatment seeking substance use disorder patients: results from an international multi-center study exploring DSM-IV and DSM-5 criteria. *Drug Alcohol Depend*. 2014 Jan 1;134:158-166.doi: 10.1016/j.drug
- 18 - van Emmerik-van Oortmerssen K, van der Glind G, Koeter MWJ, Allsop S, Auriacombe M, Barta C, et al. Psychiatric comorbidity in treatment-seeking substance use disorder patients with and without attention deficit hyperactivity disorder: results of the IASP study. *Addiction*. 2014;109(2):262-72. doi: 10.1111/add.12370.
- 19 – van de Glind G, et al. The International ADHD in Substance Use Disorders Prevalence (IASP) study: background, methods and study population. *Int Methods Psychiatry Res*. 2013 Sep;22(3):232-44. doi: 10.1002/mpr.1397..
- 20 – Ohlmeier MD et al. Comorbidity of alcohol and substance dependence with attention-deficit hyperactivity disorder (ADHD). *Alcohol Alcohol*. 2008 May-Jun;43(3):300-4. Doi: 10.1093/alcalc/agn014.
- 21 – Velez-Pastrana MC, et al., Attention Deficit Hyperactivity Disorder in Prisoners: Increased Substance Use Disorder Severity and Psychiatric Comorbidity. *Eur Addict Res*. 2020 Feb;26(4-5):179-190. Doi: 10.1159/000508829.
- 22 – Wilens TE, Morrison NR. The intersection of attention-deficit hyperactivity disorder and substance abuse. *Curr Opin in Psychiatry*. 2011 Jul;24(4):280-5. doi: 10.1097/YCO.0b013e328345c956
- 23 – Estevez N et al. Adult attention-deficit hyperactivity disorder and its association with substance use and substance use disorder in young men. *Epidemiol Psychiatry Sci*. 2016 Jun;25(3):255-66. doi: 10.1017/S2045796015000360.
- 24 – Charach A, Yeung E, Climans T, Lillie E. *J Am Acad Child Adolesc Psychiatry*. 2011 Jan;50(1):9-21. Doi: 10.1016/j.jaac.2010.09.019.
- 25 – Molina BS, et al. Adolescent substance use in the multimodal treatment of ADHD. *J Am Acad Child Adolesc Psychiatry*. 2013 Mar;52(9):250-63. Doi: 10.1016/j.jaac.2012.12.014. Dalsgaard S, Mortensen PB, Frydenberg M, Thomsen PH. ADHD, stimulant treatment in childhood and subsequent substance abuse in adulthood – a naturalistic long-term follow-up study. *Addict Behav*. 2014; 39(1):325-8. j.jaac.2012.12.014.

2015 Jun;39(3):136-40. Doi: 10.1192/pb.bp.113.043257.

33 – Olievenstein, C. *There Are No Happy Junkies*, Ed. Moraes, 1978

34 - Mucci F, Carpita B, Pagni G, Della Vecchia A, Bjedov S, Pozza A, et al. Lifetime evolution of ADHD treatment. *J Neural Transm (Vienna)*. 2021;128(7):1085-1098. doi: 10.1007/s00702-021-02336-w.

35 – Murphy P, Schachar R. Use of self-ratings in the assessment of symptoms of attention deficit hyperactivity disorder in adults. *Am J Psychiatry*. 2000 Jul;157(7):1156-9. doi: 10.1176/appi.ajp.157.7.1156.

36 – Sandra Kooij, et al. Reliability, validity and utility of instruments for self report and information report concerning symptoms of ADHD in adult patients. *J Atten Disord*. 2008 Jan;11(4):445-58. doi: 10.1177/1087054707299367.

37 – Losano OM, et al. ADHD Adult Self-Report Scale: Comprehensive Psychometric Study in a Spanish SUD Sample. *J Attent Disord*. 2020 Oct;24(12):1674-1684. doi: 10.1177/1087054716664410.

38 - van Emmerik-van Oortmerssen K, van der Glind G, van der Brink W, Smit F, Crunelle CL, Swets M, et al. Prevalence of attention-deficit hyperactivity disorder in substance use disorder patients: a meta-analysis and meta-regression analysis. *Drug Alcohol Depend*. 2012 Apr 1;122(1-2):11-9. doi: 10.1016/j.drugalcdep.2011.12.007.

39 – Liao YT, et al. Depression and severity of substance dependence among heroin dependent patients with ADHD symptoms. *Am J Addict*. 2017 Jan;26(1):26-33. doi: 10.1111/ajad.12487 .

40 – Di Lorenzo R et al. Children and adolescents with ADHD followed up to adulthood: a systematic review of long-term outcomes. *Acta Neuropsychiatrica* 2021 Dec;33(6):283-298. doi: 10.1017/neu.2021.23.