

## Original Research Article

# Evaluation of Assessment of four maize (*Zea mays* L.) genotypes using drought tolerance indices.

### Abstract

Drought tolerance ~~as such~~ is not often considered ~~as~~ an independent trait by breeders. The objective of this study was to evaluate ~~and identify drought tolerant genotypes using~~ eight drought tolerance indices ~~, namely namely~~ the Stress Susceptibility Index (SSI), the YSI, the YR (Yr), yield index (YI), tolerance index (TOL), average productivity (MP), mean geometric productivity (GMP) and stress tolerance index (STI) of maize genotypes (*Zea mays* L.). ~~A field trial was conducted to evaluate four~~ Four genotypes ~~were evaluated in the field~~ during the hot dry season of 2016 and 2017 at the irrigated perimeter of Jirataoua. Drought tolerance indices were calculated on the basis of yield under optimal and stressed conditions. The comparison of the means of drought tolerance ~~demonstrated and yield validated~~ the ~~important~~ effects of drought on yield ~~and showed~~ ~~as well as~~ significant differences between genotypes. The ~~calculated~~ correlation coefficient ~~results~~ and the principal component analysis showed that the GMP, MP and STI indices were able to discriminate ~~between~~ drought-sensitive and tolerant genotypes. Two genotypes CZH131001 and CZH142013 ~~produced high grain yield under both gave a good yield in~~ optimal and stressed conditions. Overall, ~~we concluded that~~ GMP, MP, and STI indices can be ~~used as~~ effectively ~~exploited not only for~~ drought tolerance screening ~~indices and able~~ ~~but also~~ to identify ~~a better~~ genotypes, suitable for both optimal ~~and stress~~ conditions ~~and stressed~~.

**Key words:** ~~genotypes~~, correlation coefficient, drought tolerance index, ~~genotypes~~, principal component analysis

### 1. INTRODUCTION

Water is one of the key factors in agricultural production. ~~and Although water is the most abundant molecule on the earth's surface~~, its availability has a strong influence on agricultural production [1]. It also plays an important role in the transfer of salts and nutrients [2]. Low rainfall and lack of sufficient irrigation ~~water~~ are the ~~challenges of crop production main causes of water availability for crops~~ in arid and semi-arid regions. The effects of water stress on growth and yield components are very different. Yield loss is a major concern for farmers and breeders alike. Consequently, the focus is on genotypes selected for yield performance under water stress conditions. Research has been underway since the early 1980s, with the aim of associating yield variations and their interactions with growing environment conditions with stress tolerance indices based on yield loss under water deficit compared with normal conditions [3]. The results of several previous investigations have shown that genotype x environment (G x E) interactions could be described in part by stress tolerance indices [4]. These indices provide a measure of impairment and enable adequate screening of stress-tolerant genotypes [5].

But the variation in yield potential comes from factors related to adaptation rather than drought tolerance. Thus, drought indices provide a measure of drought based on yield losses under drought conditions compared to normal conditions and used for the selection of drought-tolerant genotypes [5]. [6] proposed a cultivar stress

**Comment [H1]:** Lacks consistency and coherence. It should start from importance of the crop or subject of the study followed by problems and gaps. This may be followed by the methods or problem solving alternatives and associated literatures that support the new approaches etc. Hypothesis and purpose of the study will be followed. The paper has to be modified in such away that it would be more clear for the reader than the current status. It needs to be modified and the author or authors are expected to rewrite it.

**Comment [H2]:** Rewriting and rephrasing is required.

susceptibility index (SSI). Lower SSI values (< 1) indicate low yield variation in stressed and unstressed environments and demonstrate greater yield stability, and higher SSI values (> 1) suggest greater susceptibility [7-8-9]. [10] defined a new leading index (STI = stress tolerance index), which can be used to identify high-yielding genotypes under both optimal and stressed conditions. He suggested that selection based on STI results in genotypes with higher stress tolerance and yield potential. IL demonstrated that GM indices are mathematical derivatives of yield. Thus, genotypes ranked with high yields under both conditions rather than relative performance are performances are the best indicator for assessing drought tolerance [11]. [12] observed that genotypes with high YSI show greater stability in the environment. Based on the existed knowledge it is hypothesized that YSI is therefore a good measure of the use of effectiveness of drought tolerance indices and is more important useful for selection criteria than simply estimating absolute yield under stress. The aim of this study was to evaluate-identify drought tolerant genotypes and suitable drought tolerant screening indices, selection criteria for identifying drought tolerant genotypes with high yields under both optimal and stressed conditions.

**Comment [H3]:** Rewrite the sentences. You need to indicate the hypotheses and purpose of the study clearly..

## 2. MATERIALS AND METHODS

**Comment [H4]:** The materials and methods should be written in detail. It is better to rewrite it

### 2.1. Plant material and growing conditions

Experiments were carried out on four maize genotypes. Four Three genotypes namely, (CZH131001, CZH142013, SC303) obtained from were obtained from the (CIMMYT) and the fourth (P3K) from the National Institute of Agronomic Research of Niger (INRAN) were used in this study. P3K is widely used locally, while CZH131001, CZH142013 and SC303 are renowned for their drought tolerance (CIMMYT HARARE). The genotypes except P3K were developed for drought tolerance. The experiment was conducted at the Keguel site in the Jiratawa irrigated site in 2016 and 2017 dry season using irrigation. The experimental site is 10 km from the town Maradi, Niger, located in latitude perimeter (latitude 13°41' and longitude = 7°14') in altitude of ..... 40 km from the town of Maradi, Niger, under conditions of The maximum and minimum temperature: 45.44-34.88°C: 28.74-12.93°C and min: max relative humidity: 20.36-49.28%: 83.00-99.26%) during the hot season (March to June). The soils are sandy-loam, representing 95% of the irrigated perimeter soils. These soils have a light texture with 85.17% (w/w) sand, 13.44% (w/w) silt and 1.39% (w/w) clay. The PH<sub>eau</sub> and PH<sub>CaCl2</sub> of the soil solution are 7.20 and 6.97 respectively.

**Formatted:** Font color: Blue

**Formatted:** Font color: Blue

**Formatted:** Font color: Blue

**Formatted:** Font color: Blue

**Formatted:** Font color: Blue

**Formatted:** Font color: Blue

Drought tolerance indices were calculated using the following equations:

Ten drought tolerance indices, namely stress tolerance index (STI), mean productivity (MP), geometric mean productivity (GMP), tolerance index (TOL), stress susceptibility index (SSI), yield stability index (YSI), yield reduction rate (Yr) and yield index (YI) were calculated on the basis of yields under drought (Y<sub>s</sub>) and irrigated (Y<sub>p</sub>) conditions. Drought tolerance indices were calculated using the following equations:

**Comment [H5]:** It would be good to summarize all the information (climate and soils data) and put in the table

Indice de susceptibilité au stress (SSI) 
$$SSI = \frac{1 - \frac{Y_d}{Y_p}}{DII} \quad [6]$$

**Comment [H6]:** Include references to each formula

Yield Stability Index (YSI) 
$$YSI = Y_d / Y_p \quad [13]$$

Yield index (YI) 
$$YI = Y_d / \bar{Y}_d \quad [14]$$

Stress Tolerance Index (STI) 
$$STI = \frac{(Y_p * Y_d)}{(Y_p)^2} \quad [10]$$

Yield reduction (Yr) 
$$Yr = 1 - (Y_d / Y_p) \quad [15]$$

Productivité géométrique moyenne (GMP) 
$$GMP = \sqrt{(Y_d * Y_p)} \quad [16]$$

Tolerance index (TOL) 
$$TOL = Y_d - Y_p \quad [17]$$

Average productivity (MP) 
$$MP = (Y_p + Y_d) / 2 \quad [17].$$

### 2.2. Data Statistical analysis

Analysis of variances was undertaken using ..... Comparison of means, Phenotypic correlations was done coefficients was analysed with ..... between indices and yield and P principal component analysis (PCA) based on selection criteria (drought tolerance indices) and genotypes were carried out using XLSTAT 2019 software

version 21.1.2.56803. ~~This was done to interpret relationships between selection criteria and compare genotypes on the basis of drought tolerance indices, enabling the identification of genotypes with a certain level of drought tolerance.~~

### 3. RESULTS

#### 3.1. Comparison of genotypes based on stress resistance and tolerance indices

~~Drought tolerance and yield are complex traits affected by different factors. To find appropriate stress resistance indices for the selection of drought tolerant genotypes, the grain yield of maize genotypes under optimal and stressed conditions was measured to calculate different sensitivity and tolerance indices. Table 1 shows the mean values of various indices is given in Table 1, used to characterize the drought tolerance and adaptation of genotypes to different environments. Significant variation among genotypes was observed for grain yield under optimum and drought conditions. Drought stress reduced the grain yield of maize genotypes and the genotypes respond differently due to the effect of drought as indicated by drought indices. Among the four genotypes, CZH142013 and CZH1310001 showed high values of stress tolerance index (STI), geometric mean productivity (GMP) and mean productivity (MP) compared with genotypes SC303 and P3K. Consequently, these two genotypes with high STI, GMP and MP values can be selected as water stress tolerant genotypes. These tolerant genotypes also recorded high Yield Stability Index (YSI) and Yield Index (YI) values. In addition, these genotypes showed higher tolerance than the other two (SC303 and P3K) due to the low values of tolerance index (TOL), yield reduction (Yr) and stress susceptibility index (SSI). Consequently, these two genotypes with high STI, GMP and MP values can be selected as water stress tolerant genotypes. At the same time, the Stress Susceptibility Index (SSI) estimates the rate of yield variation for each genotype between optimal and stress conditions, relative to the average variation for all genotypes.~~

**Comment [H7]:** What was observed should be clearly written. I suggest to rewrite the findings of the study clearly.

**Comment [H8]:** Please explain the findings accordingly.

**Table 1: Average comparison of drought tolerance indices and grain yield (Kg/ha) of different maize genotypes under stressed and unstressed conditions.**

Genotypes	Yp	Yd	Yr	YSI	SSI	STI	GMP	MP	TOL	YI
CZH142013	954,4 <sup>ab</sup>	732,71 <sup>b</sup>	23 <sup>bc</sup>	76,77 <sup>ab</sup>	0,8 <sup>b</sup>	1,13 <sup>a</sup>	836,24 <sup>ab</sup>	843,56 <sup>ab</sup>	121,69 <sup>b</sup>	1,30 <sup>ab</sup>
CZH1310001	1046,77 <sup>a</sup>	1012,39 <sup>a</sup>	3 <sup>c</sup>	96,65 <sup>a</sup>	0,12 <sup>b</sup>	1,71 <sup>a</sup>	1029,43 <sup>a</sup>	1029,58 <sup>a</sup>	34,38 <sup>c</sup>	1,80 <sup>a</sup>
SC303	572,8 <sup>b</sup>	139,31 <sup>c</sup>	75 <sup>a</sup>	24,31 <sup>c</sup>	2,7 <sup>a</sup>	0,12 <sup>b</sup>	282,48 <sup>c</sup>	356,06 <sup>c</sup>	433,49 <sup>a</sup>	0,25 <sup>c</sup>
P3K	568,46 <sup>b</sup>	334,89 <sup>bc</sup>	41 <sup>b</sup>	58,9 <sup>b</sup>	1,46 <sup>a</sup>	0,3 <sup>b</sup>	436,31 <sup>bc</sup>	451,68 <sup>bc</sup>	233,57 <sup>b</sup>	0,59 <sup>bc</sup>

Means followed by the same letters in each column are not significantly different ( $p < 0.01$ ). Yp: yield under optimal conditions, Yd: yield under water stress, GMP: geometric mean productivity, MP: mean productivity, SSI: stress susceptibility index, STI: stress tolerance index, Yr: yield reduction ratio, YSI: yield stability index, YI: yield index, TOL: tolerance index.

#### 3.2. Correlation analysis

~~Phenotypic correlation coefficients of grain yield under drought stress, no stress and drought tolerant indices are given in Table 2. The correlation coefficient of grain yield under stress were positive and strong with drought tolerant indices (GMP=0.99); (M=0.99); (STI=0.99); (YI=0.99); and (YSI=0.96). In constrast, grain yield under drought were negatively correlated with (SSI=-0.96) and (TOL=-0.92).~~

Some indices have been developed on the basis of quantitative traits under optimal and stressed conditions that can be used as indicators of stress tolerance. To determine the most desirable drought tolerance criteria, phenotypic correlation coefficients between Yd, Yp and other quantitative drought tolerance indices were calculated (Table 2). As a general rule, indices with a high correlation with yield under both optimal and stressed conditions are presented as the best, as they can separate high-yielding genotypes under both conditions. An appropriate index should have a significant correlation with yield under both conditions [5].

**Comment [H9]:** Please rewrite and correct accordingly.

**Table 2: Correlation coefficients between Yp, Yd and the drought tolerance and susceptibility indices of 4 maize genotypes**

	SSI	GMP	MP	STI	TOL	YI	YSI	Yd	Yp	Yr
ISD	1,00									
GMP	-0,96 <sup>*</sup>	1,00								
MP	-0,94 <sup>ns</sup>	0,99 <sup>***</sup>	1,00							
STI	-0,93 <sup>ns</sup>	0,99 <sup>**</sup>	0,99 <sup>**</sup>	1,00						
TOL	0,97 <sup>*</sup>	-0,89 <sup>ns</sup>	-0,87 <sup>ns</sup>	-0,89 <sup>ns</sup>	1,00					
YI	-0,96 <sup>*</sup>	0,99 <sup>**</sup>	0,99 <sup>**</sup>	0,99 <sup>**</sup>	-0,92 <sup>ns</sup>	1,00				
YSI	-0,99 <sup>***</sup>	0,95 <sup>*</sup>	0,94 <sup>ns</sup>	0,93 <sup>ns</sup>	-0,97 <sup>*</sup>	0,96 <sup>*</sup>	1,00			
Yd	-0,96 <sup>*</sup>	0,99 <sup>**</sup>	0,99 <sup>**</sup>	0,99 <sup>**</sup>	-0,92 <sup>ns</sup>	0,99 <sup>***</sup>	0,96 <sup>*</sup>	1,00		
Yp	-0,88 <sup>ns</sup>	0,98 <sup>*</sup>	0,99 <sup>**</sup>	0,98 <sup>*</sup>	-0,79 <sup>ns</sup>	0,97 <sup>*</sup>	0,97 <sup>*</sup>	0,97 <sup>*</sup>	1,00	
Yr	0,99 <sup>***</sup>	-0,96 <sup>*</sup>	-0,94 <sup>ns</sup>	-0,93 <sup>ns</sup>	0,97 <sup>*</sup>	-0,96 <sup>*</sup>	-0,99 <sup>***</sup>	-0,96 <sup>*</sup>	-0,88 <sup>ns</sup>	1,00

ns. \* and \*\*: non-significant and significant at the 5% and 1% probability levels, respectively. Yp: yield under optimal conditions, Yd: yield under stressed conditions, GMP: geometric mean productivity, MP: mean productivity, STI: stress tolerance index, SSI: stress susceptibility index, Yr: yield reduction ratio, YSI: yield stability index, YI: yield index, TOL: tolerance index.

### 3.3. Principal component analysis

A PCA was performed using the tolerance indices and then the genotypes were subjected to a biplot analysis to obtain the relationships between the indices (Table 3). Many researchers have used this analysis to compare different genotypes for different criteria and different species. The results of the principal component analysis (PCA) showed that the first two components explained 95.84% and 3.75% of the total variation. The PCA revealed that the first component (F1) explained 95.84% of the variation in total yield, and was positively correlated with Yp, Yd, YSI, YI, MP, GMP and STI. This component was therefore linked to yield potential and drought tolerance.

**Table 3: Principal component analysis results for grain yield of different maize genotypes under optimal (Yp) and stressed (Yd) conditions, geometric mean productivity (GMP), mean productivity (MP), stress tolerance index (STI), stress susceptibility index (SSI), yield reduction rate (Yr), yield stability index (YSI), yield index (YI) and stress tolerance index (TOL).**

Tolerance index	F1	F2
Yp	0,307	0,508
Yd	0,322	0,107
Yr	-0,317	0,298
YSI	0,316	-0,306
STI	0,318	0,252
GMP	0,321	0,180
MP	0,319	0,266
TOL	-0,303	0,525
YI	0,322	0,107
SSI	-0,317	0,306

F2  
(3,  
75)

F1 (95,84 %)

Figure 1: Biplot drawn on the basis of first and second axes obtained from principal component analysis using stress susceptibility index (SSI), yield reduction rate (Yr), yield index (YI), stability index (YSI), stress tolerance (TOL), mean productivity (PM), geometric mean productivity (GMP), stress tolerance index (STI) and yield under optimal (Yp) and stress (Yd) conditions of 4 maize genotypes.

#### 4. DISCUSSION

In this study drought intensity of about 0.27 % was observed and it was ~~Water stress was~~ considered as moderate based on the work of various authors. (Include references associated to similar reports). ~~moderate with an intensity of DH = 0.27.~~ Water stress is considered severe for intensity values above 0.7 [11]. [18] suggested that simultaneous evaluation of genotypes under optimal and stressed conditions would appear to be the most suitable procedure for selecting genotypes in environments frequently confronted with abiotic stresses such as water

Comment [H10]: No biplot is given in the paper.

Comment [H11]: Please summarize the major findings in this section and explain the phenomena with evidences. Use and include recent referenes

deficit. Indeed, selection based solely on the performance of genotypes under normal conditions does not necessarily lead to productivity gains under stressed conditions, and vice versa [10]. Several selection criteria have been proposed to evaluate the performance of plant species under optimal and stressed conditions. [6] proposed the water stress susceptibility index (SSI) as an indicator of a genotype's sensitivity to water stress. SSI values below 1 indicate low sensitivity to drought (or high yield stability), and values above 1 indicate high sensitivity to drought (or low yield stability). [10] defined the geometric mean productivity (GMP) and the water stress tolerance index (STI), which could be used to identify high-yielding genotypes under both normal and stressed conditions. The higher the STI value, the more drought-tolerant the genotype and the higher its yield potential. According to [19], a genotype's ability to exhibit high yield and geometric mean productivity (GMP) is linked to its drought tolerance. The value of STI as a function of yield under optimal and stressed conditions showed that the genotypes (CZH1310001 and CZH142013) with the highest yields in both environments also had the highest tolerance index (STI). These results are in line with those of [20] and [10], who respectively showed that STI values were higher in sesame and wheat genotypes with the highest yields in two contrasting environments. Similarly, working on maize, [21] found STI to be the best predictor of high yields under different environmental conditions for the selection of lines for water stress, while [22] selected from among five stress indices, the stress tolerance index (MSTI) to which they made a modification using a correction factor  $k_i$ . The STI was even suggested for the selection of tolerance to high temperatures [23]. When stress is severe, TOL, SSI and STI were more useful as indices to distinguish between resistant and susceptible, although no single indicator could clearly identify high-yielding cultivars under stressed and unstressed conditions. It has been concluded that the effectiveness of selection indices under severe stress confirms that different stress conditions influence yield under stress [24-25]. The SC303 and P3K genotypes showed susceptibility due to higher TOL and SSI values. Our results concur with those of [7] who used the stress susceptibility index (SSI) to assess drought tolerance in wheat genotypes. It was observed that both TOL and SSI were successful in selecting high-yielding genotypes under drought stress.

The correlation matrix shows that a significant positive correlation was observed between yield under stressed ( $Y_d$ ) and optimal ( $Y_p$ ) conditions with stress tolerance index (STI), mean productivity (MP), geometric mean productivity (GMP), indicating that these criteria discriminated between drought-tolerant genotypes with high yields in stressed and unstressed environments. Our results are similar to observations made by [4-22-26-27-28-29-30], who claim that these three indices (STI, MP and GMP) would be the best predictors of yields under optimal and stressed conditions. [31] also concluded that MP, GMP and STI were practical parameters for selecting high-yielding wheat genotypes under optimal and stressed conditions. [32] found that STI and GMP, which showed the strongest correlation with yield under optimal and stressed conditions, can be used as the best indices for breeding programs aimed at introducing drought-resistant hybrids. A significant negative correlation was observed between ( $Y_d$ ) with the tolerance index (TOL), the stress susceptibility index (SSI) and the yield reduction ratio ( $Y_r$ ). The tolerance index (TOL) was not significantly correlated with  $Y_p$  and MP. Consequently, indices with significant associations were also able to identify drought-tolerant genotypes. The higher the TOL and SSI values, the higher the yield production under optimal conditions, the greater the reduction in production under stressed conditions. Our results are similar to those reported by [33] on soft wheat cultivars, who conclude that the higher the TOL values, the greater the yield reduction under water stress and the higher the sensitivity to stress. Similarly, [30] working on durum wheat, suggest that selection for TOL decreases yield under water deficit and increases it under optimal water conditions.  $Y_d$  and  $Y_p$  were negatively correlated with SSI in both environments. SSI has been widely used by many researchers to identify drought-sensitive and drought-tolerant genotypes [7-30]. Since in both water regimes, yield under optimal and stressed conditions ( $Y_d$  and  $Y_p$ ) is significantly correlated with the STI and GMP factors, these two indices can be validly used to discriminate between water-stress-tolerant and water-stress-sensitive genotypes. In fact, indices that correlate with yields ( $Y_d$  and  $Y_p$ ) under both optimal and stressed conditions are considered the best because they can separate genotypes with high yields under both water regimes [10]. The latter author introduced the STI and GMP factors to select bean genotypes that are both drought-tolerant and have high yields in contrasting environments.

PCA revealed that genotypes with high F1 values (Table 3) should have high yields under both conditions. Similar results were reported by [30-34] in durum wheat and in cotton. F2 explained 3.75% of the variation in total yield and correlated positively with TOL,  $Y_r$  and SSI. F2 is associated with yield under stress conditions and stress sensitivity. Consequently, F1 and F2 can be considered as yield potential and stress susceptibility respectively. Biplot results based on F1 and F2 data for the 4 genotypes showed that genotypes CZH142013 and CZH1310001 are close to the best drought tolerance indices with high F1, but low F2 values (figure 1). On the other hand, the two genotypes (SC303 and P3K) with low F1 and high F2 values were identified as sensitive genotypes. Our results are in line with the report by [35] who found that wheat genotypes with higher F1 and lower F2 values had high yields (stable genotypes) and genotypes with lower F1 and higher F2 scores had low yields (unstable genotypes).

**Comment [H12]:** Link the discussion with the findings of your study.

**Formatted:** Font color: Red

**Formatted:** Font color: Red

## 5. CONCLUSION

The aim of this study is to evaluate selection criteria for identifying drought-tolerant genotypes with high yields under both optimal and stressed conditions. As stress tolerance indices are only mathematical derivations of the same yield data, selection based on a combination of different stress tolerance indices would be able to characterize the most productive and water-stress resistant genotypes. The positive and significant correlations obtained with Yd and Yp and MP, GMP and STI lead to the conclusion that these indices are the best for determining yield in both environments. The tolerant genotypes CZH1310001 and CZH142013 with strong correlations with these three indices showed the highest yields under both optimal and stressed conditions. Whereas genotypes (SC303 and P3K) with high values of TOL and SSI were able to produce high yields only under optimal conditions. Consequently, breeders can select genotypes that are tolerant under water-stressed conditions and compare their performance under optimal conditions using MP, GMP and STI indices.

**Comment [H13]:** Summarize the discussions and give your recommendation. You are also expected to include the limitations and significance of the study. Please rewrite this section accordingly.

## REFERENCES

1. Pospisilova J, Synkova H, Rulcova J. (2000). Cytokinins and water stress. *Bio.Planta*. 43: 321-328.
2. Ahmadi, A., Mohammadi, V., Sio-Se Mardeh, A. and Poustini, K. (2008). Evaluation of wheat yield drought resistance indices across water regimes. *Acta Biol Szeged* 52(1): 97-100.
3. Salisbury FB, Ross C. (1992). *Plant Physiology*, Ed 4, Wadsworth, Belmont, CA, USA.
4. Maleki, A., Babaei, F., Cheharsooghi, H.A., Ahmadi, J. and Dizaji, A.A. (2008). The study of seed yield stability and drought tolerance indices of bread wheat genotypes under irrigated and non-irrigated conditions. *Res. J. Biol. Sci.* 3(8): 841-844.
5. Mitra J. (2001). Genetics and genetic improvement of drought resistance in crop plants. *Curr. Sci.* 80:758-762.
6. Fischer AT, Maurer R. (1978). Drought resistance in spring wheat cultivars. I. Grain yield responses. *Aust. J. Agric. Res.* 29: 897-912
7. Clarke JM, De Pauw RM, Townley-Smith TM. (1992). Evaluation of methods for quantification of drought tolerance in wheat. *Crop Sci.* 32:728-732.
8. Guttieri MJ, Stark JC, Brien KO, Souza E. (2001). Relative sensitivity of spring wheat grain yield and quality parameters to moisture deficit. *Crop Sci.* 41:327-335.
9. Ullah I, Mehboob-Ur-Mehboob, Zafar Y. (2006). Genotypic variation for drought tolerance in cotton (*Gossypium hirsutum* L.): seed cotton yield responses. *Pak. J. Bot.* 38:1679- 1687.
10. Fernandez GCJ. (1992). Effective selection criteria for assessing plant stress tolerance. In: C.G. Kuo, editor, Adaptation of Food Crops to Temperature and Water Stress: Proceedings of an International Symposium, Taiwan. 13- 18 Aug. 1992. *Asian Vegetable Res. and Dev. Ctr.*, Shanhua, Tainan, pp 257-270.
11. Ramirez P, Kelly J. (1998). Traits related to drought resistance in common bean. *Euphytica* 99: 127-136.
12. Rajamani A. (1994). Screening *Gossypium hirsutum* genotypes for drought tolerance. *Madras Agric. J.* 81:465-468.
13. Bouslama M, Schapaugh WT. (1984). Stress tolerance in soybean. Part I: evaluation of three screening techniques for heat and drought tolerance. *Crop Sci.* 24: 933-937.
14. Gavuzzi P, Rizza F, Palumbo M, Campalino RG, Ricciardi GL, Borghi B. (1997). Evaluation of field and laboratory predictors of drought and heat tolerance in winter cereals. *Can. J. Plant Sci.* 77:523-531.
15. Golestani-Araghi S, Assad MT. (1998). Evaluation of four screening techniques for drought resistance and their relationship to yield reduction ratio in wheat. *Euphytica* 103: 293-299.
16. Kristin AS, Senra RR, Perez FI, Enriquez BC, Gallegos JAA, Vallego PR, Wassimi N, Kelley JD (1997). Improving common bean performance under drought stress. *Crop Sci.* 37: 43-50.
17. Rosielle AA, Hamblin J. (1981). Theoretical aspect of selection for yield in stress and non-stress environment. *Crop Sci.* 21: 943-946.
18. Rajaram, S., Braum, H. L. & Van Ginkel, M. (1996). CIMMYT's approach to breed for Drought tolerance. *Euphytica* 92: 147-153.
19. Toker, C. & Çagiran, M. I. (1998). Assessment of response to drought stress of chickpea. *Int. Chickpea Pigeon pea Newslett.* 11: 8-10.
20. Boureima, S., A. Oukarroum, M. Diouf, N. Cisse and P. Van Damme. 2012. Screening for drought tolerance in mutant germplasm of sesame (*Sesamum indicum*) probing by chlorophyll a fluorescence. *Environ. Exp. Bot.* 81: 37-43.

**Comment [H14]:** Lack of consistency in using dot, bracket, italics and etc. It is good to use more recent references.

**Comment [H15]:** Incomplete and not correctly written. Follow the guidelines of the journal.

21. Moghaddam, A. and Hadizadeh, M.H. (2002). Response of corn (*Zea mays* L.) hybrids and their parental lines to drought using different stress indices. *Seed and Plant*, Vol. 8 n°3: 255-272.
22. Farshadfar, E. and Sutka, J. (2002). Screening drought tolerance criteria in maize. *Acta Agronomica Hungarica*, 50(4): 411-416.
23. Porch T.G. (2006). Application of stress indices for heat tolerance screening of common bean (*Phaseolus vulgaris*). *J. Agron. and Crop Sci*, Vol. 192 n° 5: 390-394.
24. Blum A. (1996). Crop responses to drought and the interpretation of adaptation. *Plant Growth Regul.* 20:135-148.
25. Pantuwan G, Fukai S, Cooper M, Rajatasereekul S, O'Toole JC. (2002). Yield response of rice (*Oryza sativa* L.) genotypes to different types of drought under rainfed lowlands. Part 1: Grain yield and yield components. *Field Crop Res* 73: 153-168.
26. Toorchi M, Naderi R, Kanbar A, Shakiba MR. (2011). Response of spring canola cultivars to sodium chloride stress. *Ann. Bio. Res.* 2: 312-322.
27. Dehghani GH, Malekshhi F, Alizadeh B. (2009). A study of drought tolerance indices in canola (*Brassica napus* L.) genotypes. *J. Sci. Technol. Agri. Natural Reso.* 13:77-90.
28. Khalili M, Naghavi MR, Pour AR, Talebzadeh J. (2012). Evaluating of drought stress tolerance based on selection indices in spring canola cultivars (*Brassica napus* L.). *J. Agri. Sci.* 4: 78-85.
29. Khodarahmpour Z. and Hamidi J. (2011). Evaluation of drought tolerance in different growth stages of maize (*Zea mays* L.) inbred lines using tolerance indices. *Afr. J. Biotechnol.* 10:13482-13490.
30. Golabadi M, Arzani A, Mirmohammadi SAM. (2006). Assessment of drought tolerance in segregating populations in durum wheat. *Afr. J. Agric. Res.* 1:162-171.
31. İlker E., "Correlation and path coefficient analyses in sweet corn" (2011). *Turk. J. Field Crops* 16 (2): 105-107, 2011.
32. Jafari A, Paknejad F, Jami Al-Ahmadi M. (2009). Evaluation of selection indices for drought tolerance of corn (*Zea mays* L.) hybrids. *Inter. J. Plant Prod.* 3:33-38.
33. Sio-Se Mardeh A, Ahmadi A, Poustini K, Mohammadi V. (2006). Evaluation of drought resistance indices under various environmental conditions. *Field Crop Res.* 98: 222-229.
34. Chandrakant Singh, Vijay Kumar, Indivar Prasad, Vishal R. Patil, Rajkumar B.K. (2015). Response of Upland Cotton (*G.hirsutum* L.) Genotypes to Drought Stress Using Drought Tolerance Indices. *J. Crop Sci. Biotech.* 19 (1): 53 – 59.
35. Kaya Y, Palta C, Taner S. (2002). Additive main effects and multiplicative interactions analysis of yield performances in bread wheat genotypes across environments. *Turkish J. Agric. For.* 26: 275-279.