

Original Research Article

Cultural, morphological and molecular characterization of *Ustilaginoidea virens* causing rice false smut in Tamil Nadu

ABSTRACT

Rice false smut, caused by the fungus *Ustilaginoidea virens* is one of the most destructive and re-emerging disease in many rice growing areas of Tamil Nadu, India. Studies on cultural, morphological and molecular characteristics of *U. virens* may provide a more detailed look at disease control and cultivar breeding. The current study was carried out to characterize different isolates of *Ustilaginoidea virens* collected from major rice growing areas in Tamil Nadu, India. Cultural and morphological characteristics of the false smut pathogen on a different culture media as well as genetic variation were studied using specific ITS primers by PCR technique for *U. virens* isolates. The pathogen was isolated by different isolation methods, out of which inner spore dust was effective. The colony was milky white to yellowish white, fluffy, leathery and compact. The colony diameter ranged between 35.33 mm and 65.33mm. The mycelium was hyaline, septate and width of the hyphae ranged between 1.84 and 3.10 μm . The size of the conidia was also varied among the isolates and the maximum size (6.35 μm) was observed in FS 6 isolate and minimum in FS 2 (4.10 μm). The growth of fungus in different media was examined. The maximum colony diameter was observed in Potato Sucrose Agar (PSA) medium and minimum growth in Czapek Agar (CA) medium. The molecular confirmation was done through *U. virens*-specific internal transcribed spacer (ITS) region amplification using the primers, US 1-5/US3-3. The primers amplified a product of 380 bp which are specific for the fungus *U. virens*. The identity of the fungus was further confirmed through ITS sequencing which showed 98-99% sequence homology. The dendrogram showed two main clusters; cluster I comprised of five isolates (FS1, FS2, FS9, FS10 and FS3), while Cluster II consisted of five isolates (FS4, FS8, FS6, FS5 and FS7).

Key words: Characterization, false smut, media, rice, Ustilaginoidea virens

1. INTRODUCTION

Rice is one of the most important and widely grown food crops in the world. The production and productivity of rice is affected by various biotic and abiotic factors. Rice false smut, which is also known as pseudo-smut, or green smut is one of the most severe re-emerging fungal diseases in rice. False smut is caused by an ascomycetous fungus *Ustilaginoidea virens* (Cooke) (Takahashi) (Teleomorph: *Villosiclava virens*) which cause significant quantitative and qualitative yield losses in grain (Rush *et al.*, 2000). Excessive usage of nitrogen fertilizer, large scale cultivation of hybrids and high yielding variety with low disease resistant made the false smut as destructive disease of rice worldwide (Zhou *et al.*, 2014). Initially the white fungal mass emerges from the inside of a spikelet during the initial infection stage, which then changes into a light-yellow smut ball that eventually turns greenish-black. The false smut ball fills with powdery chlamydospores as it matures, and the colour changes to yellowish, yellowish orange, green, olive green, and at last greenish black when smut balls rupture, powdery dark green spores are released (Jing fan *et al.*, 2016). The pathogen produces sclerotia (sexual stage) and chlamydospore (asexual stage). The sclerotia are the main source of primary inoculum which usually develop in the autumn (Ikegami, 1961). The sclerotia germinate to produce millions of ascospores. These ascospores produce conidia which infects the rice plant during flowering stage (Biswas, 2001). During sexual cycle, chlamydospores are produced which cause secondary infection via wind and rain (Huang *et al.*, 2019).

The duration and severity of a disease epidemic are highly influenced by the weather. The disease rate will be high if the temperature ranges between 22°C and 28°C, and rainy and humid condition during the rice booting stage (Wang *et al.*, 2005). Although the pathogens begin to affect rice during the growth of the panicle, symptoms don't appear until flowering. Initially affects a small number of grains and it spreads to the entire panicle, yet in cases of severe disease incidence, the number may rise to 100% (Ladhalakshmi *et al.*, 2012). The causative organism was also identified to produce an enormous amount of mycotoxin, such as ustiloxins and ustilaginoidins which is toxic to human and animal health (Zhou *et al.*, 2012). Due to the pathogen's difficulties with artificial culturing and inoculation, research on false smut was very less in Tamil Nadu. With this in consideration, the present study was undertaken to know about cultural, morphological and molecular diversity of false smut pathogen collected from various rice growing areas of Tamil Nadu.

2. MATERIALS AND METHOD

2.1 Collection and isolation of false smut pathogen

Paddy seeds infected with false smut pathogen were collected from major rice growing areas of Tamil Nadu by conducting roving surveys. The incidence (%) of false smut per panicle was also recorded during the survey. The infected panicle having smut ball symptoms were collected in a polythene bag and bought to the laboratory. The fungus strains were isolated on Potato Sucrose Agar (PSA) plates by inner spore dust method as described by Bashyal *et al.* (2020). The smut balls were surface sterilized by dipping them in 1% sodium hypochlorite solution for 1 min followed by 70% ethanol wash for 1 min and subsequently washed three times with sterile distilled water. The smut balls were cut into two halves using a sterilized scalpel and spores were dusted from the inner surface of the ball with sterilized needle onto the Petri plates containing PSA medium amended with streptomycin sulphate @ 100 ppm. The plates were incubated at 27±1 °C for 7 days. Pure culture was obtained by hyphal tip method using sterilized needle and transferred to PSA plate. The cultures were maintained in slants at 4°C for further use and the culture was periodically transferred to fresh media. To compare the isolation efficiency of inner spore dust method, different isolation techniques such as entire smut ball tease and outer spore dust method was also done.

2.2 Cultural and morphological characterization

2.2.1 Cultural characterization

Ten different isolates of *U. virens* were isolated and used for further studies. The 8-mm mycelial discs were cut from the 3 weeks old culture and placed in the Petri plate containing PSA media. The plates were incubated at 27°C and three replications were maintained. The colony characters such as colony shape, colour and growth type were observed.

2.2.2 Growth on different media

In order to study the variations in mycelial growth, two *U. virens* isolates (FS 1 and FS 8) were grown in seven different media. The composition for these culture media were as follows: paddy straw extract agar (PSEA): 100 g paddy straw, 15 g agar and 1000 ml distilled water; rice extract agar (REA): 200 g rice, 15 g agar and 1000 ml distilled water; potato dextrose agar (PDA): 200 g potato, 20 g dextrose, 15 g agar and 1000 ml distilled water; potato sucrose agar (PSA): 200 g potato, 20 g sucrose, 15 g agar and 1000 ml distilled water; cornmeal agar (CMA): 200 g corn meal, 15 g agar and 1000 ml distilled water; oatmeal agar (OMA): 30 g oatmeal, 1000 ml distilled water; Czapek agar (CA): 30 g sucrose, 3 g NaNO₃, 1 g K₂HPO₄, 0.5 g MgSO₄, 0.5 g KCl, 0.01 g FeSO₄, 15 g agar, and 1000 ml distilled water. The 8 mm mycelial discs were cut from three-week-old culture and placed in the Petri plate containing different media and incubated at 27°C. Three replications were maintained. The colony diameter on different media was measured at different time interval *viz.* 7th, 14th and 21st days of incubation.

2.2.3 Morphological characterization

Yellow coloured false smut spores were spread over the cleaned glass-slide having a drop of distilled water using a needle and covered with a cover slip and observed under the compound microscope

under 40X. The mycelial character and septation were observed and photographed. The colour and shapes of spores were also observed. The morphological characters like width of mycelium and size of conidia of isolates were measured under phase contrast microscope-image analyzer.

2.2.4 Molecular characterization

2.2.4.1 DNA extraction

The total DNA of *U. virens* was extracted using the Cetyl Trimethyl Ammonium Bromide (CTAB) technique (Murray and Thompson, 1980) with partial modification. A pure culture of *U. virens* mycelia was inoculated in potato sucrose broth and cultured at 28°C in a rotary incubator shaker at 125 rpm for 2 weeks (Ladhalakshmi *et al.*, 2012). The mycelial mat was collected by filtering the broth through Whatman filter paper No.1 and dried under sterile condition. About 100 mg of the dried mycelium was powdered using liquid nitrogen, to which 750µl of CTAB buffer (100mM TrisHCl, pH 8.0, 1.4 M NaCl, 50mM EDTA, pH 8.0 and 2% CTAB) was added. The finely ground mycelium was added into a sterile 1.5 ml centrifuge tube, mixed thoroughly, and incubated for an hour in a water bath at 65°C with intermittent vortex. After incubation, equal volume of phenol: chloroform: isoamyl alcohol (24:1:1 v/v) was added and mixed gently and was centrifuged at 13000 rpm for 15 min at 4°C. The aqueous phase was transferred to a new eppendorf tube and equal volume of ice-cold iso-propanol was added and kept for overnight incubation at -20°C. After incubation, DNA was precipitated by centrifugation at 13,000 rpm for 15 min. Ethanol wash was given twice for the DNA pellet using 70% ethanol and air dried at 37°C. The pellet was suspended in 50 µl of the sterile distilled water and stored at -20°C.

2.2.4.2 Specific internal transcribed spacer (ITS) region analysis

The ten different isolates of *U. virens* were confirmed using the specific internal transcribed spacer (ITS) primers US1-5, Forward (CCGGAGGATACAACCAAAAAA ACTCT) and US3-3, Reverse (GCTCCAAGTGCGAGGATAACTGAAT) primer (Zhou *et al.*, 2003). The 20 µl of Polymerase Chain Reaction (PCR) mixture consisted of 0.5 µl of 2.5 mM of dNTPs, one unit of Taq polymerase, 10 pmol of each primer (US1-5/US3-3), and DNA template (20 ng). The PCR reaction was performed with following condition: an initial denaturation at 96°C for 2 min, and 30 cycles of amplification (denaturation at 96°C for 20 s, primer annealing at 53°C for 30 s and extension at 70°C for 30 s) and single cycle of final extension was done at 72°C for 7 min. PCR amplification was performed on a Thermocycler. The amplified products were then resolved in 1.2% agarose gel electrophoresis and viewed under a gel documentation unit (Alpha Imager 2000; Alpha Innotech Corporation, San Leandro, CA, USA). The amplified PCR products were sequenced using by adopting the Sanger dideoxy sequencing method.

2.3 Statistical analysis

The *in vitro* experiments were performed by using a Completely Randomized Design (CRD). The data generated from different experiments were analyzed statistically using statistical package of INDOSTAT and the ANOVA determined the probability for significant variation among the treatments. Phylogenetic analysis was carried out using the MEGA 11 software and dendrogram was constructed by UPGMA (Unweighted Paired-Group Method with Arithmetic Average) method.

3. RESULT AND DISCUSSION

3.1 Isolates and severity of rice false smut

The false smut infected panicles were collected from different regions of Tamil Nadu and totally 10 isolates of *U. virens* were collected. Among 10 different samples used for the study, maximum false smut disease incidence per panicle was observed in variety ADT 37 collected from Nallathur, Cuddalore district (7.8%) followed by var. Co 43 (6.2%) and Amman Ponni (5.0%) collected from Pollachi, Coimbatore district and Omalur, Salem district, respectively and minimum disease incidence was observed in the rice variety Super aman collected from Chinnampalli, Dharmapuri district (1.6%) (Table 1).

For isolation of pathogen different isolation techniques were followed such as outer spore dust method, entire smut ball tease, streaking and inner spore dust method. Out of these four methods, inner spore dust was found to be more effective for *U. virens* isolation. The pathogen was not able to isolate from outer spore dust and entire smut ball tease method. The isolation efficiency was lesser in streaking method compared to inner spore dust method. Hence, inner spores were used for isolation of *U. virens* and dusted on the PSA medium as reported by Bashyal *et al.* (2020) (Fig. 2a & 2b).

Table 1: Details of false smut samples collection and disease incidence per panicle

S. No	Name of the isolates	Place of sample collection		GPS coordinates	Variety	Disease incidence per panicle (%)
		District	Village			
1	FS 1	Coimbatore	Wetland, TNAU	11.0019°N 76.9267°E	ADT 45	4.6
2	FS 2	Coimbatore	Pollachi	10.6583°N 77.0085°E	Co 43	6.2
3	FS 3	Erode	Bhavanisagar	11.29 °N 77.80°E	ASD 16	2.5
4	FS 4	Cuddalore	Nallathur	12.2998°N 79.7506°E	ADT 37	7.8
5	FS 5	Madurai	Narasingam	9.9686°N 78.2029°E	ADT 36	1.8
6	FS 6	Thanjavur	Koilvenni	10.7824°N 79.3617°E	ASD 16	2.6
7	FS 7	Tiruppur	Thittuparai	11.1105°N 77.5902°E	BPT 5204	4.2
8	FS 8	Erode	Bhavani	11.4506 °N 77.6260 °E	Co 43	5.0
9	FS 9	Dharmapuri	Chinnampalli	12.0106°N 77.9644°E	Super aman	1.6
10	FS 10	Salem	Omalur	11.7291°N 78.0200°E	Amman ponna	5.2

3.2 Cultural and morphological characterization

3.2.1 Morphological characterization

The symptoms on paddy grains differed greatly among the false smut infected samples. During the initial stages of infection, the smut balls were orange or yellowish in colour then they gradually turned to dark olive-green colour at later stage of infection (Fig. 1a, 1b & 1c). The smut balls were divided into two halves and inner region containing different layers were examined. The innermost layer of the orangish yellow balls was white, whereas the outer two layers were yellow. The inner region of the dark olive-green balls was white, the second layer was yellow, and the outermost layer was dark green in olive. Nessa (2017) reported similar results with slight variations. The hyphae under compound microscope appeared hyaline and septate. The yellow smut spores appeared spherical to elliptical in shape, and no prominent spines were noticed under compound microscope (Fig. 1d & 1e). The width of the hyphae was varied among the isolates and ranged between 1.84 and 3.10 μm (Table 2). The maximum width of the hypha (3.10 μm) was observed in isolate FS 2 and minimum in FS 7 (1.84 μm). The size of the conidia also differed between isolates and ranged from 4.10 to 6.35 μm . The isolate FS 2 contained the smallest conidia, while isolate FS 6 contained the largest. The morphological characteristics of *U. virens* isolates have been agreed with the descriptions given by Verma and Singh (1988) and Baite *et al.* (2014).

Table 2. Hyphal and conidial characteristics of *U. virens*

Isolate	Width of the hyphae (μm)	Conidial size (μm)
FS 1	2.78	4.38-5.36
FS 2	3.10	4.10-6.10
FS 3	1.97	4.50-5.65
FS 4	2.56	4.87-6.27
FS 5	2.90	4.40-5.42
FS 6	2.15	4.78-6.35
FS 7	1.84	4.85-5.77
FS 8	2.42	4.65-5.69
FS 9	2.08	4.54-5.90
FS 10	2.65	4.75-5.81
CD @ 5%	0.40	0.36
SE(d)	0.21	0.18




Fig. 1. (a). False smut infected panicle; (b). Orange smut spore ball; (c). Dark olive-green smut ball; (d) & (e). Septate hyphae and smut spores viewed in compound microscope at 40x magnification

3.2.2 Cultural characteristics

The growth of *U. virens* was very slow which took 3-5 days for formation of tiny white colony. After 14 days, large colony developed on the culture plate. The colony was creamy white to milky white in colour with fluffy mycelium, flat and leathery. On observe side, mycelium produced yellow with olive green tinch. Some isolates formed chlamydo spores at the colony margin or the centre of the colony, which was yellowish orange and later became greenish (Fig. 2c - f). The results were in accordance with Ladhakshmi *et al.* (2012), Baite *et al.* (2014) and Lin *et al.* (2018). The maximum colony growth was observed in the isolate FS 6 (65.33 mm), followed by FS 5 (61 mm) and FS 1 (59.00 mm). The minimum colony diameter was observed in FS 8 (35.33 mm) (Table 3). The colony colour varied from creamy white to milky white on front side and yellowish green or olive green on observe side. Almost all isolates showed fluffy and raised growth except FS 5 and FS 6 which formed flat colony on PSA media. Two isolates FS 2 and FS 7 formed leathery mycelial growth with undulations (Fig 3a & 3b).




Fig. 2. Isolation of *U. virens* by inner spore dust method; (a). Sampling location of spore dust from false smut balls; (b). Growth of *U. virens* after 3 days; (c). 5 days old culture of *U. virens*; (d) & (e) 14 days old culture of *U. virens* (Front side and observe side); (f). Spores formed by isolate FS 8




Fig.3 Culture photo of ten isolates (FS 1 to FS 10) : a. front side b. observe side

3.2.3 Growth on different media

The colony diameter and density of *U. virens* were highly influenced by the media. The maximum colony diameter was observed on PSA media (FS 1- 43.67 mm & FS 8- 39 mm) and minimum on CA media (FS 1- 10 mm & FS 8- 19.67 mm) (Table 4). The similar results were reported by Rongtao Fu *et al.* (2013) and Baite *et al.* (2015). Rongtao Fu *et al.* (2013) reported that maximum growth of about 36 mm diameter was observed in PSA media and minimum growth was observed in CA media after 14days of inoculation. Baite *et al.* (2015) reported that on PSA media, the mycelial growth rate of the fungus was 2.54 mm/ day, whereas PDA showed comparatively slow growth. This may be due to the preference of

Table 3. Cultural characteristics of ten *U. virens* isolates

Isolates	Radial growth (mm)			Colony colour	Growth type
	10 th day	20 th day	30 th day		
FS 1	24.33 ^b (29.556)	46.00 ^b (42.705)	59.00 ^c (50.186)	Milky white	Fluffy and circular
FS 2	14.00 ^f (21.965)	25.67 ^g (30.438)	41.33 ^h (40.009)	Creamy white	Leathery and irregular with undulation
FS 3	20.67 ^{cd} (27.030)	31.67 ^e (34.241)	50.67 ^e (45.382)	Creamy white	Fluffy and circular
FS 4	11.33 ^f (19.646)	22.00 ^h (27.957)	45.00 ^g (42.130)	Creamy white to yellow	Fluffy and circular with undulation
FS 5	25.33 ^{ab} (30.219)	41.67 ^c (40.202)	61.00 ^b (51.355)	Creamy white to yellow	Flat and circular
FS 6	26.67 ^a (31.090)	50.33 ^a (45.191)	65.33 ^a (53.930)	Creamy white	Flat and circular
FS 7	22.00 ^c (27.968)	36.33 ^d (37.067)	51.00 ^e (45.573)	Milky white	Leathery, circular with undulation
FS 8	11.67 ^f (19.969)	26.67 ^g (31.090)	35.33 ⁱ (36.471)	Milky white with yellowish green	Fluffy and circular
FS 9	16.33 ^e (23.835)	29.00 ^f (32.581)	48.67 ^f (44.236)	Milky white	Fluffy and circular
FS 10	20.00 ^d (26.561)	33.00 ^e (35.055)	54.00 ^d (47.295)	Milky white	Fluffy and circular
CD @ 5%	1.278	1.391	1.016	-	-
SE(d)	0.613	0.666	0.487	-	-

Values are the means of three replications.

Figures in parentheses represent arcsine transformation.

Means in a column followed by same superscript letters are not significantly different according to DMRT

U. virens for sucrose over dextrose. The mycelial density was thick on PSA, PDA, OMA and CMA media, whereas on REA, PSEA and CA media, the density was very less (Fig. 4a & 4b).

Table 4. Mycelial growth of two isolates (FS 1 and FS 8) of *U. virens* on seven media

Media	Radial growth (mm)					
	FS 1			FS 8		
	7 th day	14 th day	21 st day	7 th day	14 th day	21 st day
PSA	21.00 ^a (27.236)	31.67 ^a (34.227)	43.67 ^a (41.344)	20.00 ^a (26.547)	27.33 ^a (31.510)	39.00 ^a (38.645)
PDA	16.00 ^{ab} (23.542)	28.67 ^b (32.354)	37.67 ^b (37.842)	17.67 ^a (24.853)	25.00 ^{ab} (29.997)	37.33 ^a (37.662)
OMA	14.33 ^{bc} (22.235)	24.00 ^c (29.319)	35.33 ^b (36.456)	12.67 ^b (20.826)	20.00 ^d (26.552)	29.33 ^b (32.790)
CMA	12.33 ^c (20.529)	21.33 ^d (27.488)	26.67 ^c (31.072)	11.33 ^{bc} (19.646)	22.67 ^{bc} (28.422)	30.67 ^b (33.626)
REA	8.67 ^d (17.091)	12.33 ^e (20.548)	23.67 ^d (29.090)	10.00 ^{cd} (18.421)	21.33 ^{cd} (27.503)	30.00 ^b (33.203)
PSEA	6.33 ^{de} (14.561)	10.67 ^e (19.024)	16.00 ^e (23.562)	9.67 ^{cd} (18.095)	15.33 ^e (23.043)	22.33 ^c (28.194)
CZ	4.00 ^e (11.473)	7.00 ^f (15.311)	10.00 ^f (18.387)	8.33 ^d (16.736)	13.67 ^e (21.693)	19.67 ^d (26.315)
CD @ 5%	2.293	1.766	1.769	2.144	1.703	1.482
SE(d)	1.059	0.816	0.578	0.999	0.776	0.690

Values are the means of three replications.

Figures in parentheses represent arcsine transformation.

Means in a column followed by same superscript letters are not significantly different according to DMRT




Fig. 4. Growth of FS 1(a) and FS 8(b) different media PSA: Potato Sucrose Agar; PDA: Potato Dextrose Agar; OMA: Oat Meal Agar; CMA: Corn Meal Agar; PSEA: Paddy Straw Extract Agar; REA: Rice Extract Agar; CA: Czapek Agar

3.2.4 Molecular characterization

The fungus' identity was validated by PCR using *U. virens*-specific primer, US1-5/US3-3 (Zhou *et al.*, 2003). The desired specific length of 380 bp band was observed in all 10 isolates and no band was formed in control lane (Fig. 5). The amplified DNA samples were purified and sequenced. The sequences were blasted in NCBI database which shows that all 10 isolates belong to *U. virens* with homology as shown in Table 5. The sequences were submitted to GenBank and accession numbers were obtained. The dendrogram constructed through UPGMA analysis clearly grouped the isolates into two major clusters (Fig. 6). The cluster I comprised of five isolates: FS1 (Wetland, TNAU), FS2 (Pollachi), FS9 (Chinnampalli), FS10 (Omalar) and FS3 (Bhavanisagar). The cluster II comprised of five isolates: FS4 (Nallathur), FS8 (Bhavani), FS6 (Koivenni), FS5 (Narasingham) and FS7 (Thitupparai). This shows that isolates taken from nearby location showed less variability and belonged to the same cluster, whereas those collected from locations farther away exhibited more variation and were grouped into separate clusters. Ze Tan *et al.* (2022) also reported variations in the

isolates of rice false smut pathogens collected from different locations compared to the same locations.

Table 5. Fungal identification based on NCBI-BLAST search (for ITS1-5.8s-ITS2 region)

Isolate	Accession number	Species identified as	Coverage identity (%)
FS 1	OR193799	<i>U. virens</i>	98
FS 2	OR223235	<i>U. virens</i>	99
FS 3	OR225647	<i>U. virens</i>	99
FS 4	OR225649	<i>U. virens</i>	98
FS 5	OR225650	<i>U. virens</i>	99
FS 6	OR225651	<i>U. virens</i>	98
FS 7	OR225654	<i>U. virens</i>	98
FS 8	OR225656	<i>U. virens</i>	98
FS 9	OR232535	<i>U. virens</i>	99
FS 10	OR232680	<i>U. virens</i>	99




Fig. 5. Polymerase chain reaction (PCR) amplification of internal transcribed spacer (ITS) regions of *U. virens* with specific ITS primers (US1-5/US3-3). Lane M – 100 bp DNA marker; lane 1- FS1; lane 2- FS2; lane 3- FS 3; lane 4- FS 4; lane 5- FS 5; lane 6- FS 6; lane 7- FS 7; lane 8- FS 8; lane 9- FS 9; lane 10- FS 10; lane C- control




Fig. 6. Dendrogram constructed by UPGMA (Unweighted Paired-Group Method with Arithmetic average) method

4. CONCLUSION

Ten fungal isolates of false smut infected panicles collected from different rice growing areas of Tamil Nadu were identified as *Ustilaginoidea virens* based on morphological and cultural characters. The fungal pathogen was further confirmed molecularly through PCR analysis using *U. virens* specific ITS primer (US1-5/US3-3). The dendrogram constructed by UPGMA method grouped the isolates into two clusters which showed the molecular diversity between the ten isolates. The information obtained from this study will help to understand the population structure and evolution of rice false smut fungus in Tamil Nadu.

REFERENCE

1. Baite MS, Sharma RK, Devi TP, Sharma PR, Kamil D. Morphological and molecular characterization of *Ustilaginoidea virens* isolates causing false smut of rice in India. *Indian Phytopathol.* 2014;67(3):222-7.
2. Baite MS, Sharma RK. Isolation technique and culture conditions of false smut pathogen (*Ustilaginoidea virens*) on rice. *Indian Phytopathology.* 2015;68(1):50-5.
3. Bashyal BM, Parmar P, Zaidi NW, Sunani SK, Prakash G, Aggarwal R. Improved methodology for the isolation of false smut pathogen *Ustilaginoidea virens* of rice. *Indian Phytopathology.* 2021; 74:249-52.
4. Biswas A. Field Reaction of Hybrid Rice Varieties to False Smut (FSm) and Kernel Smut (KSm) Diseases in West Bengal, India. *Environment and Ecology.* 2001 Mar;19(1):229-30.
5. Fan J, Yang J, Wang YQ, Li GB, Li Y, Huang F, Wang WM. Current understanding on *Villosiclava virens*, a unique flower-infecting fungus causing rice false smut disease. *Molecular Plant Pathology.* 2016;17(9):1321-30.
6. Fu R, Yin C, Liu Y, Ding L, Zhu J, Zheng A, Li P. The influence of nutrient and environmental factors on mycelium growth and conidium of false smut *Villosiclava virens*. *Afr. J. Microbiol. Res.* 2013 Feb 26;7: 825-33.
7. Huang S, Liu L, Wang L, Hou Y. Research on advance of rice false smut *Ustilaginoidea virens* (Cooke) takah worldwide: IV. Identification of rice resistance to RFS, management and prospection of RFS. *Journal of Agricultural Science.* 2019 Sep 15;11(15):275.
8. Ikegami H. Studies on the false smut of rice VII. Analysis on the occurrence of the smut balls at the different ages. *Res Bull Fac Agric Gifu Univ.* 1961; 14:27-37.
9. Ladhakshmi D, Laha GS, Singh R, Karthikeyan A, Mangrauthia SK, Sundaram RM, Thukkaiyannan P, Viraktamath BC. Isolation and characterization of *Ustilaginoidea virens* and survey of false smut disease of rice in India. *Phytoparasitica.* 2012 Apr; 40:171-6.
10. Lin X, Bian Y, Mou R, Cao Z, Cao Z, Zhu Z, Chen M. Isolation, identification, and characterization of *Ustilaginoidea virens* from rice false smut balls with high ustilotoxin production potential. *Journal of basic microbiology.* 2018 Aug;58(8):670-8.
11. Murray MG, Thompson W. Rapid isolation of high molecular weight plant DNA. *Nucleic acids research.* 1980 Oct 10;8(19):4321-6.
12. Nessa BO. *Rice false smut disease in Bangladesh: Epidemiology, yield loss and management* (Doctoral dissertation, PhD thesis, Department of Plant Pathology and Seed Science, Sylhet Agricultural University, Sylhet, Bangladesh).
13. Rush MC, Shahjahan AK, Jones JP, Groth DE. Outbreak of false smut of rice in Louisiana. *Plant disease.* 2000 Jan;84(1):100-.
14. Tan Z, Bai Z, Qin Y, Du J, Zhang R, Tian B, Yang Y, Yu Y, Bi C, Sun W, Fang A. Characterization of genetic diversity and variation in pathogenicity of the rice false smut pathogen *Ustilaginoidea virens* from a single source. *Plant Disease.* 2022 Oct 3;106(10):2648-55.
15. Verma RK, Singh RA. Variations in *Claviceps oryzae-sativae* the incitant of false smut of rice. *Indian Phytopathol.* 1988;41(1):48-50.
16. Zhou L, Lu S, Shan T, Wang P, Sun W, Chen Z, Wang S. Chemistry and biology of mycotoxins from rice false smut pathogen. *Mycotoxins: Properties, applications and hazards.* 2012:109-30.

17. Zhou YL, Izumitsu K, Sonoda R, Nakazaki T, Tanaka E, Tsuda M, Tanaka C. PCR-based specific detection of *Ustilagoidea virens* and *Ephelis japonica*. *Journal of Phytopathology*. 2003 Sep;151(9):513-8.
18. Zhou YL, Xie XW, Zhang F, Wang S, Liu XZ, Zhu LH, Xu JL, Gao YM, Li ZK. Detection of quantitative resistance loci associated with resistance to rice false smut (*Ustilagoidea virens*) using introgression lines. *Plant pathology*. 2014 Apr;63(2):365-72.

UNDER PEER REVIEW