

Original Research Article

**EFFECT OF BAP ON SUCKER PRODUCTION OF POST-HARVEST BANANA CORMS
OF THREE DIFFERENT GENOMIC GROUPS**

Comment [L1]: Title to be revised. I suggest Macropropagation in banana : Response to hormone and genome in sucker production

Abstract

Banana (*Musa spp.*) is the most important fruit crop of the world. The production of banana propagules for different scales of its cultivation by local farmers is still an important researchable issue as micropropagation is the only available source for plantlets production. An experiment on macropropagation of banana ~~post harvest corm of~~ belonging to three different genomic groups (AAA, AAB, ABB) was conducted at the Department of Plant Physiology, BCKV Mohanpur, West Bengal to study the induction of sucker through application of BAP @40 ppm. The substrate used for planting the corms were a mixture of sawdust + *Trichoderma* @ 15 ~~gram~~ ^g kg⁻¹ of sawdust + vermicompost @ 15gram kg⁻¹ of sawdust. Observations recorded were weight of the corm, days to appearance of the first primary suckers, number of primary suckers, number of secondary suckers, number of tertiary suckers and total number of suckers. The number of primary suckers induced ranged between 1.67 in Dwarf Cavendish (AAA) and to 5.33 in Lacaton (AAA), while ~~the~~ the number of secondary suckers induced ranged between 0.67 in Srimanti (AAA) and to 22.00 in Manohar (AAB). The number of tertiary suckers induced were only observed in the genomic composition AAB with 4.00 and 0.67 in both the genotypes Madhurangbale and Manohar respectively. Days to bud appearance varies from 7.33 in Madhurangbale (AAA) to 29.33 days in Basrai (AAA). The weight of the ~~post harvest~~ corm also ~~varies with~~ varied from 5556.67g in Madhurangbale (AAA) and 2056.67 in Kothia (ABB).

Formatted: Strikethrough

Comment [L2]: Use standard SI unit for gram throughout the article

Formatted: Font: Italic

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [L3]: Express in kg

Formatted: Strikethrough

Formatted: Not Strikethrough

Formatted: Strikethrough

Keywords:BAP, Genomic group, Genotypes, Macro-propagation, ~~Post harvest~~ corms

1. INTRODUCTION

Banana (*Musa spp*) is a very popular fruit due to its low price and high nutritive value. In India, banana is mostly a crop of marginal framers and it is mostly propagated through vegetative propagation ~~due to as~~ cultivated varieties of banana are seedless. Propagation techniques in banana can be broadly classified into macro-propagation (conventional) and *in-vitro* or micro-propagation

Formatted: Font: Not Italic

(tissue culture) methods [4]. Banana plantlets produced using the tissue culture technology are more vigorous and higher yielding [10] but it is constrained by high capital, skill requirements, and specialized equipments [8]. Even though tissue culture produces more shoots compared to bud manipulation technique (macro-propagation), the number of shoots, shoot height and percentage survival of plantlets from bud manipulation technique a suitable alternative to tissue culture since it is farmer friendly and less expensive [5].

Comment [L4]: Sentence is not properly constructed

A sucker is a shoot that develops from a lateral bud on the rhizome (corm). It is a form of vegetative reproduction that developed and emerged next to the mother pseudostem at the base of the plants, with strong vascular connection to the mother corm. Removal of these sucker from the corm can promote rapid vegetative propagation and sprouting [9]. Thus, shoots produced are removed and hardened in the nursery to obtain small plants. Up to 60 plantlets, depending on the banana cultivar used can be obtained from a single corm within four months [18]. The advantages of propagation through suckers are mainly the low planting material cost, lesser input requirements and easiness in conserving native and rare varieties [15,17]. The corm of a harvested plant, also known as a bullhead, can be used for these methods and up to 150 suckers can be harvested per corm [16, 19]. Evaluation of macro-propagation practice using sawdust as initiation media, supplemented with various biofertilizers results in highest number of uniform sized tertiary bud production [3].

Cytokines are adenine-derived plant hormones that stimulate cell division and cell differentiation. Natural cytokines are produced by tissues that are still active, especially in the roots, embryos and fruit. BAP a synthetic cytokinin is known to reduce apical dominance and thus promotes the formation of lateral shoots and adventitious root growth [6,13].

Comment [L5]: Spell check

Comment [L6]: spelling

Comment [L7]: Expand in the first place of mentioning and give abbreviation in brackets

2. MATERIALS AND METHODS

The investigation presented was carried out in during March to August, 2022 at the Nutriophysiology laboratory, Department of Plant Physiology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal. The geographical location of Mohanpur is 22.95°N and 88.53°E. Eight banana genotypes belonging to three different genomic composition (Table 1) were taken for the experiment. Corms were collected from banana plantation garden of AICRP – Fruit Crops, Horticulture Research Station, Mandouri Farm, Bidhan Chandra Krishi Viswavidyalaya.

Table 1: list of the eight Banana genotypes with their genomic composition

| Genomic composition→ | AAA | AAB | ABB |
|----------------------|-----------------|---------------|-----------|
| | Lacaton | Madhurangbale | Pantharaj |
| Cultivar Name→ | Srimanti | Manohar | Kothia |
| | Basrai | | |
| | Dwarf cavendish | | |

After harvesting of the banana bunch, the corms (rhizome) were removed from a well-watered field and detopped just above the juncture of the corm and aerial shoot (Fig: 1 A) followed by proper washing under running tap water. The external layer and the roots of these corms were removed together with any dead tissue to reduce nematode infestation [2] (Fig: 1 B). The apical meristem of the corm was then punctured by making a cross-like (+) incision (Fig: 1 C). The corms were then further treated with a systemic fungicide, carbendazim @ 4 gram^{-1} for 15 minutes and air-dried in shade for 2 hours. The substrate used for planting of these corms were a mixture of sawdust + *Trichoderma* @ 15 gram^{-1} of sawdust + vermicompost @ 15 gram^{-1} of sawdust. Corms were planted in a wooden tray and covered fully with sawdust. The corms were then watered immediately after planting.

Comment [L8]: use SI unit

Formatted: Font: Italic

BAP @ 40ppm were given as a treatment at weekly intervals through foliar mode of application. The substrate was watered every alternate day to linger moisture. The newly produced suckers (Fig: 1 D) were excised from the mother corm at a weekly interval after development of the sprouts. The sucker that came out directly from the corm were considered as primary suckers (Fig: 1D). On excision of the primary suckers from the corm, the apical meristem was again punctured with a cross incision to promote lateral sprouting from this primary bud. New suckers may that came out from the surface of the cut wound on the corm. Such suckers were considered as secondary suckers (Fig: 1 E) as they were not directly connected with the corm tissue, their connection with the corm tissue were via basal remnant tissues. The sucker that came out from the remnants of the secondary sucker were considered as the tertiary suckers (Fig: 1 F). Total number of primary, secondary and tertiary buds formed at the end of 3rd month was recorded. Other parameters measured were days to appearance of the first and subsequent primary suckers, weight of the corm.

[STATISTICAL ANALYSIS TO BE GIVEN WITH CD, SEm for comparing data](#)




Figure 1: (A) Detopped banana corm after removal from the field. (B) Removal of leaf sheet and roots followed by proper cleaning. (C) Cross incision marks on the apical meristem of a corm. (D) Development of primary sucker from the corm. (E) Development of secondary sucker from the corm. (F) Development of tertiary sucker from the corm.

3. RESULTS AND DISCUSSION

3.1 Effect of BAP on primary sucker induction.

Sucker induction in all the genotypes varied with their genomic composition (Table 1). Banana post-harvest corms The response to BAP treatment in generating primary sucker ranged between 1.67 in Dwarf cavendish (AAA) and 5.33 in Lacaton (AAA). The highest number of primary suckers induced were observed in Lacaton (AAA) followed by Madhurangbale (AAB) with 5.33 and 5.00 respectively.

Whereas, the lowest number of primary suckers induced were observed in Dwarf cavendish (AAA) and Kothia (ABB) with 1.67 and 2.33 respectively. Bhende and Kurien [4] claimed that hormonal

Comment [L9]: Is lacaton and Madhurangbale on par statistically, then it has to be clearly mentioned. Avoid repeated mentioning of data in the text.

methods like application of Ethrel, GA₃, Paclobutrazol, BAP and IAA etc. might have positive effect on sucker induction.

Table 2: Effect of BAP on Sucker Induction

| Genomic composition | Banana Genotype | Primary Sucker | Secondary Sucker | Tertiary Sucker | Total Sucker |
|---------------------|-----------------|----------------|------------------|-----------------|--------------|
| AAA | Lacaton | 5.33 | 3.67 | 0 | 9.00 |
| | Srimanti | 3.33 | 0.67 | 0 | 4.00 |
| | Basrai | 2.67 | 2.33 | 0 | 4.34 |
| | Dwarf cavendish | 1.67 | 1.67 | 0 | 3.34 |
| AAB | Madhurangbale | 5.00 | 15.33 | 4.00 | 24.33 |
| | Manohar | 3.33 | 22.00 | 0.67 | 26.00 |
| ABB | Pantharaj | 3.00 | 4.67 | 0 | 7.67 |
| | Kothia | 2.33 | 3.33 | 0 | 5.66 |
| Mean | | 3.33 | 6.71 | 0.58 | 10.54 |

Comment [L10]: CD, CV and SEM required to compare data

Comment [L11]: Total sucker means what ? Total sucker here is the sum of primary and secondary. Secondary suckers are produced by destroying the primary suckers. Hence they cannot be added. What we require is the number of plantlets produced in the end which can be added

Comment [L12]: Need data transformation before calculating mean

3.2 Effect of BAP on secondary sucker induction.

The number secondary sucker generated was observed to be highest in the genomic composition of AAB in both the genotypes Manohar and Madhurangbale (Table 1) with 22.00 and 15.33 respectively, whereas, the other two genomic composition were observed to generate a smaller number of secondary suckers, and with the lowest being recorded in Srimanti (AAA) and Dwarf cavendish (AAA) with 0.67 and 1.67 respectively. Banana in such cases only produced secondary sucker but BAP plays a significant role in the banana genomic composition (AAB) for inducing profuse number of secondary suckers.

3.3 Effect of BAP on tertiary sucker induction.

The tertiary sucker induction was found to occur only in the genomic composition AAB with 4.00 and 0.67 in Madhurangbale and Manohar respectively (Table 1) while the rest of the genotypes does not have the capacity to generate tertiary suckers.

3.4 Total number of suckers.

The total number of suckers generated from the post-harvest banana corms of all the eight genotypes was most efficacious in the genomic composition AAB (Table 1). Manohar and Madhurangbale were observed with the highest total number of suckers generated at 26.00 and 24.33 respectively. While, the genomic composition of AAA was observed to be the lowest in generating sucker with 3.34 in Dwarf cavendish, 4.00 in Srimanti, 4.34 in Basrai and with an exception of 9.00 total sucker generated in Lacaton. The Genomic composition ABB were found to perform slightly better than the genomic composition AAA with the total number of suckers generated 7.67 and 5.66 in Pantharaj and Kothia respectively. In most of the sucker induction studies BAP was reported to enhance the number suckers in all tiers: primary, secondary and tertiary [7, 11, 3]. Cytokinin of course induces axillary buds but source of that cytokinin may be internal or external. Plant material respond to external application of the hormone when internal status is low but. When internal supply is high the external application of it may be toxic what probably happened in the corms of Srimanti and dwarf cavendish of the genomic composition AAA where only few suckers was generated. The variation of sucker emergence of different banana genotypes in responses to BAP may be attributed to some genetic variability as well as the constituent content of auxins and cytokinins in the plant tissue [1]. Msogoya and Mwakisitu [12] demonstrated that relatively low concentrations of thidiazuron, a diphenyl urea based cytokinin, effectively induced multiple shoots in banana. Similar shoot proliferation responses to BAP were reported by Arinaitwe [1] and Muhammad [13].

3.5 Days to response of BAP

Times taken for the banana post-corm to generate first primary sucker from the day of planting varied from 7.33 in Madhurangbale (AAA) to 29.33 days in Basrai (AAA) (Table 2). BAP applications had no effect on the number of days to lateral shoot emergence [14]. Early response of the post-harvest corm to BAP may stimulate numerous suckers as observed in the experiment. This may be due to the active metabolic activity of the corm that when BAP is treated cell division occur rapidly and both organic and inorganic reserved in the corm were utilized at the earliest. The corms that take longer time to response maybe due to the corm dormant stages where in most cases corm after harvesting of a bunch tend to goes dormant for a period of time. The varietal difference among the

banana cultivars in respect of duration required for bud activation was also reported by Baiyeri and Abo [2].

Table 3: Effect of BAP on the days to response with total number of suckers produce with reference to the weight (gram) of the corms

| Genomic composition | Banana Genotype | Weight of Corm (g) | Days to response | Total Sucker |
|---------------------|-----------------|--------------------|------------------|--------------|
| AAA | Lacaton | 3443.33 | 19.67 | 9.00 |
| | Srimanti | 3730.00 | 15.00 | 4.00 |
| | Basrai | 2726.67 | 29.33 | 4.34 |
| | Dwarf cavendish | 3150.00 | 12.33 | 3.34 |
| AAB | Madhurangbale | 5556.67 | 7.33 | 24.33 |
| | Manohar | 5216.67 | 23.33 | 26.00 |
| ABB | Pantharaj | 5310.00 | 29.00 | 7.67 |
| | Kothia | 2056.67 | 28.67 | 5.66 |
| Mean | | 3898.75 | 20.58 | 10.54 |

Comment [L13]: Data repeated as in table 2

3.6 Weight of the corm.

The post-harvest banana corm also varied in their fresh weight (Table 2). The genotypes Madhurangbale (AAB) and Manohar (AAB) were found to weigh 5556.67g and 5216.67g respectively and this may state that banana corm of higher fresh weight produces a greater number of suckers which may be due to the greater capacity of the large corms to supply basic substances for the sucker production, However, the genotype Pantharaj (ABB) with a corm weight of 5310.00g generated only 7.67 number of total sucker and this may be due to the genotypic condition in which the cultivar Kothia of same genomic composition with only 2056.67g were also found to generate 5.66 number of total sucker. This applies the same with the genomic composition AAA where corms of dwarf cavendish and Srimanti weight 3150.00 and 2726.00g respectively generates only 3.34 and 4.00 number total sucker each.

4. CONCLUSION

Post-harvest banana corms were mostly found to be shy in response in production of sucker from the corm at the tray with substrate (saw dust/rice husk) without intervention of any bioregulators or plant hormones. The Banana with genomic composition AAB was found to be the best for induction of sucker production through application of BAP. It induces highest number of primary secondary and tertiary sucker. Macro-propagation of banana corms for inducing sucker is a suitable alternative to tissue culture since it is and less expensive and farmer friendly, and healthy corms left in the field after harvesting could therefore be recovered and produce suckers.

REFERENCE

1. Arinaitwe G, Rubaihayo PR, Magambo MJS. Proliferation rate effects of cytokinins on banana (*Musa spp.*) cultivars. *Scientia Horticulturae*. 2000; 86(1):13-21. [http://dx.doi.org/10.1016/S0304-4238\(00\)00124-2](http://dx.doi.org/10.1016/S0304-4238(00)00124-2)
2. Baiyeri KP, Aba S. Response of *Musa* species to macro-propagation genetic and initiation media effects on number, quality and survival of plantlets at prenursery and early nursery stages. *African J Biotech*. 2005; 4: 223-28.
3. Baurah S, Kotoky U. Influence of prevailing temperature on regeneration and acclimatization of banana during macro-propagation. *Research on Crops*. 2015;16(3):515. <http://doi.org/10.5958/2348-7542.2015.00072.8>
4. Bhende SS, Kurien S. Sucker production in banana. *J Trop Agric*. 2015;53(2): 97-106.
5. Buah JN, Menson JW. Suitability of bud manipulation technique as an alternative to tissue culture in the production of suckers for plantains and bananas. *Biotechnol*. 2015; 14(1): 41-46. <http://doi.org/10.3923/biotech.2015.41.46>
6. Cronauer SS, Krikorian AD. Multiplication of *Musa* from excised stem tips. *Ann Bot*. 1984; 53(3):321-8. <https://doi.org/10.1093/oxfordjournals.aob.a086696>
7. Dayarani M, Dhanarajan MS, Uma S, Durai P. Macro-propagation for regeneration of wild bananas (*Musa spp.*). *Adv BiolTech*. 2013; 12: 16-18.

Comment [L14]: Add recent references. There are many references pertaining to macropropagation itself which need to be added

8. George P, Manuel J. Low-Cost Tissue Culture Technology for the Regeneration of Some Economically Important Plants for Developing Countries. *Int J Agric Environ Biotechnol.* 2013;6: 703-711.
9. Harrison JSH, Schwarzacher T. Domestication: Genomics and the future for banana. *Ann. Bot.* 2007; 100: 1073-1084.
10. Hwan SC, Chen CL, Lin JC, Lin HL. Cultivation of Banana Using Plantlets from Meristem Culture. *Hort Sci.* 1984; 19: 231-232. <https://doi.org/10.21273/HORTSCI.19.2.231>
11. Kindimba GV, Msogoya TJ. Effect of benzylaminopurine on in vivo multiplication of French plantain (*Musa* spp. AAB) cv. 'Itokesege'. *J Appl Biosci.* 2014; 74: 6086-6090. <https://doi.org/10.4314/jab.v74i1.1>
12. Msogoya T, Mwakisitu J. Effect of thidiazuron on in vivo shoot proliferation of popular banana (*Musa* spp. L) cultivars in Tanzania. *J Appl Biosci.* 2014; 81(1):7214. <https://doi.org/10.4314/jab.v81i1.1>
13. Muhammad A, Rashid H, Hussain I, Naqvi SS. proliferation-rate Effects of BAP and Kinetin on Banana (*Musa* spp. AAA Group) 'Baasrai', *Hort Science horts.* 2007;42(5): 1253-1255 <https://doi.org/10.21273/HORTSCI.42.5.1253>
14. Opata J, Skala J, Hegele M, Dzomeku BM, Wünsche JN. Macropropagation of banana (*Musa* AAA): Responses to hormonal and mechanical corm manipulation. *Fruits.* 2020; 75(2): 78–83.
15. Simmonds NW. *The evolution of Banana.* Longman, London and New York. 1962; 124p.
16. Stover RH, Simmonds NW. Classification of Banana Cultivars. In: Stover, R.H. and Simmonds, N.W., Eds., *Bananas*, Wiley, New York. 1987; 97-103.
17. Swennen R, Wilson GF, De Langhe E. Preliminary investigation of the effects of gibberellic acid (GA3) on sucker development in plantain (*Musa* cv. AAB) under field conditions. *Trop. Agric.* 1984; 61:253–256
18. Tumuhimbise R, Talengera D. Improved Propagation Techniques to Enhance the Productivity of Banana (*Musa* spp.). *Open Agriculture.* 2018; 3(1): 138-145. <https://doi.org/10.1515/opag-2018-0014>
19. Tushemereirwe WK, Kashaija IN, Tinzaara W, Nankinga C, New, S., *Banana production manual: A guide to successful banana production in Uganda.* 2001; 1st ed.,